

# Uses and management of poultry litter

N.S. BOLAN<sup>1,2\*</sup>, A.A. SZOGI<sup>3</sup>, T. CHUASAVATHI<sup>1,2</sup>, B. SESHADRI<sup>1,2</sup>,  
M.J. ROTHROCK JR.<sup>3</sup> and P. PANNEERSELVAM<sup>4</sup>

<sup>1</sup>Centre for Environmental Risk Assessment and Remediation, University of South Australia, SA – 5095, Australia; <sup>2</sup>Cooperative Research Centre for Contaminants Assessment and Remediation of the Environment (CRC CARE), University of South Australia, SA-5095; <sup>3</sup>USDA-ARS, Coastal Plains Soil, Water and Plant Research Center, 2611 W. Lucas St., Florence, SC, 29501, USA; <sup>4</sup>Indian Institute of Horticultural Research, Bangalore – 560 089, India

\*Corresponding author: nanthi.bolan@unisa.edu.au

---

The poultry industry is one of the largest and fastest growing agro-based industries in the world. This can be attributed to an increasing demand for poultry meat and egg products. However, a major problem facing the poultry industry is the large-scale accumulation of wastes including manure and litter which may pose disposal and pollution problems unless environmentally and economically sustainable management technologies are evolved. Most of the litter produced by the poultry industry is currently applied to agricultural land as a source of nutrients and soil amendment. However environmental pollution, resulting from nutrient and contaminant leaching can occur when poultry litter is applied under soil and climatic conditions that do not favour agronomic utilisation of the manure-borne nutrients. This review examines the composition of poultry litter in relation to nutrient content and environmental contaminants, its value as a nutrient source, soil amendment, animal feed and fuel source, and cost-effective innovative technologies for improving its value. Poultry litter provides a major source of nitrogen, phosphorus and trace elements for crop production and is effective in improving physical and biological fertility, indicating that land application remains as the main option for the utilisation of this valuable resource. The alternative use of poultry litter; as an animal feed and fuel source, is limited by contaminants, and high moisture content, respectively. The review proposes best management practices to mitigate environmental consequences associated with air and water quality parameters that are impacted by land application in order to maintain the continued productivity, profitability, and sustainability of the poultry industry.

---

**Keywords:** poultry litter; nutrients; trace elements; antibiotics; feed management; land application

## **Introduction**

The poultry industry is one of the largest and fastest growing agro-based industries in the world. There is an increasing demand for poultry meat mainly due to its acceptance by most societies and its relatively low cholesterol content. The poultry industry is currently facing a number of environmental problems. One of the major problems is the accumulation of large amount of wastes, especially manure and litter, generated by intensive production. is voided by a layer as po Large-scale accumulation of these wastes may pose disposal and pollution problems unless environmentally and economically sustainable management technologies are evolved (Power and Dick, 2000; Kelleher *et al.*, 2002; Sharpley *et al.*, 2007).

Most of the manure and litter produced by the poultry industry is currently applied to agricultural land. When managed correctly, land application is a viable way to recycle the nutrients such as nitrogen (N), phosphorus (P) and potassium (K) in manure. However, pollution and nuisance problems can occur when manure is applied under environmental conditions that do not favour agronomic utilisation of the manure-borne nutrients (Sharpley *et al.*, 1998; Casey *et al.*, 2006; Kaiser *et al.*, 2009). The continued productivity, profitability, and sustainability of the poultry industry will likely be dependent on the formulation of best management practices to mitigate environmental consequences associated with air and water quality parameters that are impacted by land application, and the development of cost-effective innovative technologies that provide alternative to land application of poultry wastes (Kelleher *et al.*, 2002; Moore Jr., *et al.*, 2006; Szogi and Vanotti, 2009).

Confined animal production (*i.e.* beef and dairy cattle, poultry and swine) is the major source of manure by-products in many countries, including the United States of America (U.S.), Australia, and New Zealand. For example, total 2008 broiler production in the U. S. accounted for approximately 8,882 million birds (USDA, 2008). It is estimated that about 44.4 million tons of poultry manure was produced in that year, containing 2.2 million tons of N, 0.7 million tons of P and 1.4 million tons of K (McDonald *et al.*, 2009). Similarly, on the basis of 2008 broiler production (approximately 81 million birds), it is estimated that approximately 591, 300 tons of broiler manure waste is generated annually in New Zealand, which can supply around 15, 000, 6, 000 and 9, 000 tons of N, P and K nutrients, respectively (PIANZ, 2006).

Manure by-products have the potential for being recycled on agricultural land. Beneficial use through land application is based on their ability to favourably alter soil properties, such as plant nutrient availability, soil reaction (pH), organic matter content, cation exchange capacity, water holding capacity, and soil tilth. Poultry waste contains all essential nutrients including micronutrients and it has been well documented that it provides a valuable source of plant nutrients (Kelley *et al.*, 1996; Williams *et al.*, 1999; Chan *et al.*, 2008; Harmel *et al.*, 2009), especially for organic growers (Preusch *et al.*, 2002). Addition of poultry manure to soils not only helps to overcome the disposal problems but also enhances the physical, chemical and biological fertility of soils (Friend *et al.*, 2006; McGrath *et al.*, 2009). For example, continuous cultivation of arable soils often results in the deterioration of soil structure leading to reduced crop yield. Addition of poultry manure has been shown to improve the fertility of the cultivated soil by increasing the organic matter content, water holding capacity, oxygen diffusion rate and the aggregate stability of the soils (Mahimairaja *et al.*, 1995; Adeli *et al.*, 2009).

Optimum use of manure by-products requires knowledge of their composition not only in relation to beneficial uses but also to environmental implications. Environmental concerns associated with the land application of manure by-products from intensive animal operations include leaching losses of N in sub-surface drainage and to

groundwater, contamination of surface water with soluble and particulate P, reduced air quality by emission of greenhouse gases and volatile organic compounds, and increased metals input (Williams *et al.*, 1999; Ribaudo *et al.*, 2003; Harmel *et al.*, 2004; Casey *et al.*, 2006). Maintaining the quality of the environment is a major consideration when developing management practices to effectively use manure by-products as a nutrient resource and soil conditioner in agricultural and horticultural production systems (Sims and Wolf 1994; Moore *et al.*, 1995; Moore Jr. *et al.*, 2006). Most of the environmental problems associated with improper practices of land application of manure by-products have centred on the contamination of ground and/or surface water with two major nutrients, N and P (Sims *et al.*, 2005). However, manure by-products may also contain other potentially toxic trace elements, such as arsenic (As), copper (Cu) and zinc (Zn), which, to date, have received less attention (Bolan *et al.*, 1992; Jackson *et al.*, 2003; Epstein and Moss, 2006; Toor and Hunger, 2009). For example, poultry manure addition is considered to be one of the major sources of As input to soils. In the Delaware-Maryland-Virginia (Delmarva) peninsula along the Eastern shore of the US, 20–50 tons of Arsenic (As) is introduced annually to the environment through the use of As compounds (*e.g.* Roxarsone) to control coccidiosis in poultry birds in the U.S. (Christen, 2001). However, it is noteworthy that As compounds are not commonly used to control coccidiosis in most countries including Australia and New Zealand. To offset the environmental risks of manure land application, Edwards and Someshwar (2000) pointed out that ‘to reduce the risk of offsite contamination, land application guidelines should be developed that consider the total composition of the manure by-products rather than only one component, *i.e.*, N and/or P concentration’. On the other hand, the concentration of trace elements in poultry litter and its by-products could be minimized by controlling the quality of raw feed materials and reducing mineral additives in poultry diet (Van Ryssen, 2008).

Air quality has become a major environmental concern of the poultry industry. Dust, odours and bio-aerosols (*e.g.* microbes, endotoxins and mycotoxins suspended in air) generated at production, manure storage facilities and during land spreading of poultry litter constitute the most frequent source of complaints against animal-based industries (Millner, 2009). Uncontrolled decomposition of manure produces odorous gases, including amines, amides, mercaptans, sulphides, and disulphides. These noxious gases can cause respiratory diseases in animals and humans (Schiffman and Williams, 2005). Ammonia volatilisation from manure creates odour problems, and it may also contribute to atmospheric deposition and acid rain (Walker *et al.*, 2000a; 2000b). Furthermore, greenhouse gases, such as carbon dioxide, methane and nitrous oxides are also released from manure handling and storage facilities, which are implicated in ozone depletion and global warming. Improved manure handling and storage methods are needed to reduce the emission of these gases (Aneja *et al.*, 2006).

This review examines the composition of poultry litter in relation to nutrient content and environmental contaminants, its value as a nutrient source and soil amendment to improve soil fertility, alternative uses of poultry litter including animal feed and fuel source, and cost-effective innovative technologies for improving the beneficial value of poultry litter. Poultry litter provides a major source of N, P and trace elements for crop production and is very effective in improving the physical, chemical and biological fertility, indicating that land application remains as the main option for the utilisation of this valuable resource. The review proposes best management practices to mitigate environmental consequences associated with air and water quality parameters that are impacted by land application in order to maintain the continued productivity, profitability, and sustainability of the poultry industry.

## **Poultry litter production**

The quantity of poultry litter produced in a broiler unit depends on the litter (*i.e.* bedding material) management, and feed intake and its digestibility. A range of materials including wood shavings, cereal straw, husk and paper clippings are used as bedding materials (Swain and Sundaram, 2000). Three common practices are adopted for litter management in broiler units (Bernhart *et al.*, 2010). These include single use litter, partial re-use and multi-use litter. The single-use litter involves the total clean-out of the house after each flock and replacement of the bedding material. Partial re-use involves the removal of litter from the brooding section for spreading on the grower section of the house. New bedding material is then spread on the brooding section. The partially spent litter is often composted for a few days to elevate its temperature to kill pathogens. Some of the spent litter may be removed after each batch, and after 2 to 5 batches the house is totally cleaned out. With the multi-use of litter, only caked material is removed (Sistani *et al.*, 2003) and the house is disinfected. The litter in the brooding section is either left untouched or covered with 25 to 50 mm of fresh bedding material. The multi-flock litter may increase the incidence of pathogenic microbes and parasites, and produces a spent litter with a much higher concentration of nutrients (Kelley *et al.*, 1996).

The amount of total solids (dry matter) excreted by the birds can be estimated from the dry matter digestibility of the diet. Broiler chickens generally digest about 85 to 90% of the dry matter of the feed (NRC, 1994). Broiler chickens consume approximately 2.5 to 3.0 kg of dry matter up to 35 days of age (typical first thin out) and 5 to 6 kg of dry matter up to 49 days of age (typical final clean-out). On the basis of the dry matter digestibility of the diet (87.5%), it is estimated that 0.34 and 0.63 kg of solid is excreted by a 35 and 49 day old bird, respectively (FSA, 2007). At a moisture content of 90%, total manure production will be around 4 and 6 kg for 35 and 49 days old birds, respectively. The amount of any nutrient excreted can be calculated from the difference between the amount in the feed and the amount assimilated by the bird. It has been estimated that broiler chickens excrete approximately 55% of the total N, 70% of the P and 80% of the K. The amount of feed spilt during feeding can significantly affect the total amount of solid and nutrients remaining in the litter (Leytem *et al.*, 2007).

The average daily fresh manure production for broilers is about 43 kg/1000 kg live weight (ASABE, 2005). Converting this to the quantity of dry manure removed from a typical broiler house, the amount is 6.9 kg/1000 kg live weight/day for broilers. Furthermore, handling and storage factors also affect the actual quantity and quality of manure/litter generated from various types of poultry units (Malone, 1992; Maguire *et al.*, 2006). Among these are feed composition and efficiency of feed utilisation, the type of bedding material, the frequency of crust removal and total clean out operations, the number of flocks in a house between replacement of the bedding material, the final live weight of poultry and management practices. On a dry weight basis, poultry litter production ranges from 0.7 to 2.0 tons/1000 broilers/flock.

## **Composition of poultry litter**

The major components of poultry litter include the bedding material, feather, manure and the spilt feed (Kelley *et al.*, 1996; Tasistro *et al.*, 2004). The litter contains plant nutrients, such as N, P and K, trace elements, such as Cu, Zn and As, pesticide residues, pharmaceuticals such as coccidiostats, endocrine disruptors and microorganisms. As with other organic wastes, the moisture content, pH, soluble salt level, and elemental composition of poultry manure and litter have been shown to vary widely as a function of

types of poultry, diet and dietary supplements, litter type, and handling and storage operations.

#### PLANT NUTRIENTS

The major plant nutrients in poultry manure include N, P, K, calcium (Ca), magnesium (Mg) and sulphur (S) (*Table 1*). Some general observations on chemical composition of poultry manure and litter include (Edwards and Daniel, 1992; Sims and Wolf, 1994; Toor *et al.*, 2006; Guo and Song, 2009): (i) the huge variability in nutrient concentration can be attributed to a number of factors including feed use efficiency, bedding material used, litter management practices etc; (ii) the total N and P contents of poultry manures and litters are among the highest of organic amendments including manures and compost; (iii) the total N and P contents are usually lower for poultry litter than for fresh manure, which is attributed to both the losses that occur following manure excretion and the dilution effect from combining manures with bedding materials that are low in nutrients; (iv) uric acid and ammonium are the significant N components of poultry manures and litters; (v) the use of poultry manure as a soil amendment for agricultural crops will provide appreciable quantities of essential major plant nutrients (N, P and K), secondary nutrients (Ca, Mg and S) and some of the trace elements (Cu, Zn and Mo); and (vi) application of poultry manure based on crop N requirements is likely to provide more of other nutrients (especially P) than is required by the crops.

**Table 1 Nutrient contents of manures and composts (g/kg dry weight basis).**

Nutrients	Poultry manure compost		Pig manure	Mushroom	Biosolid
	Layer	Broiler	compost	compost	compost
Nitrogen	32.8	25.7	15.2	17.5	21.8
Phosphorus	10.8	6.7	6.5	7.5	10.1
Potassium	15.2	10.1	8.2	9.2	5.9
Calcium	18.5	16.2	4.2	21.5	19.7
Magnesium	6.2	3.5	3.7	5.2	4.2
Sulphur	8.5	5.2	3.4	3.5	7.0

Among these nutrients, N and P cause some environmental concerns. Four forms of N are identified in poultry litter that include complex organic N, labile organic N, ammonium and nitrate (Sims and Wolf, 1994; Sharpley and Smith, 1995; Diaz *et al.*, 2008). Complex forms of organic N in poultry litter include constituents of feathers, spilt and undigested feed, and bedding materials. Labile organic N is largely uric acid and urea. Uric acid in the fresh manure is rapidly hydrolyzed to urea by the uricase enzyme, and the urea is subsequently hydrolyzed to ammonium by urease enzyme. Nitrate is formed when the ammonium ions are oxidised during aerobic composting.

Phosphorus in poultry litter is about two thirds present as solid-phase organic P and one third as inorganic P (Edwards and Daniel, 1992; Sharpley and Smith, 1995; Sharpley *et al.*, 2004). The amount of total P in poultry litter varies with the diet and bedding material, and ranges from 0.3 to 2.4% of dry matter. Fractionation studies have shown that a large proportion of P in poultry litter is in acid soluble fraction, indicating low bioavailability (Mahimairaja *et al.*, 1995b). According to Turner and Leytem (2004), acid extractable P in raw broiler litter is dominated by inorganic (35 to 41%) and organic P forms (58 to 65%). Inorganic phosphate species in poultry manure include dibasic calcium phosphate, amorphous calcium phosphate and weakly bound water-soluble

phosphates (Sato *et al.*, 2005), while organic P in poultry litter is largely in form of phytic acid salts (Turner and Leytem, 2004).

The amounts of major nutrients in the litter produced for a broiler unit with 100 000 birds and 6.5 flocks per annum are calculated based on the nutrient concentration values given in literature (Table 2). The fertilizer equivalents (in the form of common fertilizers) of the major nutrients in the poultry litter and the total maximum area of maize crop than can be grown using these individual nutrients are also given in the table. It is important to point out that both the total amount of nutrients in the poultry litter and the maximum area of maize crop than can be grown using these nutrients depend on the nutrient concentration of the litter. For example, if we use a N content of 35 kg/ton in the litter, the total amount of N in the poultry litter generated in this broiler unit will be around 234 500 kg (6700 x 35), which will be sufficient enough to grow a maximum of 1172 hectares of maize at the N application rate of 200 kg N/ha.

**Table 2 Nutrient concentration and the estimated total amount of nutrients in the poultry litter for poultry unit with 100, 000 birds.**

Nutrients	Concentration (kg/ton)	Total amount (tons)*	Fertilizer equivalent (tons)**	Area of maize (ha)***
Nitrogen	25.7	172	Urea (374)	860
Phosphorus	6.7	45	SSP (473)	1125
Potassium	10.1	68	KCl (136)	453
Calcium	16.2	109	Lime (272)	5450
Magnesium	3.5	23	Dolomite (121)	1150
Sulphur	5.2	35	Gypsum (194)	1167

\* calculated for 6700 tons of litter

\*\* Urea (46% N); SSP – Single superphosphate (9.5% P); KCl – Potassium chloride (50% K); Lime (40% Ca); Dolomite (19% Mg); Gypsum (18% S)

\*\*\*For maize 200 kg N/ha; 40 kg P/ha; 150 kg K/ha; 30 kg S/ha; 20 kg Ca/ha; 20 kg Mg/ha

#### TRACE ELEMENTS

Trace elements in poultry litter are mostly derived directly from the birds' diet and indirectly during manure collection. A number of trace elements are added to poultry feed not only as essential nutrients but also as supplement to improve health and feed efficiency (Table 3). Essential nutrient elements participate in a wide range of enzymatic processes involving many aspects of physiological functions and the intermediary metabolism of the organism. They act as catalyst or cofactor in enzyme systems with their role ranging from relatively weak, nonspecific ion effects (metal ion activated enzymes) to highly specific association (metallo-enzymes) in which the metal is firmly attached to the protein in a fixed number of atoms per molecule. These trace elements cannot be removed without the loss of activity and basically cannot be substituted by another element (NRC, 1994).

**Table 3 Trace element concentrations in poultry manure by-products and poultry feed (Bolan *et al.*, 2004).**

No*	Concentration (mg/kg)											
	As	B	Cd	Co	Cr	Cu	Mn	Mo	Ni	Pb	Se	Zn
1						400	1800					2300
2						313	246					327
3				313			246					327
4	34.6		4.93		9.9	6.1	501		2.46	0	1.23	743

Table 3 Continued

No*	Concentration (mg/kg)											
	As	B	Cd	Co	Cr	Cu	Mn	Mo	Ni	Pb	Se	Zn
5	0.57		-	2.0	6	30.7	166	5.0		-	0.38	158
6	9.01		0.42		17.1	96.8			5.4	3.62		378
7	0.46		1.06		4.57	64.8			7.1	8.37		459
8	43	51	3	6	-	748	956	6	15	11		718
9		19	2	8	6	19	271		14	13		252
10			0.48		7.3	54.3	465	7.69	7.0	2.3		550
11		390										
12	<1.0		0.39		0.76	23			2.6			153
13	1.27		1.66		1.81	887			4.4	10.5		6980

\*1. Poultry dropping; 2 & 3. Broiler litter; 4. Poultry litter; 5. Broiler litter; 6. Dried poultry waste from caged hens without any litter; 7. Broiler/turkey litter; 8. Layer manure; 9. Poultry litter; 10. Deep-pit poultry litter; 11. Poultry manure; 12. Layer feed; 13. Broiler feed.

Because of the intensity of production, a number of feed ingredients, including trace elements such as As, Co, Cu, Fe, I, Mn, Se, and Zn, are used in poultry industry to prevent deficiencies and diseases, improve weight gains and feed conversion, and increase egg production in the case of poultry (Miller *et al.*, 1991; Tufft and Nockels, 1991; Sims and Wolf, 1994; Moore *et al.*, 1995; Powers and Angel, 2008; Burel and Valat, 2009). Efficient feed nutrient utilisation using growth promoters is a major means of reducing potential contaminants, such as N and P in poultry and swine manure (Jongbloed and Lenis, 1998). However, since some of the growth promoters contain metals, this practice is likely to result in elevated concentration of these elements in manure by-products (Nahm, 2002). For example, Cu compounds, such as copper sulphate ( $\text{CuSO}_4$ ) and copper hydroxide [ $\text{Cu}(\text{OH})_2$ ] have been (and remain so in some regions) commonly used growth promoters in swine and poultry production units (Poulsen, 1998).

Since a major portion of the trace elements ingested is excreted in faeces and urine, the concentrations of trace elements in manure by-products depend primarily on their concentrations in the diet (Krishnamachari, 1987; Miller *et al.*, 1991; Van Ryssen, 2008). Kunkle *et al.* (1981) noticed that Cu concentrations in poultry manure by-products were linearly related to Cu added in the diet and were typically concentrated 3.25 times. Similarly, Mohanna and Nys (1999) noticed that by reducing dietary Zn from 190 to 65 mg/kg in broiler poultry feed, the body retention of Zn was increased by 8 to 20% with a consequent decrease of Zn concentration in manure by 75%. Introducing highly viscous raw materials such as triticale, rye and barley at high levels in poultry diets has been shown to reduce Zn retention, thereby contributing to increased level of Zn in manure (Mohanna *et al.*, 1999).

Trace elements used in animal health remedies also eventually reach the manure. Addition of As to feed as an additive to control coccidiosis in poultry has resulted in a seven-fold increase in As level in the litter (Morrison, 1969). Jackson and Bertsch (2001) observed that more than 90% of As in poultry manure was in water-soluble form. They have identified six As species in poultry litter and litter amended soils: the original As chemicals, Roxarsone and p-arsanilic acid, used as the feed additive, and their four metabolites - arsenate, arsenite, dimethyl arsenic acid and monomethyl arsenic acid. The original As chemicals were identified as the major species indicating that they were mostly excreted chemically unaltered. Christen (2001) obtained a direct correlation between water-extractable As in soils and the amount of poultry litter applied, implicating this material as a major source of As input in soils. As indicated earlier it

is important to point out that while As is used widely in the US but it is not a common coccidiostat in most other countries.

While the type of bedding material in broiler units may influence the litter dry matter and other chemical properties, it has little effect on the metal concentration of the resulting litter. Typically, for broilers approximately 30% of the litter is bedding materials, which has trace element concentration at background levels for plant material (Nicholson *et al.*, 1999). However, recycled paper treated with boric acid has received substantial attention as a bedding material in broiler houses (Worley *et al.*, 1999; Grimes *et al.*, 2002). With this bedding material, accumulation of boron (B) can occur since Wilkinson (1997) observed that the B content in the manure by-products collected from these houses could reach as high as 290-390 mg/kg, compared to 8-15 mg/kg in conventional broiler units which use wood chips as bedding material.

The concentration of trace elements in manure can vary considerably depending on the age of animal, type of ration, housing type and waste management practice. Although such variation is normal, it makes quality assurance difficult, rendering it cumbersome to use manures as an alternative to chemical fertilizers (Eck and Stewart, 1995). The variation in chemical composition also causes uncertainties in predicting the reactions and bioavailability of manure-borne nutrients and metals in soils.

Manure addition is increasingly being recognised as a major source of both essential and non-essential element input to soils, with repeated applications having resulted in elevated concentrations of metals in soil (He *et al.*, 2009). Based on the dry matter content and elemental concentration, Nicholson *et al.* (1999) estimated the typical metal loadings from various manure sources at an application level of 250 kg/ha (Table 4). Significant quantities of Cu and Zn were added through swine (2.2 and 1.6 kg/ha, respectively), poultry (1.1 and 0.5 kg/ha) and cattle (1.0 and 0.3 kg/ha) manures. Since Cu and Zn are essential for plant growth, optimum level of poultry litter application is likely to meet the crop requirements of these nutrients. Application of cattle and poultry manures represented also an important source of Cd metal to arable soils (ca. 3-7 g/ha), although most of the Cd is typically added via atmospheric deposition and phosphate fertilizer use.

**Table 4 Estimated trace element loading (kg/ha) from 'typical' manure application at the 250 kg N/ha rate (Nicholson *et al.*, 1999).**

Element	Loading (kg/ha)					
	Cattle compost	Swine compost	Dairy slurry	Swine slurry	Broiler litter	Layer compost
As	0.01	0.01	0.01	0.01	<0.01	<0.01
Cd	0.003	0.002	0.002	0.001	0.003	0.007
Cr	0.02	0.02	0.03	0.02	0.01	0.03
Cu	0.2	1.5	0.3	1.7	0.2	0.5
Ni	0.03	0.05	0.03	0.05	0.02	0.05
Pb	0.03	0.03	0.04	0.03	0.02	0.05
Zn	0.7	2.1	0.9	2.3	1.1	2.9

Land application of manure by-products is traditionally based on N and/or P loading. Hence the level of trace element loading through manure application depends on the concentrations of both the trace element and the two major nutrients, N and P. Metals, such as Cu and Zn are strongly bound to organic matter and the land application of organic matter-rich poultry manure is likely to result in the accumulation of such metals

in soils (Schomberg *et al.*, 2009). Over-supplementation of Cu and Zn do not seem to have direct environmental effects but are potentially phytotoxic (Alva *et al.*, 2000; Novak *et al.*, 2008). The environmental risk of heavy metals such as Cu and Zn is largely dependent upon the ability of the soil to adsorb these elements and the potential for leaching or loss by soil erosion (Gupta and Charles, 1999).

The United States Environmental Protection Agency (US-EPA) regulations set ceiling concentrations of each element that may not be exceeded if the sewage sludge is to be land applied (Table 5). The US-EPA standards do not recognize differences in soil type and plant species, and their ability to retain and precipitate these elements and bioaccumulation potential (Ritter, 2000). The typical loadings of trace elements in the USA through various sources are also given in Table 5. As expected, except for As, sewage sludge would add more metals than commercial fertilizer and manure products. Poultry manure would add the highest amount to As, manure products on the average, would add about 1.6-13, 50-107, and 212-4490 times more As, Zn and Cu, respectively than the commercial fertilizers, for a particular application rate of P. However, the poultry litters generated in most other countries are unlikely to add such high levels of As and Cu because these metals are not added in high concentration in the feed.

**Table 5 US-EPA Part 503 land application guidelines for trace elements loading (kg/ha) from commercial fertilizer, sewage sludge and from various manure sources.**

Element	Permitted	Fertilizer	Sewage sludge	Dairy manure	Poultry manure	Swine manure
As	2	0.003	0.043	0.006	0.054	
Cd	1.9	0.001	0.311	0.001	0.008	0.0003
Cr	150		4.310		0.016	0.044
Cu	75	0.0004	4.025	0.350	0.086	1.796
Pb	15		5.479		0.004	0.019
Hg	0.85		0.025			
Mo	0.90	0.003	0.042	0.013		
Ni	21				0.004	0.017
Se	5		0.009		0.002	
Zn	140	0.018	6.376	0.890	1.167	1.933

\*Based on 100 kg P<sub>2</sub>O<sub>5</sub>/ha

Fertilizer (McBride and Spiers, 2001; grade 6-24-24); sewage (Caper *et al.*, 1978; P = 1.46%); dairy (McBride and Spiers, 2001; P = 0.803% for dairy); poultry (Jackson and Miller, 2000; P = 2.78%); swine (De Abreu and Berton, 1996; P = 3.25%)

It is important to recognise that trace element loadings through manure application may overestimate their actual net accumulation in soil, since a substantial portion of the trace elements in manure originates in crop uptake or soil contaminating the manure, and are therefore being recycled within a production system. Conversely metals in fertilizers, imported feedstuff or sewage sludge represent *de novo* additions to the soil (McBride and Spiers, 2001). In general, the behaviour and effects of metals added to soil in the form of animal manure are essentially similar to that applied in the form of sewage sludge. It is therefore prudent to consider adopting similar guidelines for manure application as have been described for sewage sludge.

#### ANTIBIOTICS, COCCIDIOSTATS AND PESTICIDES

Antibiotics used in poultry production system include bacitracin, bambarmycin, chlortetracycline, dihydrostreptomycin, erythromycin, lincomycin, neomycin, oxytetracycline, penicillin, spectinomycin, streptomycin, tetracycline and tylosin

(Bhattacharya and Taylor, 1975). Similarly several chemicals are used to control internal protozoan parasites that cause coccidiosis (Sims and Wolf, 1994). The excretion of the coccidiostats sulphaquinoxaline and decoquinate has been studied in caged broilers fed diets containing these two drugs (Hobson-Frohock and Johnson, 2006). Storage of broiler excreta at 23°C for 9 days showed contrasting effects; the sulphaquinoxaline content decreased about 40% with respect to initial content in the manure whereas the decoquinate concentration remained unchanged. Both coccidiostats appeared relatively stable when samples of excreta were dried in 1.0 cm thick layers in an oven at 200°C.

In layer operations, chemicals are often included in diets to control insects in the litter. Examples of some larvicides include rabon, zoalene, unistat, nicarbazin, furazolidone, and nitrofurazone and cyromazine (Bhattacharya and Taylor, 1975; Axtell, 1986; Sims and Wolf, 1994; OSU, 2010). Some of these chemical residues have been measured in poultry manure wastes (*Table 6*). The concentration of chemical residue found in poultry litter is related to the amount, frequency, retention and stability of the chemical, and the composting of the litter (Nahm, 2005; Karci and Balcioglu, 2009).

**Table 6 Commonly used antibiotics, coccidiostats and larvicides in poultry production (Sims and Wolf, 1994).**

Material	Name		Concentration (mg/kg)
	Chemical	Common	
Antibiotic	Amprolium	Amprol	0.0-77.0
	Chlortetracycline	Aureomycin	0.8-26.3
	Chlortetracycline	Aureomycin	0.1-2.8
	Neomycin sulphate	Neomycin	-
	Nicarbazin		35.1-152.1
	Oxytetracycline	Terramycin	5.5-29.1
	Penicillin	Propen	0-25
Coccidiostat	Amprolium	Amprol	-
	Zoalene	-	-
Larvicide	2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	Rabon	196-580

The consequences of land application of poultry manure containing these chemical residues have not been adequately evaluated. Decreases in the rate of decomposition of the poultry litter and the crop responses to manure-borne nutrients resulting from the land application of antibiotics- and coccidiostats-containing manures have been reported (Sims, 1997; Nahm, 2005). It is important to point out that composting of organic wastes including poultry litter is likely to reduce the concentration of most organic pesticides in the manure product (Fogarty and Tuovinen, 1991; Kawata *et al.*, 2006).

Although the use of antibiotics as growth promoters in poultry feed has been reduced in the U.S. (Singer and Hofacre, 2006), the potential environmental risks of proliferation and transport of antibiotic-resistant bacteria via land application of poultry is less documented (Kelley *et al.*, 1998; Brooks *et al.*, 2009) than human health impacts to antibiotic exposures (Powers and Angel, 2008).

#### ENDOCRINE-DISRUPTING COMPOUNDS

Endocrine-disrupting compounds (EDCs) are a group of compounds either synthesised or naturally present in the environment suspected to have adverse effects in animals and

humans. These EDC include several natural hormones excreted by the animal, and include pesticides, herbicides, plant oestrogens, and other compounds that disrupt endocrine systems (Powers and Angel, 2008). The EDC affect organisms by interfering with the binding of hormones to their receptors or disrupting the endocrine system by mimicking natural hormones (Colborn *et al.*, 1993).

Broilers can produce and excrete EDCs in the form of steroid hormones. Research has shown that broiler litter contains  $17\beta$ -oestradiol, oestrone, oestriol, and testosterone that can persist in poultry litter in measurable concentrations (Hanselman *et al.*, 2003; Jenkins *et al.*, 2006). The impact of the discharge of these compounds on the environment is poorly understood. In the U.S., a series of studies on land application of poultry litter to meet the crop nitrogen requirements of Georgia Piedmont soils under conventional till and no-till management (Jenkins *et al.*, 2006, 2009a,b), showed that the hormones  $17\beta$ -estradiol and testosterone did not appear to impact their background levels in soil, surface runoff, and subsurface drainage. These results and other studies indicate that both oestradiol and testosterone might be mineralized once they are in contact with soil. However, other studies indicated that fractions of oestradiol and testosterone appeared to resist mineralisation (Filay-Moore *et al.*, 2000; Jenkins *et al.*, 2006).

#### MICROBIAL LOAD

Poultry manure contains a large and diverse population of viruses, bacteria, fungi and protozoa. Microbial concentrations in poultry litter can exceed  $10^{10}$  cells/g (Acosta-Martinez and Harmel, 2006; Cook *et al.*, 2008; Rothrock *et al.*, 2008a), and gram positive bacteria (*i.e.* *Actinomycetes*, *Clostridia/Eubacteria*, *Bacilli/Lactobacilli*) account for nearly 90% of the microbial diversity (Lu *et al.*, 2003; Enticknap *et al.*, 2006; Lovanh *et al.*, 2007). While microbes perform a variety of different enzymatic and metabolic processes within the litter environment, two microbial groups of special interest to the poultry industry are nitrogen mineralizing microbes and pathogens.

Ammonia volatilisation results from the mineralisation of organic nitrogen in the poultry litter, namely uric acid and urea, and more than half of the nitrogen in poultry litter is lost as ammonia due to microbial activity (Brinson *et al.*, 1994; Moore *et al.*, 1996). Culture-based work identified *Bacillus* spp. (Ritz *et al.*, 2004) and *Arthrobacter* spp. (Schefferle, 1965; Kim and Patterson, 2003) as the dominant uric acid mineralising microbes in poultry litter. Based on DNA sequencing work, a single, previously unidentified, bacterial group dominated the urea mineralising bacterial clone libraries based on the urease enzyme (Rothrock *et al.*, 2008b). These specific microbial groups are present in sufficiently high concentrations (~1% of total microbial concentration; Cook *et al.*, 2008) to account for the high organic nitrogen mineralisation rates in untreated poultry (Gale and Gilmour, 1986; Moore *et al.*, 1995). Fungi, specifically *Aspergillus* spp., were recently shown to mineralise these predominant forms of organic nitrogen in poultry litter (Cook *et al.*, 2008), and this fungal activity must be taken into account.

Pathogens represent the second group of bacteria of importance to the poultry industry. Culture- and molecular-based work has shown that poultry litter is a reservoir for several zoonotic pathogens, including *Escherichia coli*, *Salmonella* spp., *Campylobacter jejuni*, *Listeria monocytogenes*, and *Clostridium perfringens* (Williams *et al.*, 1999; Terzich *et al.*, 2000; Line, 2002; Bull *et al.*, 2006; Line and Bailey, 2006; Rothrock *et al.*, 2008a). Contact between the birds and the litter represents a major route of contamination, with carcass contamination representing a dominant source of pathogen introduction into the meat processing plant (Rigby *et al.*, 1980a; Rigby *et al.*, 1980b; Izat *et al.*, 1988; De Boer and Hahne, 1990). Pathogen destruction may be required in some situations prior to land

spreading, a potential major source of pathogen introduction into the environment (Millner, 2009).

## **Uses of poultry litter**

### **AS A NUTRIENT SOURCE**

Poultry litter is often used as an organic nutrient source in forage, cereal and fibre crop production. The addition of poultry litter to tall fescue, orchard grass, bermuda grass has been shown to increase dry matter production (Sims and Wolf, 1994; McGrath *et al.*, 2009). In some cases the amount of N applied was in excess of the amount recommended for forage production, resulting in groundwater and surface water contamination through leaching and surface runoff (Sharpley *et al.*, 2007). Excessive application can cause undesirable effects on forage crops and animals consuming the forage (Mcginley *et al.*, 2003). During a seven year study in which over 18 tons/ha of broiler manure was applied annually to tall fescue used in a grazing study, Stuedemann *et al.* (1975) noticed problems with grass tetany and fat necrosis in beef cattle.

Increases in corn response to poultry manure addition have been noticed in a number of studies. In most studies, highest corn yields were obtained with the highest rate of manure addition, but the efficiency of manure N recovery by the crop decreased markedly as N rate increased, resulting in N leaching. For example, Sims (1987) obtained three year average efficiencies for N recovery of 50, 37, and 36%, respectively, for poultry manure application rates of 84, 168, and 252 kg potentially available N. The potentially available N from manure was calculated as 80% of (NH<sub>4</sub>-N + NO<sub>3</sub>-N) plus 60% of organic N (Bitzer and Sims, 1988). Similar decreases in N recovery with increasing N rate were observed in the case of fertilizer N application. In some studies, excessive application of fertilizer N or poultry manure has been shown to decrease corn yield, which has been attributed to adverse growing conditions ('salt injury') resulting from high salt concentration in the soil (Sims and Wolf, 1994).

Fertilizer studies with cotton in the U.S. showed that poultry litter is a valuable source of plant nutrients supplying N and metals Fe, Cu, Zn and Mn (Tewolde *et al.*, 2005). When poultry litter was applied in bands, N nutrition was more effective than regular broadcast; it increased lint yield and improved cotton fibre quality (Tewolde *et al.*, 2009).

Field experiments were conducted at Massey University, New Zealand to examine the fertilizer value of freshly applied and residual poultry manure (Mahimairaja *et al.*, 1995a). Fresh poultry manure, obtained from a layer unit, was mixed with elemental sulphur and phosphate rock and composted for five weeks under aerobic conditions. The value of poultry compost as an N source for cabbage was compared with urea. Both the compost and the urea were added at the start of the cabbage trial at a rate of 300 kg N/ha. After the harvest of the cabbage, maize was grown in the same plot to examine the residual value of the poultry manure. The harvestable fresh head yields of cabbage and the dry matter yield of fodder maize are presented in *Figure 1*. Although the highest yield of cabbage was obtained with the application of urea in combination with elemental sulphur and phosphate rock, the yield was not significantly different from the yield obtained by the poultry compost. The N use efficiency (kg fresh head yield per kg N applied) was slightly higher for the urea treatment than for the poultry compost (195 vs. 139). The apparent recovery of applied N was higher with the urea treatment (58.6%) than with the poultry compost (41.7%). There was no difference in maize yield between the urea and the poultry manure compost treatments. This suggests that poultry manure composts have a greater residual value than urea.




Figure 1 Effect of urea and poultry manure on harvestable yield of cabbage and dry matter yield of maize (Mahimairaja *et al.*, 1995a).

#### AS A SOIL AMENDMENT

Continuous cultivation of arable soils results in the deterioration of soil structure leading to reduced crop yield. For example, in the Manawatu region of New Zealand, continuous cultivation of maize has resulted in the deterioration of the physical conditions of the soils. A plant growth experiment was conducted in which the effects of poultry manure on the physical fertility of the cultivated soil and the growth of maize crop were examined. A soil that has undergone continuous cultivation of maize for 34 years and a pasture soil were used in the study. Poultry manure was compared with urea at an application level of 300 kg N/ha (Bolan *et al.*, 1992).

The addition of urea and poultry manure increased the dry matter yields of the maize crop in both cultivated and the pasture soils (Table 7). In the pasture soil, there was no significant difference in dry matter yields between the poultry manure and the urea treatments. However, in the case of the cultivated soil, poultry manure achieved greater yields than the urea treatment. Addition of poultry manure achieved similar dry matter yields in the pasture and the cultivated soils. The results indicated that improving the chemical fertility status of the cultivated soil alone through chemical fertilizer input is not enough to achieve the potential maximum yield of maize crop in these soils.

**Table 7** Effect of urea and poultry manure (PM) addition on dry matter yield of maize and soil properties.

Properties	Pasture soil			Cultivated soil		
	Control	Urea	PM	Control	Urea	PM
DM yield (g/pot)	130	158	155	70	120	161
Organic matter (g/kg soil)	55	57	85	25	27	79
Bulk density (kg/m <sup>3</sup> )	1060	1060	935	1125	1125	960
Water holding capacity (g/kg )	355	357	580	217	217	560
Oxygen diffusion rate (cm <sup>3</sup> /min)	21.4	18.5	26.4	10.2	8.2	25.3
Aggregate stability	35.6	34.5	41.2	12.2	13.5	40.5

The soil data (*Table 7*) indicated that the addition of poultry manure decreased the bulk density, increased the organic matter content, water holding capacity, oxygen diffusion rate and the aggregate stability of the soils. The effect of poultry manure on these physical properties was more pronounced in the cultivated soil than the pasture soil. These results indicate that the addition of poultry manure improved the physical fertility of the cultivated soil leading to increased maize growth. Compost products, including poultry litter, are used commonly as a mulching material for agricultural and horticultural crops to conserve soil moisture and to protect the surface feeding roots from drying during the summer periods (Eneji *et al.*, 2008; Agbede and Ojeniyi, 2009). Similarly the application of poultry litter has been shown to improve the biological fertility of mine tailings. Organic manures, such as poultry litter, are increasing being used in the rehabilitation of disturbed land resulting from mining and other industrial activities (Franzluebbers and Doraiswamy, 2007).

#### AS AN ANIMAL FEED

Poultry manure, either on its own or when mixed with feed grains, has been found to be a valuable feed for cattle and fish. Ruminants are able to utilise the urea nitrogen (uric acid) in poultry manure (Smith and Fries, 1973). However, the presence of foreign materials, such as plastic and glass, and feathers affects the digestibility of poultry waste and hence it is important to remove these from the litter before using it as a feed. It is also important to maintain low ash content. When large quantities of soil are removed with the litter, the ash content increased dramatically. It is recommended that poultry litter with ash contents exceeding 28% should not be fed to cattle (Williams *et al.*, 1999).

From a hygiene perspective, unprocessed poultry waste contains potential pathogenic microorganisms such as *Clostridium*, *Salmonella* and *Enterobacter spp.* Hence proper processing to reduce the number of these microorganisms or render the waste free of pathogens is required (McCaskey and Anthony, 1979; Kawata *et al.*, 2006). In addition, as noted above, feed additives such as antibiotics, arsenicals, and coccidiostats are added in poultry diet, which can be excreted as waste by-products. Furthermore, some of the fungal species that are indigenous to the manure or litter can result in the production of mycotoxins. Pathogenic microorganisms can be destroyed by chemical, fermentation, ensilation or heat processing (McCaskey and Martin, 1988; Cook *et al.*, 2008).

Although disease problems have not been reported from feeding poultry manures to farm animals under acceptable conditions but Cu toxicity has been found to be a problem, especially in sheep. When an excess of Cu is added as a growth promoter to poultry diets, it is excreted in high concentration in the manure since Cu is poorly absorbed in the bird's digestive system. Sheep are less tolerant than other livestock

species to high dietary levels of Cu when fed broiler litter (Fontenot and Webb, 1975; Sharma et al., 2005).

Processed chicken manure and litter have been used as a feed ingredient for almost 40 years in the U.S. These contain large amounts of protein, fibre, and minerals, and have been deliberately mixed into ruminant feed for these nutrients. Normally, this animal waste is used by small farmers and owners of beef and dairy herds as a winter supplement for cows and weaned calves. While the practice of feeding poultry manure to animals seems unpleasant, the use of this product is safe as long as it meets certain specifications listed in *Table 8*. For, example, the Association of American Feed Control Officials (AAFCO) has established ‘Standard Names and Definitions’ for three processed poultry waste products used as an animal feed as follows (FDA, 2009):

- Dried Poultry Waste: a processed animal waste product composed primarily of faeces from commercial poultry, which has been thermally dehydrated to moisture content not in excess of 15%. It must contain not less than 18.0% crude protein, and not more than 15% crude fibre, 30% ash, and 1% feathers.
- Dried Poultry Waste-non protein nitrogen (NPN) Extracted: a processed animal waste product composed primarily of faeces from commercial poultry which has been processed to remove part or all of the equivalent crude protein and non protein nitrogen (NPN) as urea and/or uric acid, and which has been thermally dehydrated to a moisture content not in excess of 15%. It must contain not less than 11% crude protein, and not more than 15% crude fibre, 30% ash, and 1% feathers.
- Dried Poultry Litter: a processed animal waste product composed of a processed combination of faeces from commercial poultry together with litter that was present in the floor production of poultry, which has been artificially dehydrated to a moisture content not in excess of 15%. It must contain not less than 18% crude protein, and not more than 25% crude fibre, 20% ash, and 4% feathers.

**Table 8 Specifications for using poultry manure as an animal feed and the variation in some of the characteristics.**

Characteristics	Specifications			Variation	
	Broiler	Layer	Limits	Broiler	Layer
Moisture (%)	12	12	Maximum	10-24	56 - 87
Crude protein (%)	24	22	Minimum	10-26	5 - 24
Uric acid CP (%)	60	60	Maximum	40-60	50 - 80
Fat (%)	1.5	1.5	Minimum		
Fibre (%)	15	15	Maximum		5 - 25
Ash (%)	15	25	Maximum	10-26	7 - 15
Feathers (%)	1	1	Maximum		
Calcium (%)	3.5	8	Maximum	1.5-3.0	5 - 12
Phosphorus (%)	1.5	2	Minimum	1.2-1.8	1.5-2.9
Sodium (%)	0.5	0.5	Maximum		
Silica (%)	0.5	0.5	Maximum		
Copper (%)	0.005	0.005	Maximum	0.002-0.035	0.001-0.007

The AAFCO specifications require that processed animal waste products do not contain extraneous materials such as metal, glass, nails or other harmful matter. They must be free of pathogenic organisms, pesticide residues, parasites, or drug residues, which could

be harmful to animals or could result in residues in human food products or by-products of animals at levels in excess of those allowed by State or Federal statute or regulation.

If a product contains drug residues, then the label must contain the following statement in bold face type: **'Warning: this product contains drug residues. Do not use within 15 days of slaughter and do not use 15 days prior to or during the food production period of dairy animals and laying hens.'** If the product contains 25 mg/kg or greater of Cu, the following statement in bold face type is required: **'Warning: contains high levels of copper: do not feed to sheep.'** According to AAFCO using adequately processed poultry waste in animal feed may not be aesthetically pleasing but it is safe, nutritionally valid, and environmentally sound.

Although poultry litter has been found to be more valuable as a feedstuff than as a fertiliser, the use of poultry waste as a feed ingredient is becoming a less common practice within the livestock industry. The public perception of the concept of coprophagy (animal consumption of faecal material) contributes to the limited use of this practice as a major means of recycling poultry wastes. It is also likely that concerns about infection by enterohemorrhagic strains of *E. coli*, and the perception of transmission of neurodegenerative disease such as scrapie, Creutzfeldt-Jacob disease (CJD), and bovine spongiform encephalopathy (BSE) will naturally discourage the practice of feeding farm animals with any waste derived from animal-based industries (Williams *et al.*, 1999).

#### AS A FUEL SOURCE

Poultry litter can be burnt directly as a fuel source to produce heat energy. One of the problems with using poultry litter as a fuel source is its relatively high moisture content. The moisture content should be less than 15% in order to achieve the maximum heat energy during burning. Alternatively, the anaerobic digestion of poultry waste yields biogas, a combustible gas composed of approximately 60% methane, 38% carbon dioxide, and mixture of water vapour, ammonia and hydrogen sulphide. Biogas (also known as 'producer' gas) may be used as an energy source for burning as heat or as fuel for internal combustion engines to generate electricity. Both the burnt ash material and the post-anaerobic digested solids may be used as fertiliser and animal feed supplement (Liedl *et al.*, 2006; Blake *et al.*, 2007). Few producers, however, utilise anaerobic digestion as a method of poultry waste treatment because of unfavourable economics, low biogas yield for litter-based systems and technical operational difficulties (Williams *et al.*, 1999).

Poultry litter combustion has received major attention as a method to produce heat and electricity at large centralised facilities (Kelleher *et al.*, 2002; Turnell *et al.*, 2007; Fibrowatt, 2008). In the search to make electric power from renewable 'green' sources, a number of states in the US have turned to thermal conversion of biomass. For example, Minnesota produces over two million tons of turkey and broiler waste — the fuel for 'poultry power.' Now some Minnesota turkey farmers are working with a British company (Fibrowatt) that built a manure-fired power plant in central Minnesota in 2007 (Fibrowatt, 2008). The plant burns nearly half a million tons of poultry litter every year, generating 50 Megawatts of electricity that will be sufficient enough to supply 40000 households.

Similarly, there is a proposal to build a poultry litter-based power plant in Western Australia. On the basis of the Fibrowatt experience, it is possible to produce approximately 0.700 Megawatts of electricity by burning around 7000 tons of poultry litter produced in the proposed broiler plant. This will be sufficient enough to supply around 900 households. However, a limitation to the adoption of this technology is the high capital investment and public concern for potential emission of particulate matter,

nitrogen oxides, carbon monoxide, and sulphur dioxide from large centralised combustion facilities burning poultry litter (Turnell *et al.*, 2007). Nevertheless, it is possible to utilise the litter as a fuel source to produce hot water, which will be useful for a poultry processing unit.

### **Improving the value of poultry litter**

Fresh poultry waste is difficult to handle due to its high moisture content and odour. With the introduction of barn system (confined production), it is now possible to produce relatively dry manure. Another important problem of poultry waste is the loss of N during the storage, drying, handling, and subsequent land application (Mahimairaja *et al.*, 1994; Adeli *et al.*, 2009). The loss of N occurs mainly through ammonia volatilisation and denitrification during handling and through nitrate leaching after application to land. Gaseous losses of N are of particular concern because they not only reduce the fertiliser value of poultry waste, but also form atmospheric pollutants, and deteriorate the environmental health. Excessive application of fresh poultry manure also results in the accumulation of ammonia in soils causing injury to seedlings and roots. Thus, poor management of this valuable resource could be damaging to crops and also leads to pollution of surface and groundwaters.

Appropriate technologies, which are environmentally viable and economically feasible, are needed for efficient management of poultry waste. This can be achieved through proper composting of the manure and the appropriate feed management practices.

#### **COMPOSTING**

Manure by-products undergo a number of treatment processes that include *in-situ* biological pond treatment, composting, and specific chemical treatment. Aerobic composting is most commonly practiced to overcome some of the problems associated with the handling and disposal of poultry manure. It reduces the bulkiness of the waste and yields a stabilized product suitable for handling and land application. Composting eliminates animal and human pathogens and could reduce the risks of polluting groundwater. Development of appropriate methods of composting poultry waste with suitable amendments could greatly reduce the nutrient losses and at the same time helps to minimise environmental pollution.

Use of amendments, such as straw, peat, woodchip, paperwaste, elemental sulphur and zeolite helps to reduce the N losses during composting of poultry waste. Aerobic composting with cereal straw, which contains readily decomposable carbon, is most effective in reducing the N loss (Preusch *et al.*, 2002). Nutrient-rich ecofriendly compost can be prepared by composting poultry waste with phosphate rock and elemental sulphur. This can be used as a valuable nutrient source especially for organic farming practises. Addition of small amount of elemental sulphur reduces the pH of the compost, thereby reducing the volatilisation losses of ammonia (Mahimairaja *et al.*, 1993). It also enhances the dissolution of rock phosphate and enriches the compost with P and S nutrients (Mahimairaja *et al.*, 1995b). Use of alum and zeolite to reduce ammonia volatilisation as well as P solubility of poultry litter has provided encouraging results (Kithome *et al.*, 1999; Delaune *et al.*, 2004; Guo and Song, 2009). The potential to use such amendments not only addresses environmental concerns but also improves the housing environments for the birds and workers by reducing ammonia concentrations, as well as improving the N:P ratio of the manure for crop utilisation. Hence it forms an economically viable option for poultry growers (Moore *et al.*, 1996; Sharpley *et al.*, 2007).

Although composting is basically aimed at achieving a stable manure product, it may also affect both the total content and the speciation of metals in manure by-products. For example, Sistani *et al.* (2001) observed that composting decreased Zn content in poultry manure. Moore *et al.* (1998) observed that treatment of poultry manure with alum [Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>] resulted in the immobilization of Cd, Cu and Zn, thereby decreasing their concentration in soil solution.

#### FEED MANAGEMENT

Improving feed efficiency and health of animals that minimises the incidence of disease outbreaks is an important consideration in intensive, confined animal production systems (Nahm, 2002). Efficient nutrient utilisation using growth promoters not only reduces the cost of production but also provides a major means of reducing potential contaminants, such as N and P in poultry manure. However, since some of the growth promoters contain metals, this practice is likely to result in elevated concentration of these elements in manure by-products. Precision formulation and feeding of diets to meet but not to exceed the nutritional requirements will likely be a viable component of waste management strategies.

Nutrient excretion in poultry manure results mainly from inefficiencies associated with digestion and metabolism. Addition of feed supplements and modifying feeding programme to improve nutrient efficiency can result in significant decreases in the N, P and odour of poultry manures (Nahm, 2002). Examples of these methods include (Selle and Ravindran, 2007; Silversides and Hruby, 2009): (i) addition of synthetic amino acids and reducing protein contents in the feed have resulted in a decrease in manure N contents by 10 to 27% in the broiler units; (ii) enzyme supplementation has resulted in a 12 to 15% reduction in the dry weight of broiler manure; (iii) phytase supplementation has resulted in the reduction of P in poultry manure by 25 to 60%; (iv) formulation of diet matching the requirements has reduced the N content in manure by 10 to 15%; (v) phase feeding reduced N and P in the manure by 10-33%; (vi) use of highly digestible raw materials in feed has reduced the N and P excretion by 5%; (vii) certain feed manufacturing techniques (reducing particle size and pelletising) can significantly increase dry matter digestibility, leading to reduced manure production; (viii) use of alternative coccidiostats, such as 'ionophores' (instead of arsenical compounds) can achieve drastic reduction in As level in poultry litter; and (ix) similarly the use of non-metal containing growth promoters helps to reduce the concentration of metals, metals such as Cu and Zn in poultry litter.

#### **Best management practices for the beneficial use of poultry litter**

There is an urgent need to formulate strict regulations governing the safe disposal and handling of poultry waste in order to minimise the environmental impact. Management of poultry waste must be integrated into a broader nutrient management programme in agriculture. Guidelines on specific land application, optimal loading rates, and permissible limits of nutrients, heavy metals, antibiotics, and coccidiostats in poultry waste, are needed. Research-based information on the transformation and plant availability of nutrients and heavy metals in poultry wastes is needed for efficient management of this resource as a nutrient resource and soil amendment. Recycling of poultry waste in fish and cattle feed and in power (electricity) generation should also be given consideration for efficient and profitable management of poultry waste.

The components of an effective management programme for the agricultural use of poultry litter include (RIRDC, 2009): (i) site selection; (ii) production and collection; (iii)

storage, handling and treatment; and (iv) transport and land application. Site specific optimisation of each of these components is essential to maximise the beneficial use of litter resources, thereby avoiding the pollution of the nearby environment (Sims and Wolf, 1994).

Management programmes for poultry litter must reflect both the potential value of the litter as a resource and a realistic appraisal of the negative effects litter constituents may have on the environment. High transportation costs and lack of distribution infrastructure are some of the reasons why a comprehensive approach to poultry litter management be developed to take advantage of all beneficial end uses for the diverse waste products of this industry.

#### SITE SELECTION

- Site location for a poultry unit should be based on the facilities for production, storage and treatment of wastes and the suitability of the soils on the site for land application of litter.
- Proximity to streams, ponds, and drainage pathways and an understanding of groundwater hydrology are taken into consideration.

#### COLLECTION AND STORAGE

- A waste management plan that aims to reduce the volume of waste production will facilitate the ease and efficiency of the operations of the plan.
- Litter collection should be closely linked to storage capacity so that these resources are protected from unfavourable weather conditions and maintained in good physical conditions enabling easy application.
- The unfavourable economics of litter transportation often result in limited distribution of nutrients throughout the farm, causing a build up of some nutrients (*e.g.*, P) to excessive levels in the field at short distances from the site of waste generation.

#### TREATMENT

Composting is the most common treatment aimed at achieving a stable organic manure product. It also achieves partial or complete elimination of microbial pathogens.

- Compost areas need to have an impermeable base to avoid leaching and possible groundwater contamination.
- The composting site should be in an elevated area or a bunding may be required in order to prevent extraneous runoff entering the pile and becoming contaminated.
- The compost pile should be protected from rain in order to overcome leaching of nutrients and contaminants, and from wind to overcome the problems associated with odorous gases.
- Nutrient rich runoff from the compost piles should be collected in a sump or dam and may be reused.
- Compost pile need to be carefully managed to avoid dust and odour emission. If the compost is too dry the process will be slowed and excessive dust may be generated. If the compost becomes too wet, it may become anaerobic and result in excessive odour emissions. Optimum moisture content is around 50 - 55% (wet basis).

#### LAND APPLICATION

- Timing of land application of poultry litter should be aimed at maximising crop recovery of nutrients and closely related to production patterns and storage capacities.

*Poultry litter: N.S. Bolan et al.*

- Poultry compost should be added during the active growth of the crop or immediately before planting. For example, application of poultry manure during fall and winter, when crops have not been able to utilise nutrients is normally not encouraged.
- Application should be based on balanced nutrient requirements of the crop.
- Litter should be incorporated into soil; this will reduce the gaseous losses of N and the runoff losses of nutrients and contaminants in litter.

#### ENVIRONMENTAL MONITORING

Environmental monitoring is an important component of best management practices to achieve both sustainable production and environmental protection when poultry litter is applied to land. Environmental monitoring includes regular analysis of manure samples, maintaining accurate records of all activities in the farm, and regular soil and drainage water sampling and their analysis for nutrients and other contaminants.

#### Conclusions

Poultry litter is a good organic source of nutrients for raising crops, such as maize. However, the loss of N through ammonia volatilisation is a major issue during handling and land application of poultry litter. These losses can be minimized through proper composting process in the presence of organic amendments, such as cereal straw.

The success of utilisation of poultry litter as a valuable nutrient source and soil conditioner depends on developing technologies to produce uniform value-added products, and design of equipment for uniform waste product field application. Response of pasture, arable and horticultural crops, and silviculture species to poultry waste applications and their beneficial effects on soil physical and chemical properties should be examined in order to maximise the utilisation of poultry litter. The bioavailability of nutrient and heavy metals and their movement into surface water and groundwater as influenced by the loading rate of poultry litters need to be studied further, and site specific best management practices for safe and beneficial utilisation of poultry manure for sustainable production with minimum impact on environment should be developed. This will allow technologies that provide alternative to land application of poultry wastes to be developed.

#### Acknowledgements

The senior author thanks CRC CARE for providing funding (No 2-3-09-07/08) to undertake research on landfill site remediation as part of the review was derived from this project.

#### References

- ACOSTA-MARTINEZ, V. and HARMEL, R.D. (2006) Soil microbial communities and enzyme activities under various poultry litter application rates. *Journal of Environmental Quality* **35**: 1309-1318.
- ADELI, A., TEWOLDE, H., SISTANI, K.R. and ROWE, D.E. (2009) Broiler litter fertilization and cropping system impacts on soil properties. *Agronomy Journal* **110**: 1304-1310.
- AGBEDE, T.M. and OJENIYI, S.O. (2009) Tillage and poultry manure effects on soil fertility and sorghum yield in southwestern Nigeria. *Soil and tillage research* **104**: 74-81.

- ALVA, A.K., HUANG, B. and PARAMASIVAM, S. (2000) Soil pH affects copper fractionation and phytotoxicity. *Soil Science Society of America Journal* **64**: 955-962.
- ANEJA, V.P., SCHLESSINGER, W.H., NIYOGLI, D., JENNINGS, G., GILLIAM, W., NIGHTON, R.E., DUKE, C.S., BLUNDEN, J. and KRISHNAN, S. (2006) Emerging national research needs for agricultural air quality. *Eos, Transactions of the American Geophysical Union* **87**: 25-36.
- ASABE (2005) Manure production characteristics. ASABE Standard D384.2. American Society of Agricultural and Biological Engineers. St. Joseph, MI.
- AXTELL, R.C. (1986) Fly management in poultry production - cultural, biological, and chemical. *Poultry Science* **65**: 657-667.
- BERNHART, M., FASINA, O.O., FULTON, J. and WOOD, C.W. (2010) Compaction of poultry litter. *Bioresource technology* **101**: 234-238.
- BHATTACHARYA, A.C. and TAYLOR, J.C. (1975) Recycling animal wastes as a feedstuff: A review. *Journal of Animal Science* **41**: 1438-1457.
- BITZER, C.C. and SIMS, J.T. (1988) Estimating the availability of nitrogen in poultry manure through laboratory and field studies. *Journal of Environmental Quality* **17**: 47-54.
- BLAKE, J.P., HESS, J.B. and BOCK, B.R. (2007) Nutritional and economic value of poultry litter ash as a feed stock. *Proceedings International Symposium on Air Quality and Waste Management for Agriculture Proceedings*, Broomfield, CO., ASABE Pub. No. 701P0907cd.
- BOLAN, N.S., MAHIMAIRAJA, S. and HEDLEY, M.J. (1992) Improving the fertilizer value of poultry manures. *Proceedings of the Seminar on 'Eggs '92- winning in a deregulated environment,'* Egg Producers Federation of New Zealand, pp. 1-15.
- BOLAN, N.S., ADRIANO, D.C. and MAHIMAIRAJA, S. (2004) Distribution and Bioavailability of Trace Elements in Livestock and Poultry Manure By-Products. *Critical Reviews in Environmental Science and Technology* **34**: 291-338
- BRINSON, S.E., JR., CABRERA, M.L. and TYSON, S.C. (1994) Ammonia volatilisation from surface-applied, fresh and composted poultry litter. *Plant and Soil* **167**: 213-218.
- BROOKS, J.P., ADELI, A., READ, J.J. and MCLAUGHLIN, M.R. (2009) Rainfall simulation in greenhouse microcosms to assess bacterial-associated runoff from land applied poultry litter. *Journal of Environmental Quality* **38**: 218-229.
- BULL, S.A., ALLEN, V.M., DOMINGUE, G., JORGENSEN, F., FROST, J.A., URE, R., WHYTE, R., TINKER, D., CORRY, J.E.L., GILLARD-KING, J. and HUMPHREY, T.J. (2006) Sources of *Campylobacter* spp. colonizing housed broiler flocks during rearing. *Applied and Environmental Microbiology* **72**: 645-652.
- BUREL, C. and VALAT, C. (2009) The effect of the feed on the host-microflora interactions in poultry: an overview. *Proceedings in International Symposium on Sustainable animal production - the challenges and potential developments for professional farming for 13<sup>th</sup> Congress of the International Society for Animal Hygiene* pp.365-383.
- CAPER, S.G., TANNER, J.T., FRIEDMAN, M.H. and BOYER, K.W. (1978) Multielement analysis of animal feed, animal wastes, and sewage sludge. *Environmental Science and Technology* **12**: 785-790.
- CASEY, K.D., BICUDO, J.R., SCHMIDT, D.R., SINGH, A., GAY, S.W., GATES, R.S., JACOBSEN, L.D. and HOFF, S.J. (2006) Air quality and emissions from livestock and poultry production/waste management systems, in: RICE, J.M., CALDWELL, D.F. & HUMENIK, F.J. (Eds.) *Animal Agriculture and the Environment: National Center for Manure and Animal Waste Management White Papers*, Publication No 913C0306, pp. 1-40 (St. Joseph, MI, ASABE).
- CHAN, K.Y., VAN ZWIETEN, L., MESZAROS, L., DOWNIE, A. and JOSEPH, S. (2008) Using poultry litter biochars as soil amendments, *Australian Journal of Soil Research* **46**: 437-444.
- CHRISTEN, K. (2001) Chickens, manure, and arsenic. *Environmental Science and Technology* **35**: 184A-185A.
- COLBORN, T., VOM SAAL, F.F. and SOTO, A.M. (1993) Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environmental Health Perspectives* **101**: 378-384.
- COOK, K.L., ROTHROCK, M.J., WARREN, J.G., SISTANI, K.R. and MOORE, P.A. (2008) Effect of alum treatment on the concentration of total and ureolytic microorganisms in poultry litter. *Journal of Environmental Quality* **37**: 2360-2367.
- DE ABREU, M.F. and BERTON, R.S. (1996) Comparison of methods to evaluate heavy metals in organic wastes. *Communications in Soil Science and Plant Analysis* **27**: 1125-1135.
- DE BOER, E. and HAHNE, M. (1990) Cross-contamination with *Campylobacter jejuni* and *Salmonella* spp. from raw chicken products during food preparation. *Journal of Food Protection* **53**: 1067-1068.
- DELAUNE, P.B., MOORE, P.A. JR, DANIEL, T.C. and LEMUNYON, J.L. (2004) Effect of chemical and microbial amendment on ammonia volatilisation from composting poultry litter. *Journal of Environmental Quality* **33**: 728-734.
- DAZ, D.A.R., SAWYER, J.E. and MALLARINO, A.P. (2008) Poultry manure supply of potentially available nitrogen with soil. *Agronomy journal* **100**: 1310-1317.

- ECK, H.V. and STEWART, B.A. (1995) Manures, in: REICHIGL, J.E. (Ed.) *Soil Amendments and Environmental Quality*, pp. 169-198 (Boca Raton, FL, Lewis Publishers).
- EDWARDS, D.R. and DANIEL, T.C. (1992) Environmental impacts of on-farm poultry waste disposal-A review. *Bioresource Technology* **41**: 9-33.
- EDWARDS, J.H. and SOMESHWAR, A.V. (2000) Chemical, physical, and biological characteristics of agricultural and forest by-products for land application, in: DICK, W.A. (Ed.) *Land Application of Agricultural, Industrial, and Municipal By-products*, pp. 1-62 (Madison, WI, Soil Science Society of America).
- ENEJI, A.E., INANAGA, S., LI, X., AN, P., LI, J., DUAN, L. and LI, Z. (2008) Effectiveness of mulching vs. incorporation of composted cattle manure in soil water conservation for wheat based on eco-physiological parameters. *Journal of Agronomy and Crop Science* **194**: 26-33.
- ENTICKNAP, J.J., NONOGAKI, H., PLACE, A.R. and HILL, R.T. (2006) Microbial diversity associated with odor modification for production of fertilizers from chicken litter. *Applied and Environmental Microbiology* **72**: 4105-4114.
- EPSTEIN, E. and MOSS, L.H. (2006) A comparison of characteristics of manures, biosolids, and mineral fertilizers. *Journal of Residuals Science and Technology* **3**: 35-42.
- FDA (2009) Food and Drug Administration; Center for Veterinary Medicine. The Use of Chicken Manure/Litter in Animal Feed. Available at: <http://www.pickle-publishing.com/papers/chicken-litter-animal-feed.htm>.
- FIBROWATT (2008) Power from poultry litter. Available at: <http://fibrowattusa.com>.
- FILAY-MOORE, O., HARTEL, P.G. and CABRERA, M.L. (2000) 17 $\beta$ -estradiol and testosterone in soil and runoff from grasslands amended with broiler litter. *Journal of Environmental Quality* **29**: 1694-1611.
- FOGARTY, A.M. and TUOVINEN, O.H. (1991) Microbiological degradation of pesticides in yard waste composting. *Microbiological reviews* **55**: 225-233.
- FONTENOT, J.P. and WEBB JR, K.E. (1975) Health aspects of recycling animal wastes by feeding. *Journal of Animal Science* **40**: 1267-1277.
- FRANZLUEBBERS, A.J. and DORAISWAMY, P.C. (2007) Carbon sequestration and land degradation. *Climate and land degradation in: International Workshop on Climate and Land Degradation*: 343-358 (Arusha, Tanzania).
- FRIEND, A.L., ROBERTS, S.D., SCHOENHOLTZ, S.H., MOBLEY, J.A. and GERARD, P.D. (2006) Poultry litter application to Loblolly pine forests: Growth and nutrient containment. *Journal of Environmental Quality* **35**: 837-848.
- FSA (2007) Feedlot Services Australia Pty Ltd. Available at: <http://www.fsaconsulting.net/>.
- GALE, P.M. and GILMOUR, J.T. (1986) Carbon and nitrogen mineralization kinetics for poultry litter. *Journal of Environmental Quality* **15**: 423-426.
- GRIMES, J.L., SMITH, J. and WILLIAMS, C.M. (2002) Some alternative litter materials used for growing broilers and turkeys. *World's Poultry Science Journal* **58**: 515-526.
- GUO, M. and SONG, W. (2009) Nutrient value of alum-treated poultry litter for land application. *Poultry science* **88**: 1782-1792.
- GUPTA, G. and CHARLES, S. (1999) Trace elements in soils fertilized with poultry litter. *Poultry Science* **78**: 1695-1698.
- HANSELMAN, T.A., GRAETZ, D.A. and WILKIE, A.C. (2003) Manure-borne estrogens as potential environmental contaminants: A review. *Environmental Science and Technology* **37**: 5471-5478.
- HARMEL, R.D., TORBERT, H.A., HAGGARD, B.E., HANEY, R. and DOZIER, M. (2004) Water quality impacts of converting to a poultry litter fertilization strategy. *Journal of Environmental Quality* **33**: 2229-2242.
- HARMEL, R.D., SMITH, D.R., HANEY, R.L. and DOZIER, M. (2009) Nitrogen and phosphorus runoff from cropland and pasture fields fertilized with poultry litter. *Journal of soil and water conservation* **64**: 400-412.
- HE, Z., ENDALE, D.M., SCHOMBERG, H.H. and JENKINS, M.B. (2009) Total phosphorus, zinc, copper, and manganese concentrations in Cecil soil through 10 years of poultry litter application. *Soil Science* **174**: 687-695.
- HOBSON-FROHOCK, A. and JOHNSON, H.A. (2006) Coccidiostat residues in poultry excreta. *Journal of the Science of Food and Agriculture* **34**: 37-44.
- IZAT, A.L., GARDNER, F.A. and GOLAN, F.A. (1988) Incidence and level of *Campylobacter jejuni* contamination in modern broilers production and processing systems. *Journal of Food Protection* **54**: 259-262.
- JACKSON, B.P. and BERTSCH, P.B. (2001) Determination of arsenic speciation in poultry wastes by IC-ICP-MS. *Environmental Science and Technology* **35**: 4868-4873.
- JACKSON, B.P. and MILLER, W.P. (2000) Soil solution chemistry of a fly ash-, poultry litter-, and sewage sludge-amended soil. *Journal of Environmental Quality* **29**: 430-436.
- JACKSON, B.P., BERTSCH, P.M., CABRERA, M.L., CAMBERATO, J.J., SEAMAN, J.C. and WOOD, C.W. (2003) Trace element speciation in poultry litter. *Journal of Environmental Quality* **32**: 535-540.

- JENKINS, M.B., ENDALE, D.M., SCHOMBERG, H.H. and SHARPE, R.R. (2006) Fecal bacteria and sex hormones in oil and runoff from cropped watersheds amended with poultry litter. *Science of the Total Environment* **358**: 164-177.
- JENKINS, M.B., ENDALE, D.M., SCHOMBERG, H.H., HARTEL, P.G. and CABRERA, M.L. (2009a) 17 $\beta$ -Estradiol and testosterone in drainage and runoff from poultry litter applications to tilled and no-till crop land under irrigation. *Journal of Environmental Management* **90**: 2659-2664.
- JENKINS, M.B., TRUMAN, C.C., SIRAGUSA, G., LINE, E., BAILEY, J.S., FRYE, J., ENDALE, D.M., FRANKLIN, D.H., SCHOMBERG, H.H., FISHER, D.S. and SHARPE, R.R. (2009b) Rainfall and tillage effects on transport of fecal bacteria and sex hormones 17 $\beta$ -Estradiol and testosterone from broiler litter applications to a Georgia Piedmont Ultisol. *Science of the Total Environment* **403**: 154-163.
- JONGBLOED, A.W. and LENIS, N.P. (1998) Environmental concerns about animal manure. *Journal of Animal Science* **76**: 2641-2648.
- KAISER, D.E., MALLARINO, A.P. and HAQ, M.U. (2009) Runoff phosphorus loss immediately after poultry manure application as influenced by the application rate and tillage. *Journal of Environmental Quality* **38**: 299-308.
- KARCI, A. and BALCIOGLU, I.A. (2009) Investigation of the tetracycline, sulfonamide, and fluoroquinolone antimicrobial compounds in animal manure and agricultural soils in Turkey. *Science of the total environment* **407**: 4652-4664.
- KAWATA, K., NISSATO, K., SHIOTA, N., HORI, T., ASADA, T. and OIKAWA, K. (2006) Variation in pesticide concentrations during composting of food waste and fowl droppings. *Bulletin of environmental contamination and toxicology* **77**: 391-398.
- KELLEHER, B.P., LEAHY, J.J., HENIHAN, A.M., O'DWYER, T.F., SUTTON, D. and LEAHY, M.J. (2002) Advances in poultry litter disposal technology – a review. *Bioresource Technology* **83**: 27-36.
- KELLEY, T.R., PANCORBO, O.C., MERKA, W.C. and BARNHART, H.M. (1998) Antibiotic resistance of bacterial litter isolates. *Poultry Science* **77**: 243-247.
- KELLEY, T.R., PANCORBO, O.C., MERKA, W.C., THOMPSON, S.A., CABRERA, M.L. and BRNHAT, H.M. (1996) Elemental concentrations of stored whole and fractionated broiler litter. *Journal of Applied Poultry Research* **5**: 276-281.
- KIM, W.K. and PATTERSON, P.H. (2003) Effect of minerals on activity of microbial uricase to reduce ammonia volatilisation in poultry manure. *Poultry Science* **82**: 223-231.
- KITHOME, M., PAUL, J.W. and BOMKE, A.A. (1999) Reducing nitrogen losses during simulated composting of poultry manure using adsorbents or chemical amendments. *Journal of Environmental Quality* **28**: 194-201.
- KRISHNAMACHARI, K.A.U.R. (1987) Fluorine, in: MERTZ, W. (Ed.) *Trace Elements in Human and Animal Health*, Vol 1, 5<sup>th</sup> Ed. pp. 265-416 (San Diego, CA, Academic Press).
- KUNKLE, W.E., CARR, L.E., CARTER, T.A. and BOSSARD, E.H. (1981) Effect of flock and floor type on levels of nutrients and heavy metals in broiler litter. *Poultry Science* **60**: 1160-1164.
- LEYTEM, A.B., PLUMSTEAD, P.W., MAGUIRE, R.O., KWANYUEN, P. and BRAKE, J. (2007) What aspect of dietary modification in broilers controls litter water-soluble phosphorus: dietary phosphorus, phytase, or calcium? *Journal of Environmental quality* **36**: 453-463.
- LIEDL, B.E., BOMBARDIERE, J. and CHATFIELD, J.M. (2006) Fertilizer potential of liquid and solid effluent from thermophilic anaerobic digestion of poultry waste. *Water Science Technology* **53**: 69-79.
- LINE, J.E. (2002) *Campylobacter* and *Salmonella* populations associated with chickens raised on acidified litter. *Poultry Science* **81**: 1473-1477.
- LINE, J.E. and BAILEY, J.S. (2006) Effect of on-farm acidification treatments on *Campylobacter* and *Salmonella* populations in commercial broiler houses in northeast Georgia. *Poultry Science* **85**: 1529-1534.
- LOVANH, N., COOK, K.L., ROTHROCK JR., M.J., MILES, D.M. and SISTANI, K. (2007) Spatial shifts in microbial population structure within poultry litter associated with physicochemical properties. *Poultry Science* **86**: 1840-1849.
- LU, J., SANCHEZ, S., HOFRADE, C., MAURER, J.J., HARMON, B. and LEE, M.D. (2003) Evaluation of broiler litter with reference to the microbial composition as assessed by using 16S rRNA and functional gene markers. *Applied and Environmental Microbiology* **69**: 901-908.
- MAGUIRE, R.O., PLUMSTEAD, P.W. and BRAKE, J. (2006) Impact of diet, moisture, location, and storage on soluble phosphorus in broiler litter breeder manure. *Journal of Environmental Quality* **35**: 858-865.
- MAHIMAIRAJA, S., BOLAN, N.S. and HEDLEY, M.J. (1995a) Agronomic effectiveness of poultry manure composts. *Communications in Soil Science and Plant Analysis* **26**: 1843-1861.
- MAHIMAIRAJA, S., BOLAN, N.S. and HEDLEY, M.J. (1995b) Dissolution of phosphate rocks during the composting of poultry manure. *Fertilizer Research* **40**: 93-104
- MAHIMAIRAJA, S., BOLAN, N.S. and HEDLEY, M.J. (1993) Absorption of ammonia released from poultry manure onto soil and bark and the use of absorbed ammonia in the dissolution of phosphate rock. *Compost Science and Utilisation* **1**: 10-21.

- MAHIMAIRAJA, S., BOLAN, N.S., HEDLEY, M.J. and MACGREGOR, A.N. (1994) Transformation and loss of nitrogen in poultry manures during composting. *Bioresource Technology* **47**: 265-273.
- MALONE, G.W. (1992) Nutrient enrichment in integrated broiler production systems. *Poultry Science* **71**: 1117-1122.
- MCBRIDE, M.B. and SPIERS, G. (2001) Trace element content of selected fertilizers and dairy manures as determined by ICP-MS. *Communications in Soil Science and Plant Analysis* **32**: 139-156.
- MCCASKEY, T.A. and ANTHONY, W.B. (1979) Human and animal health aspects of feeding live stock excreta. *Journal of Animal Science* **48**: 163-177.
- MCCASKEY, T.A. and MARTIN JR, J.B. (1988) Evaluation of a process for improved quality and microbiological safety of broiler litter. *Biological Wastes* **25**:209-218.
- MCDONALD, J.M., RIBAUDO, M.O., LIVINGSTON, M.J., BECKMAN, J. and WANG, H. (2009) Manure Use for Fertilizer and for Energy – Report to Congress. Economic Research Service, U.S. Department of Agriculture, Washington, D.C.
- MCGINLEY, B.C., COFFEY, K.P., HUMPHRY, J.B., SAUER, T.J. and GOODWIN, H.L. (2003) Mineral content of forages grown on poultry litter-amended soils. *Journal of Animal Science* **81**: 106-111.
- MCGRATH, S., MAGUIRE, R.O., TACY, B.F. and KIKE, J.H. (2009) Improving soil nutrition with poultry litter application in low input forage systems. *Agronomy Journal* **102**: 48-54.
- MILLER, R.E., LEI, X. and ULLREY, D.E. (1991) Trace elements in animal nutrition, in: MORTVEDT, J.J. (Ed.) *Micronutrients in Agriculture*, 2<sup>nd</sup> Ed., pp. 593- 662 (Madison, WI, Soil Science Society of America).
- MILLNER, P.D. (2009) Bioaerosols associated with animal production systems. *Bioresource Technology* **100**: 5379-5385.
- MOHANNA, C. and NYS, Y. (1999) Effect of dietary zinc content and sources on the growth, body zinc deposition and retention, zinc excretion and immune response in chickens. *British Poultry Science* **46**: 108-114.
- MOHANNA, C., CARRE, B. and NYS, Y. (1999) Incidence of dietary viscosity on growth performance and zinc and manganese bioavailability in broilers. *Animal Feed Science and Technology* **77**: 255-266.
- MOORE, P.A., DANIEL, T.C., GILMOUR, J.T., SHREVE, B.R., EDWARDS, D.R. and WOOD, B.H. (1998) Decreasing metal runoff from poultry litter with aluminium sulfate. *Journal of Environmental Quality* **27**: 92-99.
- MOORE, P.A., DANIEL, T.C., EDWARDS, D.R. and MILLER, D.M. (1996) Evaluation of chemical amendments to reduce ammonia volatilisation from poultry litter. *Poultry Science* **75**: 315–320.
- MOORE, P.A., DANIEL, T.C., SHARPLEY, A.N. and WOOD, C.W. (1995) Poultry manure management - environmentally sound options. *Journal of Soil and Water Conservation* **50**: 321-327.
- MOORE Jr, P.A., JOERN, B.C., EDWARDS, D.R., WOOD, C.W. and DANIEL, T.C. (2006) Effects of manure amendments on environmental and production problems, in: RICE, J.M., CALDWELL, D.F., & HUMENIK, F.J. (Eds) *Animal Agriculture and the Environment: National Center for Manure and Animal Waste Management White Papers*, Publication No 913C0306, pp. 1-40 (St. Joseph, MI, ASABE).
- MORRISON, J.L. (1969) Distribution of arsenic from poultry litter in broiler chickens, soil and crops. *Journal of Agricultural and Food Chemistry* **17**: 1288-1290.
- NAHM, K.H. (2002) Efficient feed nutrient utilisation to reduce pollutants in poultry and swine manure. *Critical reviews in Environmental Science and Technology* **32**: 1-16.
- NAHM, K.H. (2005) Environmental effects of chemical additives used in poultry litter and swine manure. *Critical reviews in Environmental Science and Technology* **35**: 487-513.
- NICHOLSON, F.A., CHAMBERS, B.J., WILLIAMS, J.R. and UNWIN, R.J. (1999) Heavy metal contents of livestock feeds and animal manures in England and Wales. *Bioresource Technology* **23**: 23-31.
- NOVAK, J.M., SZOGI, A.A. and WATTS, D.W. (2008) Copper and Zinc accumulations in sandy soils and constructed wetlands receiving pig manure effluent applications, in: SCHLEGEL, P., DUROSOY, S., & JONGBLOED, A.W. (Eds) *Trace Elements in Animal Production Systems*, pp. 45-54 (The Netherlands, Wageningen Academic Publishers).
- NRC (1994) Nutrient Requirements of Poultry: Ninth Revised Edition. National Research Council. Washington, D.C., National Academy Press.
- OSU (2010) Poultry Pest Management. Bulletin 853. Available at: [http://ohioline.osu.edu/b853/b853\\_5.html](http://ohioline.osu.edu/b853/b853_5.html).
- PIANZ (2006) Poultry Industry Association of New Zealand. Available at: <http://www.pianz.org.nz/>.
- POULSEN, H.D. (1998) Zinc and copper as feed additives, growth factors or unwanted environmental factors. *Journal of Animal Feed Science* **7**: 135-142.
- POWER, J.F. and DICK, W.A. (2000) Land Application of Agricultural, Industrial, and Municipal By-products. *Soil Science Society of America Inc.*, Madison, WI.
- POWERS, W. and ANGEL, R. (2008) A review of the capacity for nutritional strategies to address environmental challenges in poultry production. *Poultry Science* **87**: 1929-1938.
- PREUSCH, P.L., ADLER, P.R., SIKORA, L.J. and TWORKOSKI, T.J. (2002) Nitrogen and phosphorus availability in composted and uncomposted poultry litter. *Journal of Environmental Quality* **31**: 2051-2057.

- RIBAUDO, M.O., GOLLEHON, N.R. and AGAPOFF, J.** (2003) Land application of manure by animal feeding operations: Is more land needed? *Journal of Soil and Water Conservation* **58**: 30-38.
- RIRDC** (2009) Rural Industries Research and Development Corporation, Australia. Available at: <http://www.rirdc.gov.au/>.
- RIGBY, C.E., PETTIT, J.R., BAKER, M.F., BENTLY, A.H., SALOMONS, M.O. and LIAR, H.** (1980a) Flock infection and transport crates as sources of salmonellae in broiler chickens and carcasses. *Canadian Journal of Comparative Medicine* **44**: 328-337.
- RIGBY, C.E., PETTIT, J.R., BAKER, M.F., BENTLY, A.H., SALOMONS, M.O. and LIAR, H.** (1980b) Sources of salmonellae in an uninfected commercially-processed broiler flock. *Canadian Journal of Comparative Medicine* **44**: 267-274.
- RITTER, W.F.** (2000) Potential impact of land application of by-products on ground and surface water quality, in: DICK, W.A. (Ed.) *Land Application of Agricultural, Industrial, and Municipal By-products*, pp. 263-288 (Madison, WI, Soil Science Society of America).
- RITZ, C.W., FAIRCHILD, B.D. and LACY, M.P.** (2004) Implications of ammonia production and emissions from commercial poultry facilities: A review. *Journal of Applied Poultry Research* **13**: 684-692.
- ROTHROCK, M.J., JR., COOK, K.L., LOVANH, N., WARREN, J.G. and SISTANI, K.** (2008a) Development of a quantitative real-time PCR assay to target a novel group of ammonia producing bacteria found in poultry litter. *Poultry Science* **87**: 1058-1067.
- ROTHROCK, M.J., JR., COOK, K.L., WARREN, J.G. and SISTANI, K.** (2008b) The effect of alum addition on poultry litter microbial communities. *Poultry Science* **87**: 1493-1503.
- SATO, S., SOLOMON, D., HYLAND, C., KETTERINGS, Q.M. and LEHMANN, J.** (2005) Phosphorus speciation in manure and manure-amended soils using XANES spectroscopy. *Environmental Science and Technology* **39**: 7485-7491.
- SCHEFFERLE, H.E.** (1965) The decomposition of uric acid in built up poultry litter. *Journal of Applied Bacteriology* **28**: 412.
- SCHIFFMAN, S. and WILLIAMS, M.** (2005) Science of odor as a potential health issue. *Journal of Environmental Quality* **34**: 129-138.
- SCHOMBERG, H.H., ENDALE, D.M., JENKINS, M.B., SHARPE, R.R., FISHER, D.S., CABRERA, M. L. and MCCRACKEN, D.V.** (2009) Soil test nutrient changes induced by poultry litter under conventional tillage and no-tillage. *Soil Science Society of America Journal* **73**: 154-163.
- SELLE, P.H. and RAVINDRAN, V.** (2007) Microbial phytase in poultry nutrition. *Animal Feed Science and Technology* **135**: 1-41.
- SHARMA, M.C., JOSHI, C. and SAXEN, N.** (2005) Mineral toxicity in livestock: A review. *Indian journal of animal sciences* **75**: 753-764.
- SHARPLEY, A., MEISINGER, J.J., BREEUWSMA, A., SIMS, J.T., DANIEL, T.C. and SCHEPERS, J. S.** (1998) Impact of animal manure management on ground and surface water quality, in: HATFIELD, J.L. & STEWART, B.A. (Eds) *Animal Waste Utilisation: Effective Use of Manure as a Soil Resource*, pp.173-242 (Chelsea, MI, Ann Arbor Press).
- SHARPLEY, A.N. and SMITH, S.J.** (1995) Nitrogen and phosphorus forms in soils receiving manure. *Soil science* **159**: 253-258.
- SHARPLEY, A.N., HERRON, S. and DANIEL T.** (2007) Overcoming the challenges of phosphorus-based management challenges in poultry farming. *Journal of Soil and Water Conservation* **58**: 30-38.
- SHARPLEY, A.N., MCDOWELL, R.W. and KLEINMAN, P.J.A.** (2004) Amounts, forms, and solubility of phosphorus in soils receiving manure. *Soil science society of America journal* **68**: 2048-2057.
- SILVERSIDES, F.G. and HRUBY, M.** (2009) Feed formulation using phytase in laying hen diets. *Journal of applied poultry research* **18**:15-22.
- SIMS, J.T.** (1987) Agronomic evaluation of poultry manure as a nitrogen source for conventional and no-tillage corn. *Agronomy Journal* **79**: 563-570
- SIMS, J.T. and WOLF, D.C.** (1994) Poultry Waste Management: Agricultural and Environmental Issues. *Advances in Agronomy* **52**: 1-83.
- SIMS, J.T.** (1997) Agricultural and environmental issues in the management of poultry wastes: recent innovations and long-term challenges. *Agricultural uses of by-products and wastes. ACS Symposium series* **668**: 72-90.
- SIMS, J.T., BERGSTRÖM, L., BOWMAN, B.T. and OENEMA, O.** (2005) Nutrient management for intensive animal agriculture: policies and practices for sustainability. *Soil Use and Management* **21**: 141-151.
- SINGER, R.S. and HOFACRE, C.L.** (2006) Potential impacts of antibiotic use in poultry production. *Avian Diseases* **50**: 161-172.
- SISTANI, K.R., BRINK, G.E., MCGOWEN, S.L., ROWE, D.E. and OLDHAM, J.L.** (2003) Characterization of broiler cake litter, the by-products of two management practices. *Bioresource Technology* **90**: 27-32.

- SISTANI, K.R., ROWE, D.E., MILES, D.M. and MAY, J.D.** (2001) Effects of drying method and rearing temperature on broiler manure nutrient content. *Communications in Soil Science and Plant Analysis* **32**: 2307-2316.
- SMITH, L.W. and FRIES, G.F.** (1973) Dehydrated poultry manure as a crude protein supplement for lactating cows. *Journal of Dairy Science* **56**: 668-672.
- STUEDEMANN, J.A., WILKINSON, S.R., WILLIAMS, D.J., CIORDIA, H., ERNST, J.V., JACKSON, W.A. and JONES, J.B. JR.** (1975) Long-term broiler litter fertilization of tall fescue pastures and health and performance of beef cow. Managing livestock wastes. *Proceedings of 3<sup>rd</sup> International Symposium Livestock Wastes*, Urbana-Champaign, IL, pp. 264-268.
- SWAIN, B.K. and SUNDARAM, R.N.S.** (2000) Effect of different types of litter material for rearing broilers. *British Poultry Science* **41**: 261-262.
- SZOGLI, A.A. and VANOTTI, M.B.** (2009) Prospects for phosphorus recovery from poultry litter. *Bioresource Technology* **100**: 5461-5465.
- TASISTRO, A.S., KISSEL, D.E. and BUSH, P.B.** (2004) Spatial variability of broiler litter composition in a chicken house. *Journal of applied poultry research* **13**: 29-43.
- TERZICH, M., POPE, M.J., CHERRY, T.E. and HOLLINGER, J.** (2000) Survey of pathogens in poultry litter in the United States. *Journal of Applied Poultry Research* **9**: 287-291.
- TEWOLDE, H., ARMSTRONG, S., WAY, T.R., ROWE, D.E. and SISTANI, K.R.** (2009) Cotton response to poultry litter applied by subsurface banding relative to surface broadcasting. *Soil Science Society of America Journal* **73**: 384-389.
- TEWOLDE, H., SISTANI, K.R. and ROWE, D.E.** (2005) Broiler litter as a micronutrient source for cotton: concentrations in plant parts. *Journal of Environmental Quality* **34**: 1697-706.
- TOOR, G.S. and HUNGER, B.E.** (2009) Phosphorus and trace metal dynamics in soils amended with poultry litter and granulates. *Soil Use and Management* **25**: 409-418.
- TOOR, G.S., HUNGER, S., PEAK, J.D., SIMS J.T. and SPARKS, D.L.** (2006) Advances in the characterization of phosphorus in organic wastes: environmental and agronomic applications. *Advances in agronomy* **89**: 1-72.
- TUFFT, L.S. and NOCKELS, C.F.** (1991) The effects of stress, *Escherichia coli*, dietary ethylene-diaminetetraacetic acid, and their interaction on tissue trace-elements in chicks. *Poultry Science* **70**: 2439-2449.
- TURNELL, J.R., FAULKNER, R.D. and HINCH, G.N.** (2007) Recent advances in Australian broiler litter utilisation. *World's Poultry Science Journal* **63**: 223-231.
- TURNER, B.L. and LEYTEM, A.B.** (2004) Phosphorus compounds in sequential extracts of animal manures: Chemical speciation and novel fractionation procedure. *Environmental Science and Technology* **38**: 6101-6108.
- USDA** (2008) Agricultural Statistics 2008. National Agricultural Statistics Service. Washington, DC, United States Government Printing Office.
- VAN RYSSEN, J.B.J.** (2008) Trace elements in poultry litter: prevalence and risks, in: SCHLEGEL, P., DUROSOY, S. & JONGBLOED, A.W. (Eds) *Trace Elements in Animal Production Systems*, pp. 101-113 (The Netherlands, Wageningen Academic Publishers).
- WALKER, J.D., ANEJA, V.P. and DICKEY, D.A.** (2000a) Atmospheric transport and wet deposition of ammonium in North Carolina. *Atmospheric Environment* **34**: 3407-3418.
- WALKER, J., NELSON, D. and ANEJA, V.P.** (2000b) Trends in ammonium concentration in precipitation and atmospheric ammonia emissions at a coastal plain site in North Carolina, U.S.A. *Environmental Science and Technology* **34**: 3527-3534.
- WILKINSON, S.R.** (1997) Response of Tifway 2 Bermuda grass to fresh or composted broiler litter containing boric acid-treated paper bedding. *Communications in Soil Science and Plant Analysis* **28**: 259-279.
- WILLIAMS, C.M., BARKER, J.C. and SIMS, J.T.** (1999) Management and utilisation of poultry wastes. *Reviews of Environmental Contamination and Toxicology* **162**: 105-157.
- WORLEY, J.W., RISSE, L.M., CABRERA, M.L. and NOLAN, M.P.** (1999) Bedding for broiler chickens: Two alternative systems. *Applied Engineering in Agriculture* **15**: 687-692.