

## RESEARCH ARTICLE

# Extrusion cooking with glucose supplementation of fumonisin-contaminated corn grits protects against nephrotoxicity and disrupted sphingolipid metabolism in rats

Kenneth A. Voss<sup>1</sup>, Ronald T. Riley<sup>1</sup>, Lauren S. Jackson<sup>2</sup>, Joseph E. Jablonski<sup>2</sup>, Andreia Bianchini<sup>3</sup>, Lloyd B. Bullerman<sup>3</sup>, Milford A. Hanna<sup>3</sup> and Dojin Ryu<sup>4</sup>

<sup>1</sup>USDA, Agricultural Research Service, Russell Research Center, Athens, GA, USA

<sup>2</sup>FDA National Center for Food Safety and Technology, Summit-Argo, IL, USA

<sup>3</sup>Department of Food Science and Technology, University of Nebraska, Lincoln, NE, USA

<sup>4</sup>Department of Nutrition and Food Sciences, Texas Woman's University, Denton, TX, USA

**Scope:** Fumonisin B<sub>1</sub> (FB<sub>1</sub>) is a mycotoxin found in maize and maize-based foods. It causes animal diseases and is a suspected risk factor for cancer and birth defects in humans. Extrusion cooking reduces FB<sub>1</sub> concentrations in maize however toxicity caused by unknown degradation or FB<sub>1</sub>-matrix reaction products might persist.

**Methods and results:** To test the efficacy of extrusion to reduce FB<sub>1</sub> toxicity, *Fusarium verticillioides* fermented corn (= maize) grits (Batch-1= 9.7 ppm FB<sub>1</sub>; Batch-2= 50 ppm FB<sub>1</sub>) were extruded without (Batch-1E; Batch-2E) or with 10% glucose supplementation (Batch-1EG; Batch-2EG). FB<sub>1</sub> concentrations were reduced 64% (Batch-2E) to 94% (Batch-1EG) after cooking. When the uncooked and processed grits were fed (50% w/w in rodent chow) to rats for up to 8 weeks, FB<sub>1</sub> intakes averaged 354, 103, and 25.1  $\mu\text{g}/\text{kg}$  body weight/day for Batch-1, Batch-1E and Batch-1EG and 1804, 698, and 222  $\mu\text{g}/\text{kg}$  body weight/day for the Batch-2, Batch-2E and Batch-2EG, respectively. Nephrotoxicity including apoptotic lesions and elevated sphingoid base concentrations decreased in a dose-dependent manner in groups fed Batch-1, Batch-1E, Batch-2, Batch-2E, or Batch-2EG and was absent in the Batch-1EG group. **Conclusion:** Extrusion cooking, especially with glucose supplementation, is potentially useful to reduce FB<sub>1</sub> concentrations and toxicity of FB<sub>1</sub>-contaminated maize.

Received: January 28, 2011

Revised: March 15, 2011

Accepted: March 21, 2011

## Keywords:

Extrusion cooking / Fumonisin / In vivo toxicity / Sphinganine / Sphinganine 1-phosphate

## 1 Introduction

Fumonisin is a mycotoxin produced by *Fusarium verticillioides*, *F. proliferatum*, other *Fusarium* and some species of

the *Aspergillus niger* complex [1–3]. They are common contaminants of maize and maize-based products but have also been found in a variety of other foodstuffs [3]. Fumonisin B<sub>1</sub> (FB<sub>1</sub>) is the most predominant and thoroughly studied congener [1, 2]. Its toxic and carcinogenic effects in animals include leukoencephalomalacia in *Equidae*, pulmonary edema in swine, and hepato- and nephrotoxicities in a variety of piscine, avian and mammalian species [2, 4, 5]. FB<sub>1</sub> is a liver and kidney carcinogen in rodents [6, 7]. The impact of FB<sub>1</sub> or other fumonisins on human health is unknown although evidence suggests that FB<sub>1</sub> is a risk factor for birth (neural tube) defects or cancer in populations consuming large amounts of contaminated maize [1, 2, 8, 9].

**Correspondence:** Dr. Kenneth A. Voss, USDA Agricultural Research Service, Toxicology & Mycotoxin Research Unit, 950 College Station Road, Athens, GA 30605, USA

**E-mail:** Ken.Voss@ars.usda.gov

**Fax:** +1-706-546-3116

**Abbreviations:** FB<sub>1</sub>, Fumonisin B<sub>1</sub>; deoxy Sa, 1-deoxysphinganine; NDF-FB<sub>1</sub>, N-(1-deoxy-D-fructosyl)-FB<sub>1</sub>; Sa, sphinganine; Sa 1-P, sphinganine 1-phosphate; So, sphingosine

FB<sub>1</sub> is generally stable during cooking and therefore efforts to reduce its concentrations in foods have yielded mixed results [10, 11]. Extrusion cooking is a process combining high temperature and pressure that has been shown to significantly reduce fumonisin concentrations in maize. Greater reductions are achieved when a reducing sugar is added to the mixture prior to processing [11–13]. However, analytical results alone are insufficient to determine whether cooking reduces toxicity as unknown thermal decomposition or other reaction products, including difficult to detect matrix-associated or bound fumonisins [14, 15] might contribute to toxicity.

In earlier studies, FB<sub>1</sub> concentrations in spiked (30 ppm) or *F. verticillioides*-fermented (33–48 ppm) grits were reduced 10–28% by cooking alone and 75–85% by cooking with 10% glucose supplementation using a single-screw extruder [13]. Extrusion with glucose partially reversed toxicity of the less contaminated fermented grits when they were fed to rats, suggesting that the process was potentially useful for reducing fumonisin exposure [16].

The purpose of this investigation was to determine the effectiveness of cooking *F. verticillioides*-fermented corn (= maize) grits by twin-screw extrusion to reduce FB<sub>1</sub> bioavailability and toxicity. The bioassay was more rigorous than an earlier one [16] in that it (i) tested the process on grits having FB<sub>1</sub> concentrations both below (9.7 ppm, Batch-1) and above (50 ppm, Batch-2) that for which partial protection against toxicity was demonstrated, (ii) evaluated toxicity over a longer, 8 wk exposure period, (iii) included hematology and clinical chemistry indicators of kidney and liver dysfunction as experimental endpoints and (iv) utilized changes in target organ sphingoid base concentrations as a biomarker of exposure.

## 2 Materials and methods

### 2.1 Grits

Contaminated corn grits were prepared as previously reported [13]. Briefly, two batches were fermented with *F. verticillioides* M-2552 until contamination levels of about 10 (Batch-1) or 50 (Batch-2) ppm FB<sub>1</sub> were achieved.

### 2.2 Extrusion

Extrusion conditions are described in detail by Jackson et al. [17]. Briefly, Batch-1 and Batch-2 grits (20% moisture content, dry weight basis) were extruded without (Batch-1E, Batch-2E) or with 10% food-grade glucose (ADM Corn Processing, Decatur, IL, USA) supplementation (Batch-1EG, Batch-2EG) using a model CTSE-V twin-screw apparatus (C.W. Brabender, South Hackensack, NJ, USA) with conical mixing configuration at 160°C, 40 rpm and 60 g/min feed rate. A portion of the uncontaminated grits (20% moisture

content, dry weight basis) was similarly processed without adding glucose. The extruded grit preparations were ground and they and the uncooked grits were stored frozen. FB<sub>1</sub> and hydrolyzed FB<sub>1</sub> (HFB<sub>1</sub>) concentrations in the grits before and after processing were quantified by HPLC with fluorescence detection following derivatization with *o*-phthalaldehyde (Sigma, St. Louis, MO, USA). The FB<sub>1</sub>-glucose reaction product *N*-(1-deoxy-*D*-fructos-1-yl)-FB<sub>1</sub> (NDF-FB<sub>1</sub>) [18] was quantified by LC-MS analysis [13].

### 2.3 Animals

Male Sprague–Dawley rats (Harlan Industries, Indianapolis, IN, USA), 3 wk of age were acclimated for 2 wk and then randomly assigned to eight groups ( $n = 10$ /group) having mean body weights of  $73 \pm 3.1$  g (SD) to  $74 \pm 3.7$  g. The animals were individually housed in stainless-steel, wire-mesh cages in an environmentally controlled room having a 12-h light/dark cycle. Fresh tap water and the appropriate control or test diet were provided ad libitum.

### 2.4 Diets

Animals were fed standard rodent diet (TD2019, Teklad, Madison, WI, USA) during acclimation. The basal feed (Certified Teklad TD.09.119, <0.1 ppm FB<sub>1</sub>) used to mix the control and test diets was formulated to ensure adequate amounts of protein, vitamins, lipotropes and other nutrients. Basal feed and the grits preparations were mixed (50% w/w) using a Patterson Kelley V blender with intensifier bar. FB<sub>1</sub> concentrations of each diet (Table 1) were calculated on the basis of the FB<sub>1</sub> concentrations in the raw or processed grits, which in turn were determined by chemical analysis [17]. The diets were stored frozen and dispensed fresh each wk.

### 2.5 Bioassay

Procedures were approved by the USDA-ARS Russell Research Center Institutional Animal Care and Use Committee. The groups were fed uncooked uncontaminated grits (uncooked control group), extruded uncontaminated grits (extrusion control group) or one of the test diets (Table 1). The animals were observed at least once per day. Body weights and food consumptions were measured weekly. After 3 and 8 wk of feeding, five animals/group were fasted overnight in metabolism cages for collection of urine samples. Thereafter, they were lightly anaesthetized (Isoflurane USP, Isoflo<sup>®</sup>, Abbott Laboratories, Chicago, IL, USA) and blood collected (periorbital plexus) for hematology and serum chemistry determinations. Following blood collection, the animals were euthanized (CO<sub>2</sub> inhalation) and examined by necropsy. The kidneys, liver, brain, heart,

lungs with trachea, testes, adrenals and spleen were weighed. Representative specimens of these and other major organs were fixed in 10% neutral buffered formalin. Additional samples of liver and kidney were collected and frozen ( $-80^{\circ}\text{C}$ ) for sphingolipid analyses.

### 2.5.1 Clinical laboratory studies

Hematology, serum chemistry and urinalysis were conducted on all animals. Hematology profiles were obtained using a Bayer Advia 120 Hematology Analyzer and commercial reagents (Siemens Healthcare Diagnostics, Deerfield, IL, USA). Profiles included erythrocyte count, hemoglobin, hematocrit, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, reticulocyte count, total and differential leukocyte counts, platelet count and mean platelet volume. Clinical chemistry procedures were automated (Hitachi Model 912) and done using commercial reagents (Roche Diagnostics Corporation, Indianapolis, IN, USA). Serum enzyme determinations were aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, lactate dehydrogenase and  $\gamma$ -glutamyl transpeptidase. Other parameters were total protein, albumin, triglycerides, cholesterol, glucose, total bilirubin, blood urea nitrogen, creatinine and electrolytes (sodium, potassium, calcium, bicarbonate, chloride phosphorus and anion gap). Urine samples were screened for color, turbidity, pH, protein, glucose, ketones, bilirubin and blood using Siemens Multistix<sup>®</sup> (Siemens Healthcare Diagnostics, Tarrytown, NY, USA) as well as for specific gravity (TS refractometer) and by microscopic examination.

### 2.5.2 Histopathology

Hematoxylin and eosin (H & E)-stained kidney specimens from all animals were examined in a random order and without knowledge of the animals' group assignments. Lesions consistent with fumonisin exposure [4] were scored on a scale of 0 = no effect; 1 = few apoptotic cells in an unremarkable parenchyma indicating a minimal change; 2 = mild, generalized effect of FB<sub>1</sub> exposure; 3 = more obvious, moderate effect and 4 = severe effect with overt necrosis and fibrosis [16]. Hematoxylin and eosin-stained liver (all groups, both time periods) and heart, testes, spleen, adrenals and lung (all groups, 8 wk only) specimens also were examined.

### 2.5.3 Tissue sphingoid base profiles

The concentrations of free sphinganine (Sa), sphingosine (So), 1-deoxysphinganine (deoxy Sa), sphinganine 1-phosphate (Sa 1-P) and sphingosine 1-P (So 1-P) in the kidneys of all animals and in the livers of the control, extruded control,

Batch-2E and Batch-2EG groups were determined using a previously reported LC-MS method [19] as modified for deoxy Sa [20].

### 2.5.4 Statistical analysis

Statistical procedures were done using SigmaStat software (Jandel Scientific, San Rafael, CA, USA). Each test group was compared with the control groups and the two other groups fed grits preparations made from the same batch. Continuous, homogenous data were analyzed by ANOVA and differences among groups identified post hoc using Student–Newman–Keul's method. Nonparametric data including kidney scores were analyzed using the Kruskal–Wallis Test and Fisher's exact test was used to evaluate pathology incidence data. All tests were two-tailed and significance was determined at  $p < 0.05$ .

## 3 Results

### 3.1 Extrusion and diets

Analytical procedures and results for the effects of twin-screw extrusion cooking on FB<sub>1</sub> in maize grits are reported in detail elsewhere [17]. Briefly, extrusion reduced FB<sub>1</sub> concentrations by 72 and 64% for Batch-1E and Batch-2E, respectively (Table 1). Glucose supplementation enhanced reductions so that FB<sub>1</sub> concentrations were reduced by 94% for Batch-1EG and 89% for Batch-2EG. As a result of dilution (50% w/w) with basal feed, the FB<sub>1</sub> concentrations ranged from 0.3 to 4.9 ppm for the lower level Batch-1 series (Batch-1, Batch-1E and Batch-1EG) and 2.9 to 25 ppm for the higher level Batch-2 series (Batch-2, Batch-2E and Batch-2EG) of experimental diets. Uncooked and extruded control diets contained  $< 0.1$  ppm FB<sub>1</sub>. Processing did not affect HFB<sub>1</sub> concentrations, which were low ( $< 0.5$  ppm) in all diets. Minor amounts of the glucose reaction product, NDF-FB<sub>1</sub>, were present in the uncooked control diet. Extrusion lowered its dietary concentration from 0.4 ppm (Batch-1) to  $< 0.1$  ppm (Batch-1E) and from 2.1 ppm (Batch-2) to 0.5 ppm (Batch-2E). Extrusion with glucose supplementation did not affect NDF-FB<sub>1</sub> in Batch-1E but increased its amount in Batch-2E: concentrations were 0.5 ppm in the Batch-1EG and 3.9 ppm in Batch-2EG diets.

### 3.2 Body weight and food consumption

No differences in body weight were found between the uncooked control and other groups. Body weights of the extruded control group were significantly higher than in the Batch-2 group at wk 2 and the Batch-2, Batch-2E and Batch-2EG groups at wk 3. No other differences were found. Group mean weights (g) at wk 8 were as follows: uncooked

control =  $380 \pm 23$  (SD), extruded control =  $383 \pm 23$ , Batch-1 =  $364 \pm 31$ , Batch-1E =  $366 \pm 26$ , Batch-1EG =  $377 \pm 38.0$ , Batch-2 =  $364 \pm 23$ , Batch-2E =  $373 \pm 25$  and Batch-2EG =  $359 \pm 27$ .

Weekly food consumption of the Batch-1 and Batch-2 groups was significantly lower than in the two control groups through wk 3 (data not shown). Food consumption of rats fed Batch-1 also was significantly less when compared with the Batch-1E and Batch-1EG groups and reduced during weeks 1 and 2 in rats fed Batch-2 compared with those fed Batch-2E or Batch-2EG. No significant differences were found after wk 3.

### 3.3 FB<sub>1</sub> intake

Daily FB<sub>1</sub> intakes ( $\mu\text{g}$  FB<sub>1</sub>/kg body weight/day) were calculated using the formula: ((weekly food consumption (g)  $\times$  FB<sub>1</sub> concentration ( $\mu\text{g}/\text{g}$ ))/average body weight (kg) during the week)/7. For Batch-1 (Table 1), extrusion reduced daily FB<sub>1</sub> intake about 70% and extrusion with glucose reduced FB<sub>1</sub> daily intake >90%. Reductions of 60–88%

**Table 1.** FB<sub>1</sub> concentrations (ppm) of the diets prepared from two batches of *F. verticillioides* fermented grits that were uncooked, extruded (-E), or extruded with 10% glucose supplementation (-EG) for 3 or 8 wks and estimated daily FB<sub>1</sub> intake of rats fed the diets for 3 or 8 wk

| Diet                                       | FB <sub>1</sub> <sup>a)</sup><br>(ppm) | Estimated daily FB <sub>1</sub> intake during weeks <sup>b)</sup> |                              |                                |
|--------------------------------------------|----------------------------------------|-------------------------------------------------------------------|------------------------------|--------------------------------|
|                                            |                                        | 1–3 ( <i>n</i> = 10)                                              | 4–8 ( <i>n</i> = 5)          | 1–8 ( <i>n</i> = 5)            |
| <i>Fermented Grits Low FB<sub>1</sub></i>  |                                        |                                                                   |                              |                                |
| Batch-1                                    | 4.9                                    | 468 <sup>c</sup> C<br>± 27.0                                      | 288 <sup>d</sup> C<br>± 34.2 | 354 <sup>d</sup> C<br>± 150    |
| Batch-1E                                   | 1.4                                    | 143 <sup>c</sup> D<br>± 28.7                                      | 80 <sup>d</sup> D<br>± 6.9   | 103 <sup>d</sup> D<br>± 34.4   |
| Batch-1EG                                  | 0.3                                    | 33.6 <sup>c</sup> E<br>± 7.7                                      | 19.3 <sup>d</sup> E<br>± 1.7 | 25.1 <sup>cd</sup> D<br>± 10.5 |
| <i>Fermented Grits High FB<sub>1</sub></i> |                                        |                                                                   |                              |                                |
| Batch-2                                    | 25                                     | 2419 <sup>c</sup> C<br>± 391                                      | 1439 <sup>d</sup> C<br>± 212 | 1804 <sup>d</sup> C<br>± 560   |
| Batch-2E                                   | 9.0                                    | 923 <sup>c</sup> D<br>± 38.2                                      | 563 <sup>d</sup> D<br>± 57.3 | 698 <sup>d</sup> D<br>± 201    |
| Batch-2EG                                  | 2.9                                    | 298 <sup>c</sup> E<br>± 50.2                                      | 179 <sup>d</sup> E<br>± 22.5 | 222 <sup>d</sup> D<br>± 63.7   |

a) Dietary concentration (wet weight basis) as determined by chemical analysis of the grits [17]; control and extruded control diets, <0.1 ppm.

b) Calculated from body weight, food consumption and FB<sub>1</sub> concentration data, see text for formula; 1–8 wk results do not include animals necropsied after week 3. cdValues in ROWS not sharing lower-case superscript are significantly different,  $p < 0.05$ . CDEBatch-1 and Batch-2 series groups not sharing upper-case superscripts (in COLUMNS) are significantly different,  $p < 0.05$ .

were achieved by extrusion and extrusion with glucose, respectively, of Batch-2. Exposures ( $\mu\text{g}$  FB<sub>1</sub>/kg body weight/day) declined as the study progressed. Daily intakes of all groups were reduced significantly (39–44% lower) during weeks 4–8 compared with weeks 1–3 and, with one exception (Batch-1EG), to the entire 8-wk feeding period.

### 3.4 Clinical laboratory studies

There were no exposure-related hematology or serum chemistry findings. Urinalysis revealed signs of kidney dysfunction in the animals fed Batch-2 or Batch-2E, which included traces of blood and increased numbers of (presumably sloughed kidney tubule) cells at wk 3 (data not shown). These findings were persistent but markedly less severe at wk 8.

### 3.5 Necropsy and organ weights

There were no treatment-related gross lesions. Absolute kidney weights (g) of animals fed Batch-1, Batch-1E or Batch-1EG were not different from those fed the uncooked control diet (Table 2). Absolute kidney weights of the Batch-1 group were however significantly lower than those of the extruded controls at wk 3. Kidney weights of the Batch-2 group were significantly reduced relative to both control groups and the Batch-2EG group at wks 3 and 8.

Relative kidney weights (% body weight) (Table 2) of the Batch-1 group were significantly lower than those of the two control, Batch-1E and Batch-1EG groups at week 3. Relative kidney weights of rats fed Batch-2 were decreased significantly at both wks 3 and 8 when compared with the controls. At wk 3, relative kidney weights of the Batch-2E and Batch-2EG groups were not different from those of the Batch-2 or the uncooked control groups but were less than those of the extruded control group. At wk 8, relative kidney weights of animals fed Batch-2 or Batch-2E were significantly lower than those fed the Batch-2EG or the two control diets.

### 3.6 Histopathology

Treatment-related effects were limited to the kidneys. Extrusion and, to a greater extent, extrusion with glucose reduced nephrotoxicity although protection was complete only in rats fed Batch-1EG. Specifically, the number of animals ( $n = 5/\text{group}$ ) in the Batch-1 series exhibiting apoptotic proximal tubule cells or apoptosis together with other features consistent with FB<sub>1</sub> nephropathy at wk 3 were as follows: Batch-1 = 5; Batch-1E = 4; Batch-1EG = 1 and the two control groups  $\leq 1$ . The group incidences of rats with lesions at wk 8 were as follows: Batch-1 = 4; Batch-1E = 2; Batch-1EG and the control groups = 0. Mean histopathology severity scores (Fig. 1) for the Batch-1 series decreased at


**Table 2.** Absolute and relative (% body weight) kidney weights of rats fed diets prepared from uncontaminated grits or two batches of *F. verticillioides* fermented grits that were uncooked (-E), or extruded with 10% glucose supplementation (-EG) for 3 or 8 wk

| Diet                                       | Week 3                             |                                 | Week 8                          |                   |
|--------------------------------------------|------------------------------------|---------------------------------|---------------------------------|-------------------|
|                                            | Absolute (g)                       | % Body weight                   | Absolute (g)                    | % Body weight     |
| <i>Controls</i>                            |                                    |                                 |                                 |                   |
| Uncooked                                   | 2.07 <sup>ab</sup> / <sup>AB</sup> | 0.81 <sup>a</sup> <sup>AB</sup> | 2.40 <sup>a</sup> <sup>AB</sup> | 0.66 <sup>A</sup> |
|                                            | ± 0.08                             | ± 0.04                          | ± 0.19                          | ± 0.03            |
| Extruded                                   | 2.25 <sup>a</sup> <sup>A</sup>     | 0.87 <sup>a</sup> <sup>A</sup>  | 2.51 <sup>a</sup> <sup>B</sup>  | 0.69 <sup>A</sup> |
|                                            | ± 0.23                             | ± 0.05<br>(4)                   | ± 0.19                          | ± 0.02            |
| <i>Fermented Grits Low FB<sub>1</sub></i>  |                                    |                                 |                                 |                   |
| Batch-1                                    | 1.79 <sup>b</sup>                  | 0.72 <sup>b</sup>               | 2.26 <sup>a</sup>               | 0.65              |
|                                            | ± 0.09                             | ± 0.02                          | ± 0.08                          | ± 0.05            |
| Batch-1E                                   | 2.06 <sup>ab</sup>                 | 0.82 <sup>a</sup>               | 2.49 <sup>a</sup>               | 0.71              |
|                                            | ± 0.17                             | ± 0.05<br>(4)                   | ± 0.31                          | ± 0.04            |
| Batch-1EG                                  | 2.11 <sup>ab</sup>                 | 0.87 <sup>a</sup>               | 2.54 <sup>a</sup>               | 0.71              |
|                                            | ± 0.27                             | ± 0.08<br>(4)                   | ± 0.37                          | ± 0.09            |
| <i>Fermented Grits High FB<sub>1</sub></i> |                                    |                                 |                                 |                   |
| Batch-2                                    | 1.73 <sup>C</sup>                  | 0.70 <sup>C</sup>               | 2.03 <sup>C</sup>               | 0.58 <sup>B</sup> |
|                                            | ± 0.08                             | ± 0.03<br>(4)                   | ± 0.11                          | ± 0.04            |
| Batch-2E                                   | 1.83 <sup>BC</sup>                 | 0.76 <sup>BC</sup>              | 2.17 <sup>BC</sup>              | 0.61 <sup>B</sup> |
|                                            | ± 0.18                             | ± 0.06                          | ± 0.17                          | ± 0.04            |
| Batch-2EG                                  | 1.86 <sup>B</sup>                  | 0.75 <sup>BC</sup>              | 2.37 <sup>AB</sup>              | 0.69 <sup>A</sup> |
|                                            | ± 0.18                             | ± 0.07                          | ± 0.16                          | ± 0.04            |

Note: values are the mean ± SD; *n* = 5 animals/group unless indicated otherwise in parentheses. abc Control and Batch-1 series groups not sharing lower-case superscripts are significantly different, *p* < 0.05; ABC Control and Batch-2 series groups not sharing upper-case superscripts are significantly different, *p* < 0.05. Fumonisin B<sub>1</sub> concentrations of the diets are given in Table 1.

week 3 in the order Batch-1 > Batch-1E > Batch-1EG ≥ two control groups and at week 8 in the order Batch-1 > Batch-1E > Batch-1EG = two control groups. The number of animals (*n* = 5/group) in the Batch-2 series having nephropathy were as follows: Batch-2 and Batch-2E = 5; and Batch-2EG = 4 at wk 3 and Batch-2 = 5; Batch-2E = 3; and Batch-2EG = 2 at week 6. The severity of lesions in the Batch-2 series groups was greater than in their Batch-1 counterparts and the controls. Scores for the Batch-2EG group were none the less lower than those of the Batch-2 and Batch-2E groups at wk 3 and wk 8 (*p* < 0.05). Scores for the latter groups were equal at wk 3 while at wk 8, the mean severity score for the Batch-2E group was between those of the Batch-2 and Batch-2EG groups.

When average daily FB<sub>1</sub> intake during weeks 1–3 or 4–8 and corresponding pathology scores of the groups were



**Figure 1.** Histopathology scores for severity of renal lesions consistent with fumonisin exposure in rats fed diets containing (50% w/w) uncooked (uncooked) or extruded (extruded) uncontaminated maize grits, or two batches of uncooked (Batch-1, Batch-2) FB<sub>1</sub>-contaminated grits, contaminated grits after extrusion (Batch-1E, Batch-2E) or contaminated grits after extrusion with glucose supplementation (Batch-1EG, Batch-2EG) for 3 or 8 wk. Values indicate the group mean score, *n* = 5; error bars denote standard deviation. See text for explanation of scores. Control and Batch-1 series groups not sharing lower-case superscripts are significantly different and Control and Batch-2 series groups not sharing upper-case superscripts are significantly different, *p* < 0.05.

considered (Fig. 2), it was evident that the scores were dose dependent and, for all groups, both daily intakes and histopathology scores were lower at wk 8.

### 3.7 Sphingoid bases


Sphingolipid metabolism was disrupted in the kidneys of all animals fed uncooked Batch-1 and Batch-2 grits as indicated by significant increases in the concentrations of Sa, So, their 1-P metabolites (Table 3) and total Sa (Sa plus Sa 1-P) (Fig. 3). Lesser, but statistically significant increases also were found in the Batch-1E and Batch-2E groups. Concen-

trations of individual sphingoid bases and total Sa of the Batch-1EG group were not elevated. Sphingoid base concentrations of the Batch-2EG group tended to be lower than in the Batch-2E and Batch-2 groups although, due to relatively large standard deviations, statistical differences were not always demonstrated. Sphingoid base concentrations of the Batch-2EG group were in contrast up to 88-fold higher (Sa, wk 8) than in the control groups and 28-

(Sa, week 3) to 69-fold (total Sa, wk 8) higher than in the Batch-1EG group.

Total Sa concentrations in kidneys increased in a (FB<sub>1</sub> daily intake) dose-dependent manner (Fig. 4) but the magnitude of the changes was diminished at week 8. A high degree of correlation (Fig. 5) was found between total Sa and mean histopathology scores at both times. Furthermore, higher pathology scores were associated with lower total Sa concentrations at week 8 than at wk 3.

Sa, Sa 1-P, So and So 1-P in the livers of rats fed Batch-2 and Batch-2E were not elevated and deoxy Sa was not found.



**Figure 2.** Group mean histopathology scores (from Fig. 1) plotted against group mean daily FB<sub>1</sub> intakes as determined from weekly body weight, food consumption and diet FB<sub>1</sub> concentration data.


## 4 Discussion

The efficacy of extrusion cooking, with and without glucose supplementation, using a twin-screw extrusion configuration to reduce FB<sub>1</sub> concentrations and in vivo toxicity of contaminated grits was tested. The experiment complemented an earlier but less rigorous 3-wk rat feeding bioassay of fermented grits containing  $\geq 33$  ppm FB<sub>1</sub> after single-screw extrusion with or without glucose supplementation [13, 16]. As before, diet formulations were based solely on FB<sub>1</sub> concentrations. The lower amounts of FB<sub>2</sub>, FB<sub>3</sub>, HFB<sub>1</sub> and NDF-FB<sub>1</sub> present in the grits [17] were not considered because these fumonisin species were not toxic when evaluated by various rodent bioassays [21–24]. Furthermore, some evidence suggests FB<sub>2</sub> and FB<sub>3</sub> do not accumulate in

**Table 3.** Kidney sphinganine (Sa), Sa 1-phosphate (Sa 1-P), sphingosine (So) and So 1-phosphate (So 1-P) concentrations (nmol/g) of rats fed diets prepared from uncontaminated grits or two batches of *F. verticillioides* fermented grits that were uncooked, extruded (-E), or extruded with 10% glucose supplementation (-EG) for 3 or 8wk

| Diet                                       | Week 3 <sup>a)</sup>  |                     |                     |                     | Week 8              |                      |                     |                     |
|--------------------------------------------|-----------------------|---------------------|---------------------|---------------------|---------------------|----------------------|---------------------|---------------------|
|                                            | Sa                    | Sa 1-P              | So                  | So 1-P              | Sa                  | Sa 1-P               | So                  | So 1-P              |
| <i>Controls</i>                            |                       |                     |                     |                     |                     |                      |                     |                     |
| Uncooked                                   | 4.49 <sup>b,c</sup> B | 0.96 <sup>b</sup> B | 8.00 <sup>b</sup> B | 0.28 <sup>b</sup> B | 1.60 <sup>b</sup> B | <0.30 <sup>b</sup> B | 4.12 <sup>b</sup> B | 0.65 <sup>b</sup> B |
|                                            | ±3.51                 | ±0.81               | ±2.77               | ±0.16               | ±0.81               | ±0                   | ±1.96               | ±0.20               |
| Extruded                                   | 2.56 <sup>b</sup> B   | 0.74 <sup>b</sup> B | 7.93 <sup>b</sup> B | 0.25 <sup>b</sup> B | 1.50 <sup>b</sup> B | <0.30 <sup>b</sup> B | 4.14 <sup>b</sup> B | 0.58 <sup>b</sup> B |
|                                            | ±0.73                 | ±0.97               | ±1.89               | ±0.20               | ±0.22               | ±0                   | ±0.78               | ±0.16               |
| <i>Fermented Grits Low FB<sub>1</sub></i>  |                       |                     |                     |                     |                     |                      |                     |                     |
| Batch-1                                    | 316 <sup>d</sup>      | 306 <sup>d</sup>    | 33.9 <sup>c</sup>   | 18.7 <sup>d</sup>   | 120 <sup>d</sup>    | 40.7 <sup>d</sup>    | 17.4 <sup>c</sup>   | 9.35 <sup>d</sup>   |
|                                            | ±99.1                 | ±143                | ±7.84               | ±7.72               | ±38.2               | ±24.0                | ±4.09               | ±4.84               |
| Batch-1E                                   | 181 <sup>d</sup>      | 89.3 <sup>c</sup>   | 29.5 <sup>c</sup>   | 6.98 <sup>c</sup>   | 67.4 <sup>c</sup>   | 6.30 <sup>c</sup>    | 20.0 <sup>c</sup>   | 2.71 <sup>c</sup>   |
|                                            | ±96.6                 | ±71.1               | ±6.01               | ±4.73               | ±22.3               | ±6.57                | ±7.86               | ±1.56               |
| Batch-1EG                                  | 6.32 <sup>c</sup>     | 1.03 <sup>b</sup>   | 8.81 <sup>b</sup>   | 0.37 <sup>b</sup>   | 1.92 <sup>b</sup>   | <0.30 <sup>b</sup>   | 4.60 <sup>b</sup>   | 0.40 <sup>b</sup>   |
|                                            | ±4.14                 | ±1.28               | ±0.55               | ±0.34               | ±0.66               | ±0                   | ±0.76               | ±0.09               |
| <i>Fermented Grits High FB<sub>1</sub></i> |                       |                     |                     |                     |                     |                      |                     |                     |
| Batch-2                                    | 443 <sup>D</sup>      | 581 <sup>C</sup>    | 43.3 <sup>D</sup>   | 23.7 <sup>D</sup>   | 265 <sup>C</sup>    | 115 <sup>D</sup>     | 45.6 <sup>D</sup>   | 17.7 <sup>D</sup>   |
|                                            | ±74.7                 | ±141                | ±5.76               | ±5.65               | ±144                | ±23.1                | ±20.6               | ±5.28               |
| Batch-2E                                   | 275 <sup>C</sup>      | 402 <sup>C</sup>    | 28.7 <sup>C</sup>   | 19.2 <sup>D</sup>   | 175 <sup>C</sup>    | 96.5 <sup>D</sup>    | 26.3 <sup>C</sup>   | 18.1 <sup>D</sup>   |
|                                            | ±29.5                 | ±95.0               | ±4.51               | ±4.52               | ±52.6               | ±35.1                | ±6.79               | ±8.13               |
| Batch-2EG                                  | 177 <sup>C</sup>      | 49.0 <sup>B</sup>   | 24.4 <sup>C</sup>   | 4.71 <sup>C</sup>   | 132 <sup>C</sup>    | 17.2 <sup>C</sup>    | 23.2 <sup>C</sup>   | 4.38 <sup>C</sup>   |
|                                            | ±85.0                 | ±76.4               | ±7.51               | ±5.23               | ±43.0               | ±19.6                | ±8.67               | ±2.03               |


a) Values are the mean (±SD) for  $n = 5$  animals. bcd Control and Batch-1 series groups not sharing superscript are significantly different,  $p < 0.05$ . BCD Control and Batch-2 series groups not sharing superscript are significantly different,  $p < 0.05$ .




**Figure 3.** Total concentrations of sphinganine (Sa) plus sphinganine 1-phosphate (Sa 1-P) in the kidneys of rats fed diets containing (50% w/w) uncooked (uncooked) or extruded (extruded) uncontaminated maize grits, or two batches of uncooked (Batch-1, Batch-2) FB<sub>1</sub>-contaminated grits, contaminated grits after extrusion (Batch-1E, Batch-2E) or contaminated grits after extrusion with glucose supplementation (Batch-1EG, Batch-2EG) for 3 or 8 wk. Values indicate the group mean score,  $n = 5$ ; error bars denote standard deviation. See text for explanation of scores. Control and Batch-1 series groups not sharing lower-case superscripts are significantly different and Control and Batch-2 series groups not sharing upper-case superscripts are significantly different,  $p < 0.05$ .

the tissues of rats exposed to these mycotoxins via the diet [25]. The bioavailability and toxic potential of matrix bound FB<sub>1</sub> or unknown FB<sub>1</sub> reaction or breakdown products in the cooked grits are unknown. However, because the *in vivo* effects were tightly correlated with the diets' measurable (not bound to matrix) FB<sub>1</sub> concentrations, it is unlikely that unknown fumonisin reaction products or matrix bound forms remaining in the extruded products made a significant contribution to toxicity. This issue requires further study.

FB<sub>1</sub> concentrations of the diets made with uncooked Batch-1 or Batch-2 grits were not high enough to cause liver lesions, influence clinical chemical variables or elevate liver sphingoid base concentrations. The Batch-1 and Batch-2 diets were however clearly nephrotoxic and effects on rela-



**Figure 4.** Group mean total kidney sphinganine concentrations (Sa ± Sa 1-P; from Fig. 3) plotted against group mean daily FB<sub>1</sub> intakes as determined from weekly body weight, food consumption and diet FB<sub>1</sub> concentration data.



**Figure 5.** Group mean histopathology scores for kidney lesions consistent with fumonisin exposure (from Fig. 1) plotted against group mean total kidney sphinganine concentrations (Sa ± Sa 1-P; from Fig. 3).  $R$  = linear correlation coefficient.

tive organ weight, histopathology and sphingoid base concentrations were consistent with well-defined apoptotic and biochemical outcomes following exposure to FB<sub>1</sub> [2, 4–7] or FB<sub>1</sub>-contaminated culture materials [16, 25, 26]. That kidney weights and histopathology scores of the Batch-1EG and control groups did not differ constitutes clear evidence that extrusion with glucose prevented toxicity of the less contaminated Batch-1 grits. Extruding Batch-1 grits without glucose was only partially protective, as was extrusion or extrusion with glucose supplementation of the more highly contaminated Batch-2 grits.

Sa increases in target tissues as a consequence of ceramide synthase inhibition by FB<sub>1</sub>, the mechanistic trigger causing sphingolipid metabolism disruption [25, 27]. The increases are dose-dependent and occur at lower doses or

earlier (time-course) than the appearance of microscopic lesions [2, 4, 26]. Consequently, elevated sphingoid base concentrations are a more sensitive indicator of low-dose FB<sub>1</sub> exposure than tissue pathology [24–26]. Sphingolipid metabolism in the kidneys of rats fed the uncooked Batch-1 and Batch-2 grits was clearly disrupted. Extrusion and extrusion with glucose reduced the sphingoid base elevations in the Batch-1 series of groups. Concentration decreased in the order: Batch-1 > Batch-1E > Batch-1EG = controls. Therefore, extrusion with glucose reduced the FB<sub>1</sub> concentration of Batch-1 grits to levels that, when mixed 1:1 with the basal diet, were not only insufficient to induce nephropathy but also too low to disrupt sphingolipid metabolism. Kidney sphingoid base concentrations in the Batch-2 series of groups exhibited a similar pattern of reduction by extrusion and greater reduction by extrusion with glucose. However, sphingoid base concentrations of the Batch-2 series of groups were higher than their Batch-1 counterparts and, with the exception of Sa 1-P of the Batch-2E group at wk 3, were consistently elevated ( $p < 0.05$ ) compared with the control groups.

Altogether, histopathology and sphingoid base findings established that extrusion with glucose supplementation using a twin-screw apparatus was an effective processing method for reducing FB<sub>1</sub> in contaminated grits and consequently for lowering daily intakes and reducing toxicity. The consequences of chronic exposure to maize-based foods processed by extrusion with glucose is unknown but should be investigated and consider the potential consequences arising from the high sugar content of the cooked product. In this regard, clinical laboratory results and microscopic examination of liver and other tissues in this study did not reveal evidence of any “non-fumonisin” effects.

The findings have also provided additional information on the relationship between sphingolipid metabolism disruption and toxicity. Both microscopic pathology scores and total Sa were tightly correlated with daily FB<sub>1</sub> intakes. Therefore, the lower scores and lesser sphingoid base elevations in the animals at 8 wk were likely a consequence of the lower doses (i.e. lower daily intakes) incurred after week 3 because of the animals’ growth, rather than a consequence of physiological adaptation. Second, while the severity of kidney pathology was correlated with total Sa concentrations at both time periods, it is interesting that the scores indicative of the most extensive tissue injury were associated with lower total Sa concentrations at wk 8 than at wk 3. Determining the reason for this was beyond the scope of this experiment but should be considered in the future. Progressive depletion of complex sphingolipids [8, 24, 27], compromised vitamin or nutrient utilization [8, 28], further disruptions of overall lipid metabolism and signaling, compositional changes in cell membranes [29], induction of nitric oxide [30, 31], or up- or down-regulation of cytokines [29, 32, 33] are among the factors that could be involved. It is noteworthy that deoxy Sa was not found in the livers or kidneys. Deoxy Sa, which is produced when alanine

is substituted for serine during de novo sphingolipid biosynthesis [20], accumulated in the livers of mice following exposure to FB<sub>1</sub> in other studies [20, 24]. Its absence in rat liver suggests that sphingolipid metabolism in rats and mice differ in some respects and that these differences might influence their target organ-specific responses to fumonisin exposure.

In conclusion, extrusion with 10% w/w glucose supplementation using a twin-screw configuration reduced FB<sub>1</sub> concentrations in maize grits more effectively than extrusion alone. Extrusion with glucose of grits containing ca. 10 ppm FB<sub>1</sub> significantly reduced daily FB<sub>1</sub> intakes and prevented the development of kidney lesions and disruption of sphingolipid metabolism when the processed grits were fed, 50% w/w in the formulated diet, to male rats. No evidence suggesting the formation of unknown toxic FB<sub>1</sub> reaction products was obtained. Therefore, extrusion with glucose supplementation is a potentially useful process to reduce the toxicity of fumonisin-contaminated maize.

*This study was supported in part by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service, grant #2005-35201-16329 The expert technical assistance of M. Vongkunthong, N. Stewart, J. Showker and E. Wray are gratefully acknowledged. We thank S. Bush and colleagues in the Pathology Department, University of Georgia College of Veterinary Medicine, Athens, GA for the clinical laboratory procedures. The pioneering work of S. Hendrich, P. Murphy and colleagues on the use of reducing sugars as a fumonisin detoxification strategy is acknowledged. References to their investigations are subsumed in the cited literature.*

*The authors have declared no conflict of interest.*

## 5 References

- [1] Marasas, W. F. O., Discovery and occurrence of the fumonisins: a historical perspective. *Environ. Health Perspect.* 2001, 109, 239–243.
- [2] Bolger, M., Coker, R. D., DiNovi, M., Gaylor, D. et al., Fumonisin, in: Safety Evaluation of Certain Mycotoxins in Foods, FAO Food and Nutrition Paper 74, World Health Organization, Geneva, 2001, pp. 103–279.
- [3] Scott, P. M., Recent research on fumonisins: a review. *Food Addit. Contam.* 2011, in press, DOI: 10.1080/1944099.2010.546000.
- [4] Voss, K. A., Riley, R. T., Norred, W. P., Bacon, C. W. et al., An overview of rodent toxicities: liver and kidney effects of *Fusarium moniliforme* and fumonisins. *Environ. Health Perspect.* 2001, 109, 259–266.
- [5] Voss, K. A., Smith, G. W., Haschek, W. M., Fumonisin: toxicokinetics, mechanism of action and toxicity. *Anim. Feed Sci. Technol.* 2007, 137, 299–325.

- [6] Gelderblom, W. C. A., Kriek, N. P. J., Marasas, W. F. O., Thiel, P. G., Toxicity and carcinogenicity of the *Fusarium moniliforme* metabolite, fumonisin B<sub>1</sub> in rats. *Carcinogenesis* 1991, 12, 1247–1251.
- [7] Howard, P. C., Eppley, R. M., Stack, M. E., Warbritton, A. et al., Fumonisin B<sub>1</sub> carcinogenicity in a two-year feeding study using F344 rats and B6C3F<sub>1</sub> mice. *Environ. Health Perspect.* 2001, 109, 277–282.
- [8] Gelineau-van Waes, J. B., Starr, L., Maddox, J. R., Aleman, F. et al., Maternal fumonisin exposure and risk for neural tube defects: mechanisms in an *in vivo* mouse model. *Birth Defects Res. A Clin. Mol. Teratol.* 2005, 73, 487–497.
- [9] Missmer, S. A., Suarez, L., Felkner, M., Wang, E. et al., Exposure to fumonisins and the occurrence of neural tube defects along the Texas-Mexico border. *Environ. Health Perspect.* 2006, 114, 237–241.
- [10] Jackson, L. S., Hlywka, J. J., Senthil, K. R., Bullerman, L. B. et al., Effects of time, temperature and pH on the stability of fumonisin B<sub>1</sub> in an aqueous model system. *J. Agric. Food Chem.* 1996, 44, 906–912.
- [11] Humpf, H.-U., Voss, K. A., Effects of thermal food processing on the chemical structure and toxicity of fumonisin mycotoxins. *Mol. Nutr. Food Res.* 2004, 48, 255–269.
- [12] Castelo, M. M., Jackson, L. S., Hanna, M. A., Reynolds, B. H. et al., Loss of fumonisin B<sub>1</sub> in extruded and baked corn-based foods with sugars. *J. Food Sci.* 2001, 66, 416–421.
- [13] Bullerman, L. B., Bianchini, A., Hanna, M. A., Jackson, L. S. et al., Reduction of fumonisin B<sub>1</sub> in corn grits by single-screw extrusion. *J. Agric. Food Chem.* 2008, 56, 2400–2406.
- [14] Dall'Asta, C., Mangia, M., Berthiller, F., Molinelli, A. et al., Difficulties in fumonisin determination: the issue of hidden fumonisins. *Anal. Bioanal. Chem.* 2009, 395, 1335–1345.
- [15] Motta, E. L., Scott, P. M., Bioaccessibility of total bound fumonisin from corn flakes. *Mycotoxin Res.* 2009, 25, 229–232.
- [16] Voss, K. A., Bullerman, L. B., Bianchini, A., Hanna, M. A., Ryu, D., Reduced toxicity of fumonisin B<sub>1</sub> in corn grits by single screw extrusion. *J. Food Protect.* 2008, 71, 2036–2041.
- [17] Jackson, L. S., Jablonski, J., Bullerman, L. B., bianchini, A., et al., Reduction of fumonisin B<sub>1</sub> in corn grits by twin-screw extrusion. *J. Food Sci.* 2011, in press.
- [18] Poling, S. M., Plattner, R. D., Weisleder, D., *N*-(1-deoxy-D-fructos-1-yl) fumonisin B<sub>1</sub> (FB<sub>1</sub>), the initial reaction product of fumonisin B<sub>1</sub> and D-glucose. *J. Agric. Food Chem.* 2002, 50, 1318–1324.
- [19] Zitomer, N. C., Glenn, A. E., Bacon, C. W., Riley, R. T., A single extraction method for the analyses by liquid chromatography/tandem mass spectrometry of fumonisins and biomarkers of disrupted sphingolipid metabolism in tissues of maize seedlings. *Anal. Bioanal. Chem.* 2008, 391, 2257–2263.
- [20] Zitomer, N. C., Mitchell, T., Voss, K. A., Bondy, G. S. et al., Ceramide synthase inhibition by fumonisin B<sub>1</sub> causes accumulation of 1-deoxysphinganine: a novel category of bioactive 1-deoxysphingoid bases and 1-deoxydihydroceramides biosynthesized by mammalian cell lines and animals. *J. Biol. Chem.* 2009, 284, 4786–4795.
- [21] Howard, P. C., Couch, L. H., Patton, R. E., Eppley, R. M. et al., Comparison of the toxicity of several fumonisin derivatives in a 28-day feeding study with female B6C3F<sub>1</sub> mice. *Toxicol. Appl. Pharmacol.* 2002, 185, 153–165.
- [22] Collins, T. F., Sprando, R. L., Black, T. N., Shackelford, M. E. et al., Effects of fumonisin B<sub>1</sub> in pregnant rats. Part 2. *Food Chem. Toxicol.* 1996, 36, 673–685.
- [23] Collins, T. F., Sprando, R. L., Black, T. N., Olejnik, N. et al., Effect of aminopentol on in utero development in rats. *Food Chem. Toxicol.* 2006, 44, 164–169.
- [24] Voss, K. A., Riley, R. T., Snook, M. E., Gelineau-van Waes, J. B., Reproductive and sphingolipid metabolic effects of fumonisin B<sub>1</sub> and its alkaline hydrolysis product in LM/Bc mice: hydrolyzed fumonisin B<sub>1</sub> did not cause neural tube defects. *Toxicol. Sci.* 2009, 112, 459–467.
- [25] Riley, R. T., Voss, K. A., Differential sensitivity of rat kidney and liver to fumonisin toxicity: organ specific differences in toxin accumulation and sphingoid base metabolism. *Toxicol. Sci.* 2006, 92, 335–345.
- [26] Burns, T. D., Snook, M. E., Riley, R. T., Voss, K. A., Fumonisin concentrations and in vivo toxicity of nixtamalized *Fusarium verticillioides* culture material: evidence for fumonisin–matrix interactions. *Food Chem. Toxicol.* 2008, 46, 2841–2848.
- [27] Merrill, A. H., Jr., Sullards, M. C., Wang, E., Voss, K. A. et al., Sphingolipid metabolism roles in signal transduction and disruption by fumonisins. *Environ. Health Perspect.* 2001, 109, 277–282.
- [28] Stevens, V. L., Tang, J., Fumonisin B<sub>1</sub>-induced sphingolipid depletion inhibits vitamin uptake via the glycosylphosphatidylinositol-anchored folate receptor. *J. Biol. Chem.* 1997, 272, 18020–18025.
- [29] Gelderblom, W. C. A., Abel, S., Smuts, C. M., Marnewick, J. et al., Fumonisin-induced hepatocarcinogenesis: mechanisms related to cancer initiation and promotion. *Environ. Health Perspect.* 2001, 109, 291–300.
- [30] Rotter, B. A., Oh, Y. N., Mycotoxin fumonisin B<sub>1</sub> stimulates nitric oxide production in a murine macrophage cell line. *Nat. Toxins* 1996, 4, 291–294.
- [31] Dresden-Osborne, C., Noblet, G. P., Fumonisin B<sub>1</sub> affects viability and alters nitric oxide production of a murine macrophage cell line. *Int. Immunopharmacol.* 2002, 8, 1087–1093.
- [32] Bondy, G. S., Barker, M. G., Lombaert, G. A., Armstrong, C. L. et al., A comparison of clinical, histopathological and cell-cycle markers in rats receiving the fungal toxins fumonisin B<sub>1</sub> or fumonisin B<sub>2</sub> by intraperitoneal injection. *Food Chem. Toxicol.* 2000, 10, 873–886.
- [33] Johnson, V. J., He, Q., Kim, S. H., Kanti, A. et al., Increased susceptibility of renal epithelial cells to TNF-alpha-induced apoptosis following treatment with fumonisin B<sub>1</sub>. *Chem. Biol. Interact.* 2003, 145, 297–309.

## Erratum

**Extrusion cooking with glucose supplementation of fumonisin-contaminated corn grits protects against nephrotoxicity and disrupted sphingolipid metabolism in rats.**

By Kenneth A.Voss et al., *Mol. Nutr. Food Res.* 2011, vol. 55, issue S2, pp. S312–S320

DOI: 10.1002/mnfr.201100067

In Section 3.6 of the Results, the superscript in the top panel of Fig. 1 (group mean histopathology scores for severity of lesions consistent with fumonisin exposure at week 8) for the group designated “Uncooked” should read “a A” not “ab A”.

In Section 3.6 of the Results, the number of animals per group in the Batch-2 series having nephropathy are given for “week 3 and week 8” not “week 3 and week 6”.