

## Structural and Fractal Dimensions are Reliable Determinants of Grain Yield in Soybean

A.A. Jaradat<sup>1</sup>, D. Surek<sup>2</sup> and D.W. Archer<sup>1</sup>

<sup>1</sup> United States Department of Agriculture (USDA)-Agricultural Research Service (ARS), 803 Iowa Avenue, Morris, MN 56267, USA

<sup>2</sup> Soil and Water Research Institute, Ankara, Turkey

\* Corresponding author (abdullah.jaradat@ars.usda.gov).

### Abstract

*Soybean [Glycin max (L.) Merr.] plants grown under five management strategies differed significantly in their geometric structures, and were classified with 75 to 100% correct classification, based on differences in their fractal dimension ( $Do$ ), midday differential canopy temperature ( $dT$ ), and canopy light penetration [ $\log(I/I_0)$ ]. Single soybean plants grown under a conventional system using moldboard tillage developed complex geometric structures, with significantly larger  $Do$  (1.477) values and grain yield (11.2 g per plant) as compared to plants grown under an organic system with strip tillage ( $Do = 1.358$ , and grain yield = 2.32 g per plant). Across management strategies,  $Do$  of single plants was a function of stem perimeter, circularity, and volume, and plant dry weight; whereas grain yield  $m^{-2}$  was a function of  $Do$ , plant dry weight and volume, and stem circularity. Knowledge of how plants respond to single and multiple management strategies will help agronomists develop better predictive models and will help farmers refine management practices to optimize yield.*

### Introduction

Plant size and architecture are important factors in determining crop productivity [1]; however, researchers are faced with the problem of developing reliable models for plant geometric structure and its relationship to yield and productivity, especially for plants with complex structures such as soybean [2, 3]. One approach to solving this problem is to use fractal analysis to provide new avenues of understanding the functional implications of the branching patterns in

relation to optimum space exploration by plants [1]. The fractal dimension ( $Do$ ) is considered [2] an effective tool for quantifying plant structure, measuring the structural response to cultural practices and modeling plant canopies.

The reproductive period is most sensitive to altered source strength and crop growth rate since it is the time during which important yield components are formed [4]. Changes in fractal dimension of several crops (e.g., corn and soybean) were found to be highly significant over time [2] reflecting the level of complexity in skeletal structure of single plants as the growth stages advanced. Several methods were used to quantify the relationships between soybean growth and development using growth analyses; however, limited information exists on the response of soybean's fractal dimension to management strategies. The objectives of this 2-yr study were to quantify the impact of management strategies on soybean's structural dimensions and  $Do$  during the reproductive growth stage, and to identify predictors of grain yield ( $gm^{-2}$ ).

### Materials and methods

Digital imagery [5] and analysis procedures [2, 6] were used to capture, measure, and statistically analyze several morphological traits of individual soybean plants grown under five management strategies, i.e., combinations of conventional (C) or organic (O) cropping system, conventional (C) or strip (S) tillage, recommended fertilizer rate (Y) and 2- or 4-yr crop rotation (Fig.1); for example, CCY4 is the management strategy with conventional cropping system (C), conventional tillage (C), with N fertilizer based on soil analysis (Y) and 4-yr crop rotation. Light interception by plant canopy [ $\log(I/I_0)$ ] and midday differential canopy temperature ( $dT$ ) were estimated as

described by Jaradat et al. [7]. The fractal analysis procedure employed the box count concept as outlined by Foroutan-pour et al. [2], where the fractal dimension ( $Do$ ) is constrained to be in the range of  $1.0 \leq Do \leq 2.0$ . A value of 1.0 indicates that the image is completely differentiable and that of 2.0 indicates that the image is irregular. Yield components (listed in Fig. 1) were measured on the same plants subjected to fractal dimension analysis and were used in subsequent statistical analyses.

The principal components (PC) option in the Nonlinear Iterative Partial Least Squares module and canonical discriminant (CD) analyses were used to analyze the standardized structural dimensions of individual soybean plants. Linear combinations of the original variables (i.e., leaf and stem structural dimension, in addition to  $\text{Log}(I/I_0)$ ,  $Do \cdot LAI$  [2], and  $dT$ ) that account for as much of total variation in the data set as possible were constructed. Canonical discriminant analysis was used to find the dimensions along which plants grown under different management strategies differ and to find classification functions to predict group membership on the first (R1) and the second (R2) canonical discriminant functions. The CD module was used to assess the variation among plants produced under two or more management strategies relative to the average variation found within all plants regardless of management strategies [8].

The impact of plant structural dimensions on  $Do$  and grain yield  $\text{gm}^{-2}$  was studied using the regression option in artificial neural networks (ANNs), then the models were subjected to sensitivity analysis to evaluate the relative importance of each variable in explaining  $Do$  or grain yield  $\text{gm}^{-2}$ . In this analysis, each predictor was treated in turn as if it were not available in the ANN model and the average value of that predictor was used. A sensitivity ratio was calculated by dividing the total ANN error when the predictor was treated as “not available” by the total ANN error when the actual value of the predictor was used. If the ratio is  $>1.0$ , then the predictor made an important contribution to  $Do$  or grain yield  $\text{gm}^{-2}$ . The higher the ratio, the more important the predictor is [8]. Additionally, we calculated the correlation coefficient ( $r$ ) and a ratio between the standard deviation (SD-ratio) of original and model data; higher  $r$  values and lower SD-ratio values are indicators of better model performance [8]. Sensitivity analysis was performed by generating response curves for each predictor to study its

relationship with  $Do$  and with seed yield ( $\text{gm}^{-2}$ ), while all other predictors were set at their mean value.

## Results

### Principal components regression

Slightly more than 50% of total variation in the whole data set was explained by the first two principal components (PCs; Fig. 1). Distinct separation between plants grown under organic and conventional systems was achieved on the basis of single plant characteristics, most of which were positively associated with conventional cropping system, conventional tillage and fertilizer application. Thousand-seed weight was the only variable associated with organic cropping system, strip tillage and no fertilizer treatment. Leaf area loaded on the third PC and accounted for additional 10% of total variance (data not presented). Grain yield  $\text{m}^{-2}$  was positively and closely associated with the fractal dimension, pods  $\text{m}^{-2}$ , and stem circularity, and to a lesser extent with the remaining plant structural dimensions on PC1 which explained 33% of total variation.




Fig. 1. Joint plot of six components of management strategies, ten soybean characteristics and the fractal dimension on the first two principal components in PC analysis.

A cumulative variance of 43% was explained by all independent variables in the PC regression (Fig. 2). Larger  $Do$  values, when multiplied by leaf area index ( $LAI$ ) [2], were positively associated with conventional cropping system and conventional tillage; whereas large values of  $dT$  and  $\text{log}(I/I_0)$  (i.e., less light interception by plant canopy) were associated with organic cropping

system (Fig. 2). The large  $LAI*Do$  values were loaded positively on PC1 along with most stem and leaf characteristics; however, stem width, leaf width, leaf perimeter, and stem perimeter were more closely associated with  $LAI*Do$  than the remaining stem and leaf structural dimensions.




Fig. 2. Mean separation and variance in grain yield per plant accounted for by differences among management strategies and soybean characteristics as predicted by the first two components in principal components regression. Means followed by the same letter do not differ significantly at the 5% level of probability.

A total of 51% of total variation in the dependent variables, accounted for by the first two PCs, explained 87% of total variation in grain yield per plant, which ranged from 2.32 (OSY4) to 11.2 g (CCY4) with significant differences among all management strategies. There were significant differences in grain yield per plant due to the tillage component, whether associated with conventional or organic systems, and due to crop rotation (2- vs. 4-yr) whether associated with conventional or strip tillage. Plants grown under CCY4 produced the largest grain yield (11.2 g), followed by CSY4 (9.82 g); whereas those grown under OCY4 and OSY4 produced the least (5.37 and 2.32 g per plant, respectively).

### Discriminant analysis

Discrimination among plant samples grown under five management strategies (Fig. 3) was clearly achieved using plant structural dimensions and three derived statistics (i.e.,  $dT$ ,  $Do$  and  $\log(I/Io)$ ). Two canonical discriminant roots accounted for a total of 92% of total variation and discriminated among plant samples with 75.0 (CSY4) to 100.0% (CCY2 and OSY4) correct

classification. The first canonical root (R1) was dominated by leaf circularity (i.e., ratio of minor to major axes), leaf area,  $\log(I/Io)$ ,  $Do$  and  $dT$ , accounted for the majority of variation (84%) and totally separated samples grown under organic system (i.e., OCY4 and OSY4, with 95.5 and 100.0% correct classification, respectively) from those grown under conventional system (CCY2, CCY4 and CSY4, with 100.0, 83.3, and 75.0% correct classification, respectively).




Fig.3. Canonical discriminant analyses and percent correct classification of soybean plants grown under five management strategies based on single plant structural dimensions,  $\log(I/Io)$ ,  $Do$  and  $dT$ .

Separation between the latter three groups along the second canonical root (R2), with 8% of total variation, ranged from 75 (CSY4) to 83.3% (CCY4). CAN2 was dominated by stem-related variables and there was clear overlap between plants grown under CCY4 and CSY4, on one hand, and those grown under CCY2. The three derived statistics (i.e.,  $Do$ ,  $dT$  and  $\log(I/Io)$ ) were closely associated with leaf circularity and leaf area, whereas stem structural dimensions were independent.

### Prediction of $Do$

Calibration and validation regression models were developed to predict  $Do$  as a function of  $dT$  are presented in Table 1. Correlation coefficients ( $r$  values) between measured and predicted grain yield using  $Do$  as a predictor were non-significant during the first two growth stages (data not presented); however,  $r$  values increased steadily from 0.74 (RGS3) to 0.96 as the plants approached maturity (RGS6); the respective  $r$ -values for the validation models were smaller (0.65 to 0.94)

albeit significant ( $p < 0.05$ ); however, the validation models performed very poorly during the first two reproductive growth stages. The intercept and slope the regression models increased steadily as plants approached maturity, the intercept approaching zero and the slope approaching unity.

Table 1. Calibration (C) and validation (V) partial least squares (PLS) regression models predicting soybean plant fractal dimension ( $Do$ ) as a function of midday differential canopy temperature ( $dT$ ) at four reproductive growth stages (RGS3 – RGS6, a and b are intercept and slope of regression models, respectively; \*,  $p < 0.05$ ).

Growth stage		PLS regression model			
		C	r	V	r
RGS3	a	0.68	0.74*	0.78	0.65*
	b	0.52		0.46	
RGS4	a	0.53	0.79*	0.58	0.74*
	b	0.63		0.60	
RGS5	a	0.29	0.89*	0.34	0.86*
	b	0.79		0.76	
RGS6	a	0.11	0.96*	0.14	0.94*
	b	0.92		0.89	

### Neural network and sensitivity analyses

A Multi-Layer Perception Neural Network (MLPR-NN) with 9:9-13-7-1:1 layers was the best neural network to predict soybean fractal dimension ( $Do$ ), and a General Regression Neural Network (GR-NN) with 13:13-43-2-1:1 neurons was the best to predict soybean grain yield  $\text{gm}^{-2}$  as a function of three plant traits and the fractal dimension ( $Do$ ). The MLPR neural network identified four independent variables with significant contribution in predicting both  $Do$  and grain yield  $\text{m}^2$  (Table 2). Plant dry weight was an important variable in predicting  $Do$  and grain yield  $\text{m}^2$ . A much simpler multi-layer perception neural network, with 13 hidden layers, was capable of predicting  $Do$  as compared to the more complex general regression neural network, with 43 hidden layers, needed to predict grain yield  $\text{gm}^{-2}$ . However, almost correlation coefficients were found between measured and modeled data for  $Do$  (0.87) and grain yield  $\text{gm}^{-2}$  (0.89) (Table 2).

The relationship between four predictors (Table 2) and  $Do$  was quantified and a regression equation was developed to predict  $Do$  as a function of each predictor while holding each of the remaining predictors at its mean value. Plant dry weight, stem volume, stem circularity and stem perimeter (Fig.

4) displayed different, albeit large and significant, effects on  $Do$ . The quadratic effect of stem volume was not significant. A plant dry weight of 6-7 g is capable of producing a maximum  $Do$  of 1.45-1.46; however,  $Do$  did not respond positively to any further increases in the plant dry weight beyond this level.

Table 2. Statistics of the Multi-Layer Perception Neural Network (MLPR-NN) with 9:9-13-7-1:1 layers predicting soybean fractal dimension ( $Do$ ), and of the General Regression Neural Network (GR-NN) with 13:13-43-2-1:1 neurons predicting soybean grain yield  $\text{gm}^{-2}$  as a function of three plant traits and the fractal dimension ( $Do$ ).

Variable	Ratio and (rank)		Test statistics	
	$Do$	Grain yield, $\text{gm}^{-2}$	$Do$	Grain yield, $\text{gm}^{-2}$
Plant dry weight, g	1.57(1)	1.6(2)		
Plants volume	1.49 (2)	1.35 (3)		
Plant circularity	1.35 (3)	1.35 (4)		
Plant perimeter	1.14 (4)			
$Do$		1.73 (1)		
Mean			1.43	172
S.D.			0.06	56.0
S.D.			0.65	0.63
Ratio				
r-value			0.87	0.89

Plant dry weight was the most important variable in predicting  $Do$ , followed by plant volume, plant circularity and plant perimeter; whereas  $Do$  was the most important variable, followed by plant dry weight, plant volume and plant circularity, in predicting grain yield  $\text{gm}^{-2}$ . The SD-ratios for  $Do$  (0.646) and for grain yield  $\text{m}^{-2}$  (0.632) were relatively similar.




Fig. 4. Sensitivity analyses of plant dry weight, stem volume, stem circularity and stem perimeter as predictors of fractal dimension ( $Do$ ) of soybean plants.

On the other hand,  $Do$  responded linearly to plant volume and, in a piecewise fashion, to stem circularity (i.e., ratio of minor to major axes, with a

breakpoint at  $Do=1.424331$ ) and stem perimeter (with a breakpoint at  $Do=1.4164$ ). Similarly, a nonlinear regression equation was developed to predict grain yield ( $\text{gm}^{-2}$ ) as a function of each predictor (i.e.,  $Do$ , plant dry weight, plant volume, and stem circularity, Table 2) while holding each of the remaining predictors at its mean value (Fig. 5). Positive and significant relationships were found among grain yield and each predictor, and the nonlinear portion of the regression equations was significant except for  $Do$ .

## Discussion

Short growing seasons in the upper Midwestern USA present serious time limitations on crop growth, in which soybean crop needs to establish and maximize canopy coverage rapidly to exploit available light [3]. Crop plants have been shown to adjust their architectural traits (Table 2) in response to management practices [2] and plant architecture, as characterized by  $Do$ , has been shown to impact grain yield in many crops [1].

Different management practices created a range of microenvironments in which soybean plants developed different architectures, as reflected by their  $Do$ ,  $dT$  and  $\log(I/I_0)$  values and on the large percentage of correct classification (75.0-100.0%). Further evidence on how grain yield responded to adjustments in plant architecture, which in turn responded to components of different management practices, is quantified in Fig. 1. The largest grain yield per plant (11.2 g) was positively associated with  $Do$ , conventional system, and conventional tillage, and was a result of maximum plant growth and development under the favorable conditions created by the CCY4 management strategy (Fig. 3).

The PLS regression models, especially during RGS3 to RGS6 (Table 1), succeeded in predicting  $Do$  as a function of midday differential canopy temperature ( $dT$ ), the value of which depends on air temperature, but will differ from it due to canopy characteristics, thermal characteristics and thermal conditions near the soil surface [10]. Reliability of the predictive equations (expressed as r-values) increased as the plants grew and changed the microenvironment within the canopy, and with time.






Fig. 5. Sensitivity analyses of fractal dimension ( $Do$ ), plant dry weight, plant volume, and stem circularity as predictors of grain yield  $\text{gm}^{-2}$  of soybean.

The response curves generated by the ANN models provided valuable information about the relationships among grain yield  $\text{m}^{-2}$  and a set of predictors beyond the information provided by simple correlation and regression models [9]. We

identified important  $Do$  and seed yield predictors using ANN models in an attempt to develop timely management practices that may help create optimum plant geometric structures (expressed as  $Do$ ) capable of maximizing light interception and midday differential canopy temperature, and thus producing the largest grain yield.

## References

- [1] G.P. Cheplick. Size and architectural traits as ontogenetic determinants of fitness in a phenotypically plastic annual weed (*Amaranthus albus*). 2002. *Plant Species Biol.* 17, 71-84.
- [2] K. Foroutan-Pour, P. Dutilleul, and D.L. Smith. Effect of plant population density and intercropping with soybean on the fractal dimension of corn plant skeletal images. 2000. *J. Agron. Crop Sci.* 184, 89-100.
- [3] D.B. Egli. Flowering, pod set and reproductive success in Soya bean. *J. Agron.* 2005. *Crop Sci.* 191, 283-291.
- [4] P. Pedersen, and J.G. Lauer. Soybean growth and development in various management systems and planting dates. 2004. *Crop Sci.* 44, 508-515.
- [5] F.J. Adamsen, T.A. Coffelt, J.M. Nelson, E.M. Barnes, and R.C. Rice, R.C. Method for using images from color digital camera to estimate flower number. 2000. *Crop Sci.* 40, 704-709.
- [6] W. Rasband. ImageJ. 1.33a National Institute of Health, USA. <http://rsb.info.nih.gov/ij/> java 1.5.0. 2004.
- [7] A.A. Jaradat, D.W. Archer, J. F-M. Johnson, S.J. Vankampen, S.W. Wagner, and J.J. Eklund. Sampling strategies for crop yield assessment within and among crop rotations. Proceedings of the Seventh International Conference on Precision Agriculture, July 25-28, 2004, Minneapolis, Minnesota. 2005 CDROM. 2005.
- [8] StatSoft Inc. STATISTICA (data analysis software systems) Version 7.1. [www.statsoft.com](http://www.statsoft.com). 2005.
- [9] StatSoft Inc., 2005b. Electronic statistics textbook (Tulsa, OK, USA). <http://www.statsoft.com/textbook/stathome.html>. (last accessed 3 July, 2006).
- [10] C.J. Birch, B. Andrieu, C. Fournier, J. Vos and P. Room. Modelling kinetics of plant canopy architecture – concepts and applications. 2003 *Eur. J. Agron.* 19, 519-533.