

**The Safety of AQUAFLO[®] (50% Florfenicol; Type A Medicated Article) Administered in Feed to Sunshine Bass (Female White Bass *Morone chrysops* × Male Striped Bass *M. saxatilis*)**

Jim Bowker*, David L. Straus¹, Molly Bowman, Cindy Ledbetter¹, Dan Carty, Miranda Dotson, Andrew J. Mitchell¹, and Bradley D. Farmer¹

*U.S. Fish and Wildlife Service, Aquatic Animal Drug Approval Partnership Program
4050 Bridger Canyon Road, Bozeman, Montana 59715, USA*

¹*Harry K. Dupree – Stuttgart National Aquaculture Research Center
U.S. Dept. of Agriculture, Agricultural Research Service
2955 Hwy 130 East, Stuttgart, AR 72160, USA*

Bacterial diseases are a major problem in aquaculture and account for significant losses (Clarke and Scott 1989; Frerichs and Roberts 1989; Bjorndal 1990). Therefore, the use of antimicrobial therapy plays an important role in aquaculture (Klontz 1987; Alderman 1988). Florfenicol (FFC) is a potent, broad-spectrum, antibacterial agent with bacteriostatic and bactericidal properties that is active against a variety of Gram-positive and Gram-negative bacteria (Horsberg et al. 1996).

AQUAFLO[®] (Intervet/Schering-Plough Animal Health Corp., Roseland, New Jersey USA) is a feed premix containing 50% florfenicol. Worldwide, AQUAFLO[®] has been approved in more than 20 countries including Norway, Japan, Chile, Canada, and the U.S., to control mortality in a variety of cultured fishes due to diseases associated with infectious bacterial pathogens. In the U.S., AQUAFLO[®] has been approved by the U.S. Food and Drug Administration (FDA) for use to control mortality in (1) catfish due to enteric septicemia associated with *Edwardsiella ictaluri*, (2) freshwater-reared salmonids due to coldwater disease associated with *Flavobacterium psychrophilum*, and (3) freshwater-reared salmonids due to furunculosis associated with *Aeromonas salmonicida* (Bowker et al., in press). For all currently labeled uses in the U.S., the FDA-approved treatment regimen is to administer AQUAFLO[®] in feed at 10 mg FFC per kg fish body weight (BW) per d for 10 consecutive days.

There is considerable interest within U.S. domestic aquaculture to expand the use of AQUAFLO[®] to include therapeutic treatment of freshwater-reared, cold-, cool-, and warmwater finfish at 15 mg FFC per kg fish BW per d for 10 consecutive days. To obtain such an approval, data must be generated to show that this higher dose is safe to representative target animals. Consequently, we conducted a target animal safety (TAS) study to evaluate the safety of AQUAFLO[®] administered in feed to sunshine bass (female white bass *Morone chrysops* × male striped bass *M. saxatilis*).

Methods

The study was conducted March 17 – April 8, 2009, at the Harry K. Dupree Stuttgart National Aquaculture Research Center (U.S. Department of Agriculture, Agricultural Research Services, Stuttgart, Arkansas USA) and consisted of a 3-d acclimation

period, 20-d exposure period, and 1-d postexposure period. Test fish (mean ± SD) total length and weight were 11.1 ± 0.42 cm and 13.6 ± 1.58 g, respectively. AQUAFLO[®] was administered in feed at 0× (0 mg per kg), 1× (15 mg per kg), 3× (45 mg per kg), or 5× (75 mg per kg) the proposed maximum therapeutic dose of 15 mg FFC per kg fish BW per d for 2× (20 d) the proposed therapeutic treatment duration of 10 d. Completely randomized design procedures were used to assign each of the four exposures to 3 of 12 test tanks, and to stock 20 fish into each test tank. The study was single-blinded such that personnel involved in day-to-day data collection did not know which exposures were assigned to which test tanks. Feed samples were collected and sent to Eurofins/AvTech Laboratories (Portage, Michigan USA) to analytically verify the homogeneity and stability of FFC in the 1×-, 3×-, and 5×-medicated feeds, and to determine if there was any FFC in the 0× (control) feed.

Mortality, general fish behavior, fish feeding (appetite) behavior, water temperature, and dissolved oxygen concentration were assessed daily. In addition, source water hardness, alkalinity, and pH were measured weekly. On the postexposure day, all test fish were collected from test tanks, euthanized, measured for total length and weight, and necropsied. During necropsy, 10 fish were randomly selected from each tank and tissues collected for histological examination. Of these, two fish per tank were randomly selected for evaluation of 11 tissues (gill, liver, anterior kidney, posterior kidney, brain, heart, muscle, skin, spleen, pyloric intestine, and rectal intestine). The other eight fish per tank were evaluated for four tissues (gill, liver, anterior kidney, and posterior kidney). All tissues were examined for lesions or cellular changes (hereafter referred to as lesions) that might provide evidence of AQUAFLO[®]-induced toxicity. Lesions were scored on a six-point ordinal severity scale. Fish in 0× and 5× exposure groups were examined and compared first. If lesions were detected in one or more tissues of the 5×-exposure group that were (a) marked or severe, (b) not observed in the 0× exposure group, and (c) appeared to be AQUAFLO[®]-induced, then all relevant tissue(s) were examined in all fish representing the 3× exposure group. If lesions were not detected in the 5× exposure groups, tissues from the 3× exposure groups were not evaluated.

*Corresponding author: jim_bowker@fws.gov

Statistical analysis of mortality data was not required for no mortality occurred during the study. Gross necropsy and histopathology findings were described qualitatively. Secondary response variables (mean feed consumption, mean body weight) were analyzed by one-way ANOVA. Treatment differences were considered significant if $P < 0.05$.

Results and Discussion

During the in-life phase of the study, no fish died or exhibited signs of morbidity. Moreover, general fish behavior was characterized as normal in all tanks. Fish fed actively, broke the surface of the water while feeding, and consumed all feed offered within 10 sec; therefore, feeding (appetite) behavior was characterized as “100% of feed consumed.”

At the end of the in-life phase, fish had grown an average of 1.4 cm and 7.3 g, and overall mean total length and weight of test fish were 12.6 ± 0.45 cm and 21.0 ± 2.29 g, respectively. Mean total length ($P = 0.493$) and mean weight ($P = 0.955$) did not differ among exposure groups.

No lesions of any kind were observed in the heart, liver, or rectal intestine tissues; no degeneration was observed in skin tissue; and no inflammation was observed in posterior kidney tissue or spleen tissue in fish examined from the 0× or 5× exposure groups. Lesions not considered to affect overall health of the fish (e.g., mild or moderate) were observed in all other tissues examined (Table 1). Only posterior kidney tissue was examined in fish from the 3× exposure group; however, results were nearly identical to those observed in the 0× and 5× exposure group fish. For example, mild to moderate tubule epithelial degeneration and necrosis was observed in 100% of the 30 fish examined from the 3× exposure group.

Based on histological data as reported above, mortality data, the fact that fish fed aggressively regardless of the FFC concentration in feed offered, and that general fish behavior was considered normal, we concluded that the margin of safety was at least five times greater than the proposed therapeutic treatment concentration of 15 mg FFC per kg BW per d. No fish health and/or histological lesions were detected that might indicate that

75 mg FFC per kg fish BW per d for 20 d was not safe to sunshine bass. In addition, reviewers from FDA’s Center for Veterinary Medicine found that the data demonstrated there is an adequate margin of safety for the use of AQUAFLO[®] at a dosage of 15 mg FFC per kg fish body weigh per d for 10 consecutive days.

Acknowledgments

We thank Intervet/Schering-Plough Animal Health Corporation for supplying AQUAFLO[®] and paying for feed analysis. Beth MacConnell (FWS, retired) did histological evaluations. Tom Bell and Dave Erdahl, FWS AADAP, critically reviewed this bulletin.

References

- Alderman, D. J. 1988. Fisheries chemotherapy: a review. **In:** *Recent Advances in Aquaculture, Vol. 3*. Croom Helm, London.
- Bjorndal, T. 1990. *The Economics of Salmon Aquaculture*. Blackwell Scientific Publications, Oxford.
- Bowker, J. D., V. E. Ostland, D. Carty, and M. P. Bowman. in press. Effectiveness of Aquaflor[®] (50% florfenicol) to control mortality associated with *Streptococcus iniae* in freshwater-reared, subadult sunshine bass. *Journal of Aquatic Animal Health*.
- Clarke, R. and D. Scott. 1989. An overview of world salmon production and recent technology developments. *AAC Bulletin* **89-4**: 31-38.
- Frerichs, G. N. and R. J. Roberts. 1989. The bacteriology of teleosts. **In:** *Fish Pathology, 2nd edition*. Balliere Tindall, London.
- Horsberg, T. E., K. A. Hoff, and R. Nordmo. 1996. Pharmacokinetics of florfenicol and its metabolite florfenicol amine in Atlantic salmon. *Journal of Aquatic Animal Health* **8**:292-301.
- Klontz, G. W. 1987. Control of systemic bacterial diseases in salmonids. *Salmonid*, **December**: 5-13 .

Table 1. Relative number of sunshine bass per exposure condition in which mild to moderate lesions were detected in the 0× and 5× (75 mg FFC per kg fish BW) exposure groups.

| Tissue | Sample size | Exposure (mg FFC per kg fish BW per d) | |
|---|-------------|--|------|
| | | 0 | 75 |
| Muscle - degeneration of fibers | 6 | 33% | 67% |
| Spleen - melanomacrophages | 6 | 100% | 83% |
| Pyloric intestine - epithelial degeneration | 6 | 33% | 17% |
| Pyloric intestine - epithelial necrosis | 6 | 33% | 17% |
| Gill - epithelial lifting | 30 | 97% | 100% |
| Gill - telangiectasias | 30 | 67% | 47% |
| Gill - inflammation | 30 | 93% | 83% |
| Anterior kidney - melanomacrophages | 30 | 13% | 3% |
| Posterior kidney - tubule epithelium degeneration | 30 | 100% | 100% |
| Posterior kidney - tubule epithelium necrosis | 30 | 97% | 100% |