

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



FARM SHEEP RAISING FOR BEGINNERS

F. R. MARSHALL and R. B. MILLIN
Of the Animal Husbandry Division



Has been rev.
--see rev.ed.
binders at
end of file.

FARMERS' BULLETIN 840

UNITED STATES DEPARTMENT OF AGRICULTURE

Contribution from the Bureau of Animal Industry
A. D. MELVIN, Chief

Washington, D. C.

July, 1917

Show this bulletin to a neighbor. Additional copies may be obtained free from the
Division of Publications, United States Department of Agriculture

LAMBS and wool are in strong demand and prospects are good for profit in raising sheep on the farm.

Several million acres of land in the United States which produce good summer feed for sheep are not grazed at present.

Sheep raising does not require expensive equipment or heavy labor, but does require study and continuous attention.

Early fall is the best time to start a flock. Good-grade ewes and a pure-bred ram are the best for beginners.

The beginner may acquire experience with less than 20 ewes, but for economy of time and fencing, and to assure proper care, flocks of 60 or more ewes are better.

In most cases lambs are most profitable if made ready for market at about 4 months, weighing 65 to 75 pounds.

Unless the flock has a very large territory to range over it is necessary to make divisions of the pasture or to use seeded forage crops. This permits the change of grazing ground necessary to insure the health and thrift of the lambs.

FARM SHEEP RAISING FOR BEGINNERS.

CONTENTS.

	Page.		Page.
The outlook for sheep raising	3	Management at breeding time	11
Sectional prospects for sheep production.....	4	The flock in winter.....	12
Requirements for sheep raising	5	The lambing season	14
Returns from sheep raising	7	The flock in summer.....	17
Starting the flock.....	8	Preparing lambs for market.....	22
Size of flock.....	10		

THE OUTLOOK FOR SHEEP RAISING.

AN IMPORTANT change in the extent and character of the American farm sheep industry began in 1915. Higher prices for lambs and wool in that year and the years following attracted wide attention to this branch of animal husbandry. Agricultural developments of former years, with their cheaper grains and lower values of meat animals, had not shown the economical advantages of the mutton and wool sheep as a quick source of income, produced mainly from pasture, forage crops, or roughages, and with a very low labor requirement.




FIG. 1.—Such pasturage as is shown in this illustration is excellent for sheep. The grazing of sheep on such land yields a good income and improves the pasturage for larger stock.

The higher prices of sheep products following 1914 were caused in part, but not mainly, by market conditions resulting from the war. The supply of lamb and mutton had been decreasing for some time in spite of the growing demand, particularly for lambs. Wool values were advancing before the outbreak of the war. Although the world consumption of wool was increasing, no foreign country, with the exception of South Africa, seemed able to increase its exports. Increased supplies of wool in the future must come chiefly from farm flocks. In the United States conditions for farm sheep raising are more favorable than in any country which has not already developed to the point at which sheep are necessary for intensive farming.

Prospective values for lambs and wool and the special economies incident to their production insure for farm sheep raising a large and permanent place, either on those farms where sheep raising is made a specialty or where flocks form a permanent part of a system of mixed farming.

SECTIONAL PROSPECTS FOR SHEEP PRODUCTION.

In the Eastern States the large and numerous flocks of earlier years were kept almost entirely for wool production. Subsequently the increasing wool supplies from other sections and from abroad, together with the demand for other agricultural products of higher value, brought about a decline in the number of farm sheep in these States. The market demand for mutton and lamb at that time was very limited, and when it became broader the cheaply produced western supplies were for some time equal to all requirements. To-day the western shipments have not only ceased to increase but have actually grown less as a result of the reduction of the range area.

In New England particularly, while many new flocks have recently been started, hesitancy has been due to a wrong interpretation of former statistics of farm sheep in that section. The decline that once occurred in New England flocks has but slight relation to present conditions and prospects. The sheep raising of the present is planned to market lambs at from 4 to 5 months of age, and wool, though important, is not the primary consideration. The full and economical utilization of New England farm labor, pastures, hay, and silage calls for more and larger flocks to supply the near-by markets. The present production can probably be multiplied three times without materially lessening other live-stock production.

Throughout the entire length of the Appalachian Mountain Range in Pennsylvania, Maryland, Virginia, West Virginia, Kentucky, Tennessee, and North Carolina there are large areas of land comparatively low in value and well suited for sheep raising.

In the hillier sections of northern Arkansas and southern Missouri, and in the cut-over timber regions of the Gulf States, there are also large areas of comparatively cheap lands which furnish favorable conditions for the keeping of large flocks of sheep at comparatively low cost. Similar opportunities are found in the cut-over timberlands in Michigan, Wisconsin, Washington, and Oregon.

On the higher priced lands of the corn belt a profitable system of sheep raising is being worked out along the lines followed on the intensively farmed areas in England and Scotland. While land values in this section are much higher on the acre basis than in the regions above referred to, there is comparatively little difference in the value of the amount of land required per head for sheep. While few farms in this section are likely to be devoted exclusively to commercial sheep raising, the different labor requirements for cattle and swine make it desirable to keep at least 1 ewe to 2 acres. This should add materially to the net income from the farm.

On western irrigated farms there seems likely to be developed an intensive sheep industry. The alfalfa and other forages produced on these lands come nearer to being satisfactory as a sole ration for sheep than for any other stock. The use of irrigated pastures or the rotation of forages will provide excellent summer feed and at the same time avoid the cost of labor for harvesting where there is not an opportunity of using the open range or forest reserve for grazing at that time of year.

REQUIREMENTS FOR SHEEP RAISING.

SOIL AND CLIMATE.

Sheep are naturally the inhabitants of high and dry areas. They thrive, however, on any except wet, swampy land. The fine-wool breeds especially prefer drier lands, while one or two of the British breeds are particularly adapted to lowlands. Sheep raising has been successfully carried on in areas of tropical temperature with low rainfall, but their rearing in high temperatures with a high rainfall has not been fully demonstrated.

PASTURE AND FEED.

Sheep naturally graze over rather wide areas and seek a variety of plants. This habit particularly adapts them to being kept in large numbers on lands of sparse vegetation or furnishing a variety of grasses or other plants. They do better on short and fine grasses than on coarse or high feed. They will eat a good deal of brush and, if confined to small areas, will do a fair job at cleaning up land. When used in this way, or on land producing only brush, they can not be expected to prove very satisfactory in the production of good lambs or good wool.

The cheapest and best feed for sheep is pasture such as described, or sown forage crops of cereals, rape, etc. Frequent changes of grazing ground are necessary to health and maximum thrift when pastures do not offer a wide range. This calls for fencing to subdivide permanent pastures, or for tight fencing around large runs in which they are to be kept. Movable fences may be largely used for carrying sheep on smaller areas of forage crops.

Grain feeding is seldom profitable when good grazing is to be had. Under some conditions flocks can be kept in good condition and lambs marketed without the use of any grain. One hundred pounds of grain in a year for one ewe and her lambs is the maximum that is likely to be used profitably under any conditions. The largest quantity may be used with ewes dropping lambs before pasture is ready and for the lambs at that time, but the feeding that is most economical and most likely to keep the flock in good condition is that which provides frequent changes of good pastures and grazing crops and winter rations mainly of good leguminous hays, with some succulent feeds, reserving what grain is to be used to feed in winter and after the lambs are born.

Silage or roots furnish cheap feed and are especially useful in keeping ewes in good condition during the winter. Too free use of roots for ewes in lamb is sometimes considered to increase the losses of young lambs, and the exclusive use of silage as a roughage has been shown to be unsafe, either for the ewes themselves or for the lambs to be dropped.¹

BUILDINGS AND FENCES.

In any part of the United States the main essentials of sheep barns are dryness and freedom from drafts. Unless lambs are to be dropped in cold weather, no expense to provide warmth is necessary, as the buildings should seldom be closed. Protection from winter rains and heavy snowfalls is desirable, but the best results may be expected when ewes are allowed access to a dry bed in the open.

Fences to hold sheep should be of woven wire, boards, or rails. Barbed or smooth wire can not be used satisfactorily, though a 36-inch woven-wire fence at the ground with two or three strands of wire is commonly used.

The construction, planning, and cost of a variety of barns and sheds for sheep and of dog-proof fences is discussed fully in Farmers' Bulletin 810, "Equipment for Farm Sheep Raising."

LABOR.

The amount of labor required to keep a farm flock in the condition necessary to insure maximum returns and lowest cost of production

¹ Pennsylvania Experiment Station Bulletin 144.

varies according to systems followed in different sections. In all cases the amount of labor is small in proportion to that required by other live-stock products of equal value. Feeding the sheep in winter is light labor, and the manure need not ordinarily be removed from pens oftener than once in six weeks during the time the flock is housed.

However, sheep raising should not be engaged in with an idea that little attention is required. The wants of sheep are numerous and varied, and frequent attention is required to forestall conditions that will result in ill health or lack of thrift. With a large flock at lambing time frequent attendance day and night is necessary to avoid losses of ewes and young lambs. While their habits are quite different from those of other farm animals, sheep are an interesting study. Sheep management can be learned and understood by anyone who is willing to observe carefully and think and attend to the details as attention is required.

RETURNS FROM SHEEP RAISING.

The gross annual returns from ewes of breeding age may be expected to range from \$8 to \$15 a head, depending upon the percentage of lambs raised, the weights of the fleeces, and the values for these products. The lamb and wool yields depend largely upon the breed selected. With ewes of any one of the medium-sized mutton breeds 115 per cent of lambs can be raised, and 150 per cent is not infrequently reached. Lambs are most in demand when fat at a weight of from 65 to 80 pounds. These weights and sufficient fatness can be obtained at from 4 to 5 months of age with very little grain feeding, and before the lambs eat much of the forage in pasturage, if the ewes' feed produces a continuous and plentiful supply of milk.

The wool returns vary from 7 to 11 pounds per ewe. The larger mutton breeds yield more, as do also the fine wools, but the value per pound of the latter is usually less on account of the greater proportion of natural grease or yolk present.

It is difficult to estimate satisfactorily the net returns from a flock of ewes. In comparison with cattle and swine, sheep can be made to yield practically the same net returns on the value of the land, if well cared for, and if kept on lands reasonably well adapted for sheep raising.

For farms of all-arable land the Illinois Experiment Station¹ has recommended a plan of live-stock production which includes 80 ewes (1 ewe to 2 acres) along with 22 breeding cows and 12 brood sows. Arable land of the best class when used exclusively for sheep can be made to support from 5 to 8 ewes (with their lambs until marketed)

¹ Circular entitled "Facts Regarding Mixed System of Farming."

per acre. On pastures suitable for either cattle or sheep five ewes may be considered the equivalent of one cow or steer, and the winter feed required for one breeding cow not in milk would be equivalent to that needed for about eight ewes.

STARTING THE FLOCK.

TIME TO START.

Late summer or early fall is the most favorable time to make a start in sheep raising. Ewes can be procured more readily at this time, and when purchased can be kept on meadows, grain stubble fields, or late-sown forage crops to get them in good condition for breeding. Experience with the ewes through fall and winter will also render a beginner more capable of attending to them at lambing time. It is seldom possible to buy any considerable number of bred ewes at reasonable prices.

SELECTION OF STOCK.

The inexperienced sheep raiser should begin with grade ewes of the best class available and a pure-bred ram. The raising of pure-bred stock and the selling of breeding rams can best be undertaken by persons experienced in sheep raising. The selection of the type and breed of sheep should be made by considering the class of pasture and feeds available and the general system of farming to be followed, along with the peculiarities of the breeds and the condi-




FIG. 2.—A good type of a pure-bred mutton sire.

tions and kind of feeding and management for which each has been especially developed.¹

It is highly advantageous for all, or a majority, of the farms in a neighborhood to keep the same breed of sheep, or at least to continue the use of rams of the same breed. After a decision has been made as to a suitable breed, the aim should be to obtain ewes that are individually good and that have as many crosses as possible of the breed selected. With such a foundation and the continuous use of good pure-bred rams of the same breed, the flock will make continuous improvement.

In looking for ewes of desired types and breeding it will often be found impossible

to get them near at home at a reasonable price. Ewes from the western ranges can be obtained directly from a stockyard market. For the most part the range ewes are of Merino breeding. First-cross ewe lambs and less often older stock bred on the range and sired by rams of the down or long-wool breeds are sometimes obtainable. These, or even the Merino ewes, furnish a foundation for the flock that can be quickly graded up by using rams of the breed preferred. The lambs from Merino ewes and mutton rams grow well and sell



FIG. 3.—A 2-year-old mouth. The two large teeth in the center are the first pair that come in at about 12 months of age. The next, or 2-year-old pair, show one on each side of the center pair. The two small teeth on the right are lamb teeth.

¹Farmers' Bulletin 576, entitled "Breeds of Sheep for the Farm," discusses the adaptability of each of the common breeds. It is obtainable from the Department of Agriculture upon request.

well if well cared for, but the yield is less than when ewes with some mutton blood are used. The sheep from the range are less often infested with internal parasite than are farm sheep, and in the large shipments there is opportunity for closer selection.

AGE OF EWES.

Yearling or 2-year-old ewes are preferable to older stock. Ewes with "broken mouths"—that is, those that have lost some of their teeth as a result of age—can be purchased cheaper than younger ones, but are not good property for inexperienced sheep raisers.

Until a sheep is 4 years old its age can usually be told within a few months. The lambs have small, narrow teeth, known as milk teeth. At about 12 months of age the two center incisors are replaced by two large, broad, permanent teeth. At about 24 months two more large teeth appear, one on each side of the other pair (see fig. 3). Another pair appears at 3 years of age, and the last, or corner teeth, come in at about the end of the fourth year, and the sheep then has a full mouth. Heavy or light feeding has considerable effect upon the exact time of appearance. After a sheep becomes 4 years old the exact age can only be estimated. As age advances, the adult teeth become shorter and the distance between them increases. The normal number of teeth may be retained until 8 or 9 years of age, but more often some are lost after the fifth year.

In buying ewes, particularly those from the range, it is desirable, when possible, to examine the udders to see that they are free from lumps that would prevent the ewes from being milkers. It is necessary to guard also against buying ewes that are useless as breeders because of the ends of the teats having been clipped off at shearing.

SIZE OF FLOCK.

Persons wholly inexperienced with sheep will do well to limit the size of the flock at the start. A beginner can acquire experience quite rapidly with 8 or 10 ewes. It is very doubtful, however, whether anyone should make a start with sheep unless the arrangement of the farm and the plan of its operation allow the keeping of as many as 30 ewes, and in most cases 60 or more will be handled better and more economically than a very small flock. The number of ewe lambs that can be kept for breeding each year should be about one-half the number of breeding ewes. Old ewes should be discarded when 5 years old. When this is done and the poorest of the ewe lambs are sold a flock will ordinarily double in size in three years. After two seasons' experience it will be a good plan to buy more ewes when good ones can be obtained at a fair price.

The economical disadvantage of a very small flock lies in the fact that the hours of labor are practically the same for a dozen or 20

ewes as for the larger flock. The fencing to allow desirable change of pastures or to give protection against dogs is about the same in either case, so that the overhead charges per ewe are much smaller in the case of the larger flock. Furthermore, the small flock on a farm having larger numbers of other animals is unlikely to receive the study and attention really needed or that would be given to one of the chief sources of the farm income.

MANAGEMENT AT BREEDING TIME.

THE EWES.

The period of gestation in sheep is 145 days. Ewes should be mated to drop their first lambs when about 24 months old. The first few cool nights in late summer or early autumn cause the ewes to come in heat, although some breeds come in heat at almost any time of the year. These periods in which the ewes will breed last from 1 to 3 days and recur at intervals of from 14 to 19 days. At the time the ewes are bred they should be gaining in weight. Feeding to produce this condition for breeding is commonly called "flushing." The main purpose of flushing the ewes is to secure a larger lamb crop and to have the lambs dropped as near the same time as possible, but it also brings the ewes into good condition for the winter. To accomplish this the ewes are changed from scant to abundant pastures of timothy, bluegrass, or rape. Rank watery fall growths of clover are of little use for this, as they often bring the ewe in heat several times and are not particularly fattening. Often some grain is fed as a supplement to the pastures. Corn is not especially good for this, oats being much better. Pumpkins strewn over the fields are excellent. At this time any large locks of wool or dung tags about the tail should be removed.

THE RAM AT BREEDING TIME.

Beginning about a month before the breeding season, the ram should be given some extra grain. Two parts of oats and one of bran by bulk form an excellent mixture. Oats alone are also very good. If the ram is thin the following mixture is excellent: Corn, 5 parts; oats, 10; bran, 3; and linseed-oil meal, 2 parts, by weight.

The number of ewes a ram will serve depends largely upon his age and the way he is handled. A ram lamb may serve from 5 to 15 ewes, depending upon his maturity. A yearling may serve from 15 to 25, while a mature ram well cared for should serve from 40 to 60 if allowed to run loose with the flock. By permitting him to be with the ewes an hour morning and evening more ewes can be bred. If the ram is old or injured or is to be bred very heavily, another ram may be used to locate the ewes in heat and thus save the older ram

from the necessary work of circulating through the flock. A bag or a piece of cloth tied under the belly prevents the "teaser" from serving the ewes. If the ram is allowed to run in the field with the ewes he may be made to mark the ones he has served, so that the approximate dates of lambing can be determined. A daub of special branding paint that later will scour out of the wool can be applied every day or two to the left side of his chest and brisket for the first two weeks, on the right side for the next two, and on the middle for the last two weeks of the season. Different colors of paint may also be used, but under no consideration should any mixture containing tar be used. When the ram is not in the flock he will be quieter and more easily handled if one or two ram or wether lambs or bred ewes are kept running with him.

FALL FEEDING.

Stubble and stalk fields may well form the principal means of sustenance for the breeding flock in the fall if they are used before the rains injure their feeding value. Fence strips in plowed fields may also give good grazing for a few days. Clover and grass pastures may well be left until the stubble and stalk fields have been used. For regions where the winters are open a heavy stand of well-cured bluegrass will help very much in carrying the flock through the winter in good condition. Green rye pastures in the late fall give considerable succulence and furnish exercise for the flock. In the South velvet beans will be found of great help in carrying the flock into January.

The shepherd should train himself to read the condition of his sheep by feeling the bone of the loin or back. At no time while they are in lamb should ewes be allowed to lose in weight. In open wet fall seasons there is danger of waiting too long to start feeding. A rank growth of soft grass may appear to be good feed, but the real need of the flock should be determined by a closer examination of the actual condition of all or a representative number of the ewes.

THE FLOCK IN WINTER.

WINTER FEED.

Winter management has a very important relation to the returns from the flock. The feeding should be such as will produce the most vigorous lambs and at the same time keep the wool in good condition. Leguminous hays, straws, and cornstalks usually form the main part of economical winter rations. Clover, alfalfa, or cowpea hay, if of good quality, may be used as the sole feed until near lambing time, from 3 to 3½ pounds being sufficient for ewes weighing less than 150 pounds. Oat and wheat straw are better than rye or barley straw. The beards of the latter are likely to prove trouble-

some. Cornstalks placed where the ewes can eat off the leaves may be used as a part of the roughage ration. If this ration is made up largely of cornstalks or straw, a nitrogenous concentrate should also be used. Timothy hay is not good sheep feed.

Such succulent feeds as roots or silage are desirable in keeping the ewes in good health. The use of silage will often materially reduce the cost of the ration, but silage can not safely be used without any hay. Only silage from well-matured corn should be used for sheep, and caution should be exercised to guard against feeding spoiled, frozen, or moldy silage. It is not advisable to feed more than 3 pounds per head daily of this feed.

For bred ewes, roots, particularly turnips, should be used sparingly until after lambing. Each of the following rations contains approximately the amount of the various nutrients required daily for ewes of from 120 to 145 pounds in weight:

(1)	(3)
2 pounds alfalfa or cowpea hay,	2½ pounds alfalfa,
2 pounds corn silage,	2 pounds silage.
½ pound shelled corn.	(4)
(2)	1 pound oat straw,
2 pounds alfalfa,	2 pounds corn silage,
2 pounds corn stover (amount eaten).	½ pound oil meal,
	½ pound corn.

Where the ewes can run on fall wheat or rye during the winter months the pasture should be supplemented by some dry or concentrated feed. Silage or roots are not desirable when the pasturage is soft or green. One-half pound of cottonseed meal contains the daily requirement of protein for pregnant ewes. When price suggests the use of this concentrate, the other feeds should be of a carbonaceous character. One-quarter pound of cottonseed meal per day and a selection of other feeds will be better than a ration containing a larger amount of cottonseed meal.

EXERCISE IN WINTER.

If the lambs are to be born strong and vigorous, a moderate amount of exercise is necessary for the ewes during the winter. This can be obtained by scattering their roughage over a field and allowing them to work back and forth over it while eating, or by feeding some of the roughage some distance away from their shelter. If winter pastures are used, no other arrangement for exercise is necessary. At no time should the pregnant ewes be forced to wade through deep mud or snow, neither should they be chased by dogs nor forced to jump over boards nor to pass through narrow doors, as such treatment is sure to cause loss of lambs or of both ewes and lambs.

If fleeces are allowed to become soaked with rain or wet snow, colds and pneumonia will surely result. Dry snow, on the other hand, has no ill effect, as the ewes readily shake it off.

THE LAMBING SEASON.**IMPORTANCE OF CARE DURING LAMBING.**

The lambing season is the shepherd's harvest time, and the size and quality of the crop practically determine the profits. A large crop of good lambs is the base of good financial returns, while a small crop of lambs means less profit, and if they are inferior in quality great skill and care are necessary to make any profit. At this time extra attention must be given to the ewes and lambs. In no other way can time be used to better advantage on the farm. If a record of the date of service has been kept, the approximate date of lambing can readily be foretold, for the ewes will generally carry their young about 145 days (five days less than five months).

CARE OF THE EWES.

Heavy grain feeding just before lambing is likely to cause udder troubles. At this time the wool around the udder should be clipped short to allow the lamb to find the teats readily. Just before lambing the ewe becomes restless and appears sunken in front of the hips. She should be put into a separate pen, which may be made of two light panels fastened together by a hinge and set in a corner.¹

These panels permit the ewe to see the other members of the flock and prevent her from becoming excited or nervous. Their use also prevents other sheep from trampling on the lamb, and the ewe has a good chance to get acquainted with her lamb at the start, thus avoiding the danger of disowned lambs later. These lambing pens should be in a well-ventilated room that is free from drafts and as warm as it can be made without artificial heat. In very cold weather a blanket thrown over the lambing pen will insure sufficient warmth to give the lamb a good chance in the first few hours, which are important ones.

TROUBLES AT PARTURITION.

Well-fed ewes seldom have much trouble in lambing, but there may always be need of assistance for a few ewes. If the ewe strains for half an hour without delivering the lamb, aid may then be given. The normal position of the lamb at birth is to have the forelegs extended with the head lying between them. If the lamb is not in the proper position, the shepherd should correct it by inserting the hand and arm into the vulva and effecting the change. When such assistance is needed the shepherd should first trim his finger nails and rub vaseline or oil upon his hand. In either case, when the position is correct the lamb can usually be successfully

¹ These panels are described in Farmers' Bulletin 810.

delivered by looping a string around the front feet and pulling outward and downward as the ewe strains. If the womb and vagina have been lacerated by the operation, it is well to use a solution composed of $\frac{1}{2}$ ounce of zinc sulphate and 2 ounces of tincture of opium in a quart of water at blood heat. This should be poured into the womb by means of a rubber tube and funnel. If the ewe seems weak a stimulant should be given.

WEAK LAMBS.

The lamb that is born strong and vigorous, with a good dam, will need little care. If the shepherd is present at the birth of a weak lamb, he should wipe away the phlegm or membrane from the nostrils of the lamb, and, if not already broken, the navel cord should be severed. Blowing into the mouth and nostrils and slapping gently on the ribs, first on one side and then on the other, will often save the life of a lamb that is apparently dead.

In cold weather lambs may get chilled and die unless prompt remedies are used. Wrapping the lamb in hot flannel cloths, which are renewed as often as necessary, is an excellent method of warming it. Another method is to place it for a few minutes in water as hot as the hand can bear; then remove, dry with cloths, and wrap up for an hour or two in fresh cloths or a sheepskin to complete the drying process. In any case milk should be given freely and the lamb returned to the ewe and allowed to suck as quickly as possible. If it does not suck when held to the teat, an infant's nursing bottle and nipple may be used. A few teaspoonfuls each hour for a few hours will usually give strength to enable the lamb to nurse without assistance.

DISOWNED LAMBS.

Little trouble is experienced with disowned lambs where lambing pens are used. With a ewe that refuses to own her lamb it is sometimes sufficient to draw some of the milk and rub it upon her nose and also upon the rump of the lamb. A heavy milking ewe with only one lamb can sometimes be made to adopt an orphan or the disowned lamb of a lighter milking ewe. When there is difficulty in having a ewe adopt another lamb after losing her own, the skin of the dead lamb may be fastened over the lamb to be adopted.

YOUNG-LAMB TROUBLES.

Well-nourished lambs from well-fed ewes have few troubles, but some troublesome conditions are to be expected in any flock. The causes and remedies of the more common ones are given below.

Constipation is indicated by straining and distress and may be remedied by a teaspoonful of castor oil. White scours can best be

cured by giving one-fourth of an ounce of cooking soda, 1 ounce of sulphate of magnesia, and a pinch of ginger in a small quantity of flaxseed tea or gruel. This should be followed in about four hours with 2 ounces of linseed oil. Indigestion is shown by distress and frothing at the mouth. A liberal dose of castor oil will effect a cure in most cases.

Sore eyes are of rather common occurrence. The eyes appear covered with a milky scum, or, in bad cases, become an angry red. In either case tears are apt to flow profusely. An eyewash of silver nitrate or 15 per cent argyrol will clear them up in a few applications. A very tiny drop of pure sheep dip is also recommended. Sore mouths are sometimes caused by scabs around the lips. These scabs should be rubbed off and sheep dip or a medium-strength solution of copper sulphate applied.

DOCKING THE LAMBS.

Docking, or removing the tail, is best done at the age of 10 to 14 days. When correctly done it adds much to the appearance and cleanliness of the lamb and raises the selling price at the market. For this purpose knives, either sharp or dull, chisels, and patented docking irons have all been used and recommended. When a sharp-edged tool is used the pain is slight, but unless some care is taken the lambs may lose considerable blood. Docking irons which burn through the tail may be used, and thus reduce the loss of blood to a minimum; but if used too hot the wound will be slow in healing. With any of these instruments the cut should be made about 1 inch from the body as measured on the underside of the tail. The lamb should be held with the rump resting upon the top of a panel or pen partition or upon a board if the hot irons are used. When docking with the hot iron the operator should work with the right hand, holding the tail in his left and pushing it toward the body. This will leave loose skin above the cut to close over the wound. Pine tar may be applied if flies are bad.

CASTRATION.

The ram lambs may well be castrated at the time they are docked. Both operations should be done early on a bright, cool morning. In castrating, the lamb is held in the same position as for docking. The hands and knife or shears should be disinfected. Unless both testicles can be felt, the operation should be delayed. The lower third of the scrotum should be cut off. The testicles then may be removed by pulling them straight out. In large flocks the testicles are removed by pulling with the teeth, as it is very difficult to grasp them with the fingers and it is necessary to do the work as quickly as possible.

TREATMENT OF EWES AFTER LAMBING.

The shepherd should watch the ewe's udder closely to see that it is in good condition, for good lambs can not be raised from ewes not milking freely. Ewes that have lambed should be kept in lambing pens from one to three days and then turned in a pen by themselves where they can be given special feed and care. After lambing they should be fed lightly at first, being put on full feed about the third or fourth day. At this time it is economy to feed heavily enough to produce a large flow of milk for the lambs. Heavy-milking ewes can make good use of from 1 to 2 pounds of grain per day. Experiments conducted at the Wisconsin experiment station showed that when ewes were on good pasture there was no extra gain made by the lambs when the ewes were fed grain.

THE FLOCK IN SUMMER.**SHEARING.**

Shearing is generally done in late spring or early summer, after lambing. It should be done on a warm day, so that the ewes may not become chilled. Formerly shearing was done mostly by the use of hand shears, but in most flocks of large size power shearing machines are now used. For small flocks under 50 head hand-power machines are the most economical. The machines are more rapid, smoother work is done, and the ewes are injured less. It is easier to learn to use them, and more wool is obtained than where hand shears are used.

The tags or dung locks should be removed from the fleece, and then it should be rolled up, not too tightly, skin side out, and tied with paper twine. Wool buyers prefer this method of tying to that done with wool boxes.

If the lambing is late the ewes may be sheared before lambing, but great care must be used in handling them. It is better to do the shearing after lambing. In either case it should be done before hot weather sets in.

DIPPING.

Sheep are dipped to free them from ticks, lice, and other skin parasites. A convenient time for dipping is shortly after shearing in the spring. Less dip per animal is needed and the weather is usually more favorable at this time than at any other season. The dipping should be done in the morning of a clear, quiet, warm day, so that the sheep will be dry by night and will not catch cold. Every member of the flock should be dipped, and it is well to spray the inside of the sheep barn with dip at this time. Any standard dip solution can be successfully used, if the manufacturer's directions are followed. To insure the eradication of sheep ticks the

sheep should be dipped a second time about 24 days after the first dipping. About 10 days should be allowed to elapse after shearing, so that shear cuts may have time to heal before dipping.

CULLING THE EWE FLOCK.

The summer or early fall, soon after the lambs have been weaned or marketed, is the best time to dispose of ewes that are not considered desirable for another year's breeding. The ewes that are to raise the next crop of lambs can then be prepared for fall breeding. Ewes of the mutton breeds do not ordinarily breed well nor keep in good condition after 5 years of age. Their usefulness, however, depends more upon the condition of their teeth than upon their actual age. Fine-wool ewes usually remain useful to a later age. It is a good plan to sell aged ewes before they become too run down to be valuable to the butcher. The ewes that give the most milk and raise the best lambs are likely to be quite thin at this time and should not be judged by their appearance.

Nonbreeding ewes, poor milkers, light shearers, and mothers of inferior lambs should be marked as their defects are discovered and should be disposed of at this time. Their places should be filled by the best individuals among the yearling ewes and from the best breeding older ewes.

WEANING THE LAMBS.

If lambs are sold at from 3 to 5 months of age, they may run with their dams until that time. The lambs to be kept for breeding pur-




FIG. 4.—A group of yearling ewes kept to replace the older ones culled out in the fall. These ewes have been cared for to produce good fleece and full development for use as breeders.



FIG. 5.—Sheep on good summer pasture with shade. Sheep need shade in summer days. They graze most toward evening and early in the morning. The sheep shown are part of a Louisiana grade flock having two crosses of mutton sires on native ewes. The average fleece weight of the native ewes was 3 pounds and of the three-quarter breds 5.6 pounds. (Photograph from Louisiana Extension Service.)

poses should be weaned at the same time and put on fresh pastures where there is no danger of stomach worms. When the weaning is done at this time the ewes can be put in better condition for the fall breeding. Ram lambs left in the flock worry the ewes and may get some of them in lamb. When lambs are to be kept on the farm the best method of weaning is to leave them on the old pasture for three or four days and remove the ewes to a scanty pasture to check their milk flow. As soon as the lambs cease fretting for their dams they may be moved to fresh pastures where the ewes have not been. Ewes with large udders should be partially milked once every three days until they go dry.

SUMMER PASTURES.

The breeding flock in summer needs little but good pasture, shade, salt, and plenty of fresh water. Bluegrass is one of the most popular pastures, but is likely to be too dry in late summer and too unbalanced in its food nutrients for ideal feed. It is at its best when used in the spring and fall and supplemented by forage crops in the summer. Alfalfa is sometimes pastured in the summer, but is better used when cut and fed as hay in the winter. There is serious danger of loss from bloating when sheep are grazed on alfalfa or clover. Sweet clover is worse than the red and alsike in this regard. Rape makes an excellent supplement for bluegrass, but is a forage crop rather than a summer pasture, though it may well supplement bluegrass. Soy beans are good, and if the flock is changed to another part of the field when most of the leaves have been eaten off, the plants will

make further growth for later use. Cowpeas are good for the older sheep, though unpalatable to lambs. Bermuda grass, when kept short, is especially good when reinforced by lespedeza and bur clover, which grow at different seasons from the Bermuda grass and here find their best use as a sheep pasture. The aftermath of grain and timothy fields furnishes feed for many flocks and helps greatly to bring down the cost of carrying the flock through the summer.

AVOIDING STOMACH WORMS.

In many farming sections the flockmaster's most serious troubles are likely to be caused by internal parasites, the effects of which are particularly evident during the later part of the pasture season. Of these parasites the stomach worm is the most common and troublesome. It occurs wherever climatic conditions and methods of keeping the flock are favorable for its development, which means on most farms, and probably all. No practicable means of entirely avoiding infection with this parasite has been discovered, but by proper arrangement of the summer pasturage it is possible to keep the numbers of the worms below the danger point. A knowledge of the development of this parasite affords a basis for the changing of pastures that insures a healthy condition of the flock.

The stomach worms live in the fourth stomach.¹ They are from one-half to 1¼ inches long and have a fine red stripe running in spirals from end to end of the body. Their eggs pass out in the droppings of the sheep and hatch in a few hours, days, or weeks, according as the temperature is high or low. At temperatures lower than about 40° F. development is arrested. The larva which hatches from the egg crawls up on the grass blades when they are moist, and after attaching itself to the blade may be swallowed by some animal. The eggs are frequently killed by freezing or drying, but the larvæ will sometimes live for months and can withstand repeated freezings. After being taken into the body of a ruminating animal they develop into the mature worms. Cattle and goats also act as hosts to these worms but usually are not so seriously affected as sheep.

The injurious action of stomach worms may be attributed to two things: First, the loss of blood extracted by the parasites and the loss of nutritive materials which may be absorbed by them from the alimentary fluids; and, second, the destruction of red corpuscles by a poisonous substance secreted by the parasites which is absorbed into the blood. Lambs that are affected become pale, thin, and weak and may either die or continue for a long time in poor condition and fail to grow as they should. The absorption of blood from the lambs

¹The information here given with regard to stomach worms is mostly taken from Bureau of Animal Industry Circular 157, written by B. H. Ransom.

by the parasites is most easily shown in the white paperlike appearance of the skin and membranes of the mouth and eyes, while watery swellings often develop under the jaws.

Treatment of infected lambs will bring about recovery if given in time, although, as before indicated, the safest and cheapest way of combating the trouble is by preventing it. Young lambs are very unlikely to become seriously infected by larvæ from eggs dropped by older sheep in barns or yards bare of grass. On a noninfected pasture the larvæ will not ordinarily develop in any considerable numbers to the stage which will result in injury in less than 10 days or two weeks. If the flock is moved to fresh noninfected ground by that time, the danger is avoided for a further period of the same length.

It is not known how long larvæ of this parasite will continue to be dangerous, but, since freezing commonly kills unhatched eggs, a pasture in cold climates that was not used in summer and fall until after frost will be practically safe for occupancy by lambs for a limited time the following spring or summer, provided the old sheep are removed from it before the winter is over. This fact, and the desirability of obtaining the maximum amount of grazing from small areas, thereby reducing the amount of fencing needed, makes it advisable to adopt the plan of having a rotation of forage crops for summer use. Land on which fall wheat or rye has been sown will be safe for spring use and, if plowed and sown to rape or other crops for later grazing, is then also free from serious stomach-worm infection.

On farms where sheep have not been previously kept trouble from stomach worms is not likely to be serious until the second or third summer.



FIG. 6.—A flock of lambs on the way to be loaded on cars for shipment to market. These lambs are the combined lots of several members of a Tennessee shipping club.

PREPARING LAMBS FOR MARKET.**ADVANTAGES OF EARLY MARKETING.**

Under ordinary farm conditions lambs should be made ready for market at from 3 to 5 months of age. When young they make a higher rate of gain and will put on the same amount of flesh for less cost than when they are older. Then, too, they will make but small gains during the heat of summer, and at this time parasites are most troublesome and they are thus more liable to losses from this cause. Risk of accidents is always higher when the lambs are held for a long time. More feed is saved for the breeding flock, and less labor is needed, if the lambs are sold early. Better prices are obtained in the spring because of not having to meet the competition of the western lambs that are marketed during the summer and fall, and in addition the grower gets the use of his money sooner by pushing the lambs to a marketable condition as fast as possible.

TEACHING THE LAMBS TO EAT.

Every effort should be made to keep the lambs growing from the start. The first essential is to teach them to eat. Liberal feeding of lambs dropped before pastures are ready is profitable under any ordinary grain prices. This is best done through the use of a small inclosure known as a "creep," to which the lambs have access at all times, but into which the ewes can not come. The creep should contain a rack for hay and a trough for grain, so arranged that the lambs can not get their feet into them.

All feed given, especially ground feed, should be clean, fresh, and free from mold. The lambs will begin to nibble at the feed when from 10 to 16 days of age. Pea-green alfalfa of the second or third cutting is one of the most relished feeds. Flaky, sweet wheat bran probably ranks next. For the first few days these are the ideal feeds. A little brown sugar on the bran at first will make it more palatable. Linseed meal is also good when mixed with the bran. Until the lambs are 5 to 6 weeks old all their feed should be coarse ground or crushed. The Ohio Experiment Station¹ has found that for young lambs that are to be marketed a grain ration of corn is of about the same value as one of corn 5 parts, oats 2 parts, bran 2 parts, and oil will be safe for spring use and, if plowed and sown to rape or other meal 1 part. Oil meal is especially relished by lambs at this time and would be especially valuable in promoting growth rather than fat.

Such feeds as middlings are too floury for extensive use. Rye is less palatable than oats or barley. Soy beans may replace the linseed meal if they cost less. Cleanliness is an important factor in keeping the lambs growing. Always feed to an empty trough, and if it becomes soiled scrub it out with limewater.

¹ Ohio Experiment Station Bulletin 270.

RAISING ON PASTURE ALONE.

The plan of having lambs dropped after ewes go to pasture and marketing them without the use of other feed for the flock can no longer be recommended for general use. The main advantages of this plan lie in the small amount of care needed and the lower feed cost. The cost, however, depends upon the quality of the pasture and the value of the land. Late lambs that have never received grain are particularly likely to be injured by stomach worms. Lambs make smaller gains in hot weather, and there is the possibility of droughts drying up the pastures and decreasing the ewes' milk at the time of the lambs' greatest need. Feeding grain to lambs on pasture is only partially satisfactory and is particularly unlikely to be profitable with lambs that have not learned to eat before going to pasture.

Unless grass is very good and cheap, and grain very high, this plan of raising lambs can not be expected to prove continuously profitable. When grass pastures are to be used for a flock turned out when the lambs are 5 to 8 weeks old, it is desirable to have sufficient divisions to allow frequent changes without the lambs being returned to any ground previously grazed in the same season. Lambs that are 6 weeks old when sent to pasture and have received some grain can withstand a considerable degree of parasitic infection.

THE DRY-LOT METHOD.

Some breeders of pure-bred sheep have practiced a dry-lot method of raising lambs, mainly to avoid stomach-worm troubles. Under this plan the lambs do not leave the sheds or yards until they are weaned, when they are put on clean, fresh pastures. In the meantime they are fed hay and grain, and their dams are returned from the pastures two or three times each day to allow the lambs to nurse. Because they do not graze, the lambs have slight chance of becoming seriously infected with stomach worms.

Some raisers of market lambs follow the plan of keeping both ewes and lambs in dry lots. This plan also prevents serious stomach-worm infection. Where green feeds or soiling crops are grown near by and fed in the lot, the ewes milk well and the lambs grow at a profitable rate. The main advantage from such a soiling system is that it insures freedom from injury by internal parasites. Less fencing is needed if the ewes can be grazed elsewhere after the lambs are sold. If this can not be done, as much fencing will be needed for the ewes in the fall as would have been required for the spring flock.

This plan is most likely to work well where alfalfa is the main crop. Feeding in the yards prevents loss from bloat, and there is no need for plowing the land, as would be necessary if sheep were to graze on it a number of times each season.

THE FORAGE-CROP METHOD.

The practice of grazing the flock on forage crops until the lambs are sold is becoming popular where lands are high in price and where stomach worms cause trouble. Under this plan the ewes and lambs are first grazed on fall-sown wheat or rye. The land is divided to avoid the necessity of keeping the flock longer than 10 days upon the same ground. By the time the second lot of this crop is grazed down, spring-sown peas and oats can be ready and the fall-wheat ground plowed and reseeded to another cereal or to rape or soy beans for later use. Such a plan requires some labor in preparing and seeding the land, but it produces the largest amount of feed per acre and prevents trouble from the stomach worms.

In 1915 three lots of ewes with lambs were reared at the Illinois Experiment Station to test the value of the grass-pastures, dry-lot and forage-crop methods. All of the lambs were dropped about the middle of March. The lambs running on grass ate an average of 0.3 pound of grain per head daily from March 27 to July 15, those in the dry lot 0.7 pound, and those on forages 0.3 pound. The gains made and the market value of the lambs when sold are shown in the following table:

Comparison of three methods of feeding lambs at the Illinois Experiment Station, 1915.

Method.	Average weight	Selling price.	Ratio of net returns.
	<i>Pounds.</i>		<i>Per cent.</i>
Dry-lot.....	66.1	{ 17 at \$8.00... 3 at \$7.00... }	100
Pasture (with grain).....	64.4	{ 15 at \$7.75... 5 at \$7.00... }	131.8
Forage.....	72.4	20 at \$8.50...	195.7

