BARD

FINAL REPORT

PROJECT NO. US-430-81

Study of Cytoplasmic Male Sterility in Lentil

F.J. Muehlbauer, G. Ladizinsky

1985

BARD
P.O.B. 6
Bet Dagan, ISRAEL

BARD No. of Proposal US-430-81

Title of Proposed Research

Study of cytoplasmic male sterility in lentil

Name(s) of Investigator(s)

Principal Investigator Fred J. Muehlbauer

Cooperating Investigator Gideon Ladizinsky

Name and Address of Affiliated Institution(s)

Grantee Institution Washington State University

Pullman, Washington, USA

Cooperating Institution The Hebrew University-Faculty of Agriculture

Rehovot, Israel

Project's Starting Date: November 11, 1982

Project's Duration 1 year

Type of Report: Initial ☐ Comprehensive ☐ Final ☒

Signature

Principal Investigator

Institution's Authorizing Official

Fred J. Muehlbauer

_____

Table of Contents

	Page
Abstract	1
Introduction	2
Methods and Procedures	3
Results and Discussion	3
References	5
Evaluation of research achievement with respect to the original proposal	6
Description of results of cooperation	6
Theses	6
Inventions	6
Published reprints and manuscripts	6
Table 1 Cultivars and accessions used for the survey of plasmotypes and sterility restorer genes in lentil and their origin . . .	7
Table 2 Summary of % stainable pollen in crosses and reciprocals of Tekoa and PI368648 with three unclassified parents	8
Table 3 Summary of % stainable pollen in a cross and reciprocals of Tekoa and PI368648 with ILL193	9
Table 4 Summary of % stainable pollen in crosses and reciprocals of Tekoa and PI368648 with seven unclassified parents	10
Table 5 Summary of % stainable pollen in crosses of Tekoa and PI368648 with unclassified parents	11
Table 6 Summary of % stainable pollen and seed set on male sterile plants in the field, 1984	13
Table 7 F ₂ progeny tests of three hybrids involving sterile cytoplasm and presumed restorer genes	14

Abstract

Hybridization in lentil (Lens culinaris Medik) is limited by small flowers that are difficult to emasculate and pollinate. As a result, progress in breeding has been hindered and hybrid cultivars are not possible. However, in lentil crosses, involving wild and cultivated forms, we found what appeared to be gametophytic cytoplasmic male sterility composed of sterile (S) and normal (N) cytoplasms, and maintainer (ms) and restorer (Ms) genes. Under the framework of this project, we attempted to determine the distribution of N and S cytoplasms and the Ms and ms alleles in cultivated types; also we attempted to assess the potential for cross pollination in lentil through the use of male sterility.

The results of the survey of cytoplasmic types in lentil indicated a complex system with at least three cytoplasms that were tentatively designated as S, and S₂ (both sterile), and N (normal), and a single restorer gene with three alleles (Ms₁, Ms₂ and ms). The field trial of fully sterile backcross-derived plants resulted in a very low percentage (0.7%) of outcrossed seed. That amount of outcrossing was not sufficient to justify development of hybrid cultivars, but the sterility may have some use in lentil breeding programs.

Introduction

The small cleistogamous flowers of lentil (*Lens* spp) are virtually 100% self pollinating. Hybridization is difficult to achieve because the small flowers are difficult to emasculate and pollinate without damaging the stigma, style and ovary. Methods of hybridization have been described by Wilson (1972), Malhotra et al. (1978), and Muehlbauer et al. (1980). Reported success in crossing has been over 80% of pollinations in the greenhouse; however, field success is much lower and often unsuccessful because of low relative humidity, difficulties in flower manipulation to avoid damage, and difficulties finding and in transferring viable pollen. In spite of the difficulties, breeding programs have succeeded in making the crosses necessary to meet objectives. Nevertheless, certain techniques such as cyclical recurrent selection and to some extent, backcrossing has been precluded because of the crossing problems. Techniques to circumvent the problem of crossing would help broaden the range of material and methods available to breeders.

Partial sterility and complete sterility have been observed (Muehlbauer, unpublished) in segregating populations of lentil (*L. culinaris*) crosses. Also, upon examination of certain reciprocal intraspecific crosses in *L. culinaris* and *L. orientalis*, some had 50% pollen fertility, while the reciprocals were fully male and female fertile. All the F_2 plants from fully fertile F_1 's were completely fertile whereas the F_2 's from hybrids with 50% pollen fertility showed a 1:1 segregation of fully fertile plants to 50% pollen fertile plants. Tentatively we assumed two cytoplasm, S and N and two alleles of one nuclear gene Ms and ms.

Based on this assumption, the genotypes N^{msms} and S^{MSMS} are fully male fertile. Upon crossing these two types, complete male fertile hybrids will be formed when the N^{msms} genotype serves as the female parent and only 50% of the pollen will be viable when the S^{MSMS} is the female parent. It appears that the sterility manifests itself at the gametophytic phase and therefore we have termed this sterility gametophytic cytoplasmic male sterility. The 1:1 ratio in the F_2 of the partially fertile F_1 hybrid is due to the fact that on the pollen side only the MS gametes are viable, but both MS and ms are viable and functional in the ovules.

In this system, complete male sterile plants can be obtained by backcrossing the F_1 hybrid as the female parent with the N^{msms} genotype.

Based on these observations, the objectives of the study were to: 1) Survey the distribution of the S and the N cytoplasm and the MS and ms genes in lentils in order to isolate stocks for future work and to get some idea about the role and the significance of the sterility; and 2) Assess the potential of cross pollination in lentil by the aid of cytoplasmic male sterile plants. Such information is essential in any attempt to use the sterility in recurrent selection breeding strategies or to produce hybrid cultivars.

Methods and Procedures

For the survey of cytoplasm types, maintainer and restorer genes, 45 germplasm accessions were chosen for hybridization with Tekoa (presumed to have the S cytoplasm and MsMs restorers) and PI368648 (presumed to have the N cytoplasm and msms restorers). The 45 accessions (Table 1) were chosen to represent a wide range of geographical origins and morphological types. Crosses and reciprocals were made between the 45 accessions and Tekoa and PI368648 in the greenhouse in the winter and spring of 1983. Because of crossing difficulties, it was not possible to obtain the entire set of crosses.

The F₁ hybrids were grown in the greenhouse during the summer and fall of 1983 and pollen collected from flowers at or near anthesis. Pollen was fixed in a 1:3 acetic acid-ethyl alcohol for 24 hours, transferred to 70% ethyl alcohol, and stored in the refrigerator until it could be analyzed for percentage stainable pollen. In order to make counts of percentage stainable pollen, the pollen was stained either with aceto-carmin or with potassium iodide (I₂KI). Counts of percentage stainable pollen were made under the microscope at about 45x magnification.

Hybrids showing 50% stainable pollen were backcrossed to their respective male parents (presumably N^{msms}) in order to develop completely male sterile plants (s^{msms}). Seeds expected to produce completely male sterile plants were germinated in the greenhouse during April, 1983 and transplanted to the field in May, 1983. The plants, expected to be completely male sterile, were transplanted within the yield trials for red cotyledon cultivars. Because the suspected male sterile lines had yellow cotyledons, outcrosses with the red cotyledon cultivars could be easily identified. To confirm male sterility of the plants, anthers from the male sterile lines were collected during early July, squashed, and stained with potassium iodide to determine percentage stainable pollen. At maturity, the plants were harvested and percentage seed set determined.

Results and Discussion

Crosses and reciprocals for the survey of cytoplasm types were analyzed for percentage stainable pollen and the results (Tables 2-5) indicated a more complex system than originally believed. Based on the data, there appear to be at least three cytoplasm types i.e. the normal, the sterile, and the one carried by PI368648. Initially, we considered PI368648 to have the normal cytoplasm (N^{msms}) and Tekoa to have the sterile cytoplasm (S^{MsMs}). The crosses shown in Table 2 were expected if the initial evaluations were true and if the accessions used (PI360630, PI251786, and ILL8) had the normal (N) cytoplasm. The possible mechanisms is as follows:

Tekoa	S^{MsMs}	X	N^{msms}	
		↓	S^{msms} (cross)	50% stainable pollen
			N^{Msms} (reciprocal)	100% stainable pollen
PI368648	N^{msms}	X	N^{msms}	
		↓	N^{msms} (cross)	
			N^{msms} (reciprocal)	100% stainable pollen

However, when ILL193 was crossed to Tekoa and PI368648, PI368648 also appeared to have a sterile cytoplasm (Table 3) since, when PI368648 was used as a female, the resulting hybrid had about 50% stainable pollen.

A number of crosses had a high percentage of stainable pollen regardless of the direction of the cross (Table 4). This indicated that the lines used in the crosses likely had the normal cytoplasm and the sterility restorer genes as well. The possible mechanism is as follows:

Tekoa	S^{MsMs}	X	N^{MsMs}	
		↓	S^{MsMs}	(Cross)
			N^{MsMs}	(Reciprocal)
				Both fully fertile
PI368648	N^{msms}	X	N^{MsMs}	
		↓	N^{Msms}	(Cross)
			N^{Msms}	(Reciprocal)
				Both fully fertile

Other crosses made for the survey of cytoplasm types are summarized in Table 5. In those crosses we were unsuccessful in obtaining a complete set of reciprocals, however, in a number of cases PI368648 appeared to have a sterile cytoplasm.

The survey results seem to indicate three different cytoplasm types and a single nuclear restorer gene with three or more alleles.

Based on that explanation, the genotypes of three accessions would be as follows:

Tekoa	S_1	$Ms_1 Ms_1$
PI368648	S_2	$Ms_2 Ms_2$
ILL 193	N	$msms$

The dominance relations among the three alleles would be: Ms_1 -- Ms_2 -- ms . However, the dominance is apparently not always complete. Thus in the Tekoa x PI368648 ($S_1^{Ms_1Ms_2}$) hybrid, the Ms_2 allele is eliminated in the pollen, but in the reciprocal cross ($S_2^{Ms_2Ms_1}$) both alleles are viable because of the dominance of Ms_1 .

There is still some difficulty in explaining the behavior of the Tekoa X ILL4354 cross (Table 4), and there are a few cases where 50% stainable pollen was recorded for both cross directions. This may have been the result of chromosomal rearrangements between the parents.

The field trial of fully sterile backcross-derived plants showed that we can expect to get very low percentages of outcrossed seeds through the use of cytoplasmic male sterility, at least in the environment of the Palouse (Table 6). All cytoplasmic male sterile plants had 0% stainable pollen, as expected, and flowered normally during the growing season. Because the sterile plants were grown among red cotyledon cultivars, we expected that the outcrossed seeds would all be red cotyledon, since that trait is controlled by a single dominant gene. The fact that more yellow cotyledon seeds were obtained was surprising. This may have resulted from insects carrying pollen from greater distances. The yellow cotyledon cultivar nursery was within 100m and that may have been the source of pollen for those apparent crosses. However, it is possible that the yellow cotyledon seeds were the result of the plants regaining some fertility. The outcrossed seeds were located at random on the otherwise male sterile plants.

Overall, we obtained a total of 21 presumed outcrossed seeds from slightly more than 3000 flowers. Because lentils average about one seed per pod, the actual outcross percentage was estimated to be about 0.7%. That degree of outcrossing would be insufficient to justify hybrid seed for lentil production and may also be too low to justify the use of cytoplasmic male sterility in a lentil breeding program. The use of this sterility in some type of cyclical recurrent selection scheme might be feasible.

Progeny tests of F_2 plants from three hybrids are shown in Table 7. In all three crosses we expected a 1:2:1 ratio between 100% fertile:50% fertile:0% fertile plants. Obviously, we did not have a large enough population from either cross; however, overall the ratio was 13:22:7 and approximated a 1:2:1 ratio.

The cytoplasmic male sterility-restorer system in lentils is an interesting trait because of its' possible evolutionary significance. It is likely that lentil was at one time an outcrossing species or at least partially so and through evolution selfing has come to be rigidly enforced through flower structure and possibly the loss of insect attraction mechanisms.

References

1. Wilson, V. E. 1972. Morphology and technique for crossing Lens esculenta Moench. Crop Sci. 12:231-232.
2. Malhotra, R. S., H. S. Balyan, and P. R. Gupta. 1978. Crossing technique in lentils. Lentil Exp. News Serv. 5:7-8.

3. Muehlbauer, F. J., A. E. Slinkard and V. E. Wilson. 1980. Chapter 29. Lentil. In Hybridization of Crop Plants. American Society of Agronomy Monograph, Madison, Wisconsin. pp 417-426.
4. Muehlbauer, F. J. unpublished.

Evaluation of research achievement with respect to the original proposal

We originally proposed to complete the genetic analysis of the cytoplasmic male sterility restorer system in lentils, survey the distribution of the two cytoplasms and their respective restorer and maintenance genes. We also wanted to assess the potential use of male sterility for producing hybrid cultivars, and facilitating hybridization in breeding programs. Technical difficulties at the beginning of the project resulted in the loss of some genetic material and led to the request for project extension. The extension of one year made it possible to complete the survey of cytoplasm types and to carry out the field evaluation of outcrossing potential using cytoplasmic male sterility.

The survey of cytoplasm types revealed a more complex system than originally thought. The data on outcrossing potential in lentil using male sterility is probably the most accurate estimate of naturally outcrossing made in lentil to date. Unfortunately, the percentage is so low as to preclude the use of male sterility for hybrid seed production; however, it does appear to have potential for breeding, especially for methods that require numerous cross pollinations.

Overall, the major objectives of the project were achieved and new insights as to the significance of cytoplasmic male sterility in lentil were obtained. We plan to complete the genetic analysis and also will attempt to obtain another year's data on outcrossing potential using male sterility.

Description of results of cooperation

Cooperation between U. S. and Israeli counterparts made it possible to formulate effective approaches to the study of the cytoplasmic male sterility-restorer system in lentil. Potential uses of the system in breeding and for seed production were evaluated effectively because of this cooperation. The understanding we now have of the system is important to breeders of this crop world wide.

Theses

Thompson, Joan. Cytoplasmic male sterility in Lens culinaris Medik. M. Sc. thesis. Washington State University, Pullman, Washington, USA (in preparation)

Inventions

None

Published reprints and manuscripts

Further work needs to be done to clarify the cytoplasms involved and the genetics of fertility restoration before publications can be prepared. We are planning to do this. and to publish the results.

Table 1. Cultivars and accessions used for the survey of plasmotypes and sterility restorer genes in lentil and their origin.

Cultivar or accession	Origin
Brewer	USA
Redchief	USA
Laird	Canada
Eston	Canada
Precoz	Argentina
Giza 9	Egypt
LC000010	USA
LC000017	USA
LC000028	USA
Benewah	USA
Chilean 78	USA
Ramtha	Jordan
ILL 101	Morocco
ILL 254	Greece
ILL 500	Mexico
ILL 504	USSR
ILL 1861	Sudan
ILL 1877	Turkey
ILL 4351	Iran
ILL 4354	Jordan
ILL 4380	India
ILL 4399	Lebanon
ILL 4400	Syria
ILL 4401	Syria
ILL 8	Jordan
ILL 15	Jordan
ILL 28	Syria
ILL 30	Syria
ILL 193	Syria
ILL 262	Hungary
ILL 419	Chile
ILL 707	Cyprus
ILL 915	Spain
ILL 983	Chile
ILL 1042	Iran
PI212610	Afghanistan
PI251768	USSR
PI297745	Greece
PI12611	Afghanistan
PI212903	Egypt
PI343025	USSR
PI370632	Turkey
PI379368	Yugoslavia
ILL 1880	Turkey
PI370630	Turkey

Table 2. Summary of % stainable pollen in crosses and reciprocals of Tekoa and PI368648 with three unclassified parents.

Pedigree	Stainable Pollen	
	Cross	Reciprocal
	- - - - - % - - - - -	
Tekoa/PI370630	69.4	96.2
PI368648/PI370630	96.2	98.1
Tekoa/PI251786	59.0	97.7
PI368648/PI251786	96.4	99.3
Tekoa/ILL8	56.7	80.4
PI368648/ILL8	83.1	86.5

Mechanism:

Tekoa	S ^{Ms} Ms	X	N ^{ms} ms	
Cross		S ^{Ms} ms		50% stainable pollen
Reciprocal		N ^{Ms} ms		100% stainable pollen
PI368648	N ^{ms} ms	X	N ^{ms} ms	
Cross		N ^{ms} ms		100% stainable pollen
Reciprocal		N ^{ms} ms		

Table 3. Summary of % stainable pollen in a cross and reciprocals of Tekoa and PI368648 with ILL193.

Pedigree	Stainable Pollen	
	Cross	Reciprocal
	----- % -----	
Tekoa/ILL193	52.0	100.0
PI368648/ILL193	55.3	93.9

Table 4. Summary of % stainable pollen in crosses and reciprocals of Tekoa and PI368648 with seven unclassified parents.

Pedigree	Stainable Pollen	
	Cross	Reciprocal
	----- % -----	
Tekoa/ILL4399	97.1	98.4
PI368648/ILL4399	97.4	100.0
Tekoa/ILL101	86.1	88.1
PI368648/ILL101	89.7	89.4
Tekoa/Benewah	98.7	100.0
PI368648/Benewah	98.1	95.0
Tekoa/PI297745	90.2	92.7
PI368648/PI297745	99.0	95.6
Tekoa/ILL1877	92.1	98.3
PI368648/ILL1877	77.5	99.0
Tekoa/ILL4351	71.4	99.4
PI368648/ILL4351	98.4	97.4
Tekoa/Precoz	98.8	97.4
PI368648/Precoz	97.1	99.7

Mechanism:

Table 5. Summary of % stainable pollen in crosses of Tekoa and PI368648 with unclassified parents.

Pedigree	Stainable Pollen	
	Cross	Reciprocal
	----- % -----	
Tekoa/ILL254	----	72.1
PI368648/ILL154	99.0	93.2
Tekoa/ILL915	----	81.3
PI368648/ILL915	85.0	89.8
Tekoa/ILL419	----	98.4
PI368648/ILL419	52.9	53.1
Tekoa/ILL504	----	100.0
PI368648/ILL504	97.9	100.0
Tekoa/PI212611	75.7	99.4
PI368648/PI212611	----	----
Tekoa/Chilean 78	----	96.0
PI368648/Chilean 78	99.0	----
Tekoa/ILL262	----	69.2
PI368648/ILL262	99.4	----
Tekoa/ILL28	----	84.6
PI368648/ILL28	47.1	----
Tekoa/ILL707	----	98.7
PI368648/ILL707	95.9	95.6
Tekoa/Giza 9	96.4	99.0
PI368648/Giza 9	----	99.7
Tekoa/Redchief	59.6	99.3
PI368648/Redchief	97.7	----
Tekoa/Eston	86.5	----
PI368648/Eston	97.2	99.7
Tekoa/ILL4400	----	71.5
PI368648/ILL4400	48.9	52.5
Tekoa/ILL4401	97.2	99.4
PI368648/ILL4401	41.4	----
Tekoa/Laird	79.0	----
PI368648/Laird	----	99.7

Table 5. Summary of % stainable pollen in crosses of Tekoa and PI368648 with unclassified parents (continued).

Pedigree	Cross	Stainable Pollen	
			Reciprocal
		- - - - - % - - - - -	
Tekoa/ILL4354	----		57.7
PI368648/ILL4354	97.1		89.5
Tekoa/PI370632	----		99.3
PI368648/PI360632	97.0		99.3
Tekoa/PI212902	83.4		----
PI368648/PI212902	----		----
Tekoa/ILL15	----		36.6
PI368648/ILL15	55.6		53.6
Tekoa/Brewer	93.6		----
PI368648/Brewer	99.0		99.0
Tekoa/PI343025	99.0		99.4
PI368648/PI343025	96.9		----
Tekoa/ILL4351	71.4		99.4
PI368648/ILL4351	98.4		97.4
Tekoa/ILL4380	----		----
PI368468/ILL4380	98.2		99.7
Tekoa/ILL30	----		59.2
PI368648/ILL30	59.0		16.0
Tekoa/ILL1861	----		----
PI368648/ILL1861	97.5		99.0

Table 6. Summary of % stainable pollen and seed set on male sterile plants in the field, 1984.

Plant Number	Stainable	No. Flowers	No. Seeds	Seed Set ^{1/}	Cotyledon Color	
					red	yellow
	---%---			---%---		
Rep I						
172-1-1	0	33	0	0	-	-
172-1-2	0	29	0	0	-	-
172-2-1	0	98	0	0	-	-
172-2-2	0	210	0	0	-	-
172-5-1	0	136	0	0	-	-
172-5-2	0	164	0	0	-	-
172-5-3	0	94	0	0	-	-
172-6-1	0	160	0	0	-	-
172-6-2	0	96	0	0	-	-
172-6-3	0	133	3	2.3	2	1
172-6-4	0	24	0	0	-	-
	Total	1177	3	0.3		
Rep II						
172-2-3	0	131	2	1.5	-	2
172-2-4	0	86	1	1.2	-	1
172-2-5	0	12	0	0	-	-
172-5-4	0	150	2	1.3	-	2
172-5-5	0	253	0	0	-	-
172-5-6	0	53	0	0	-	-
172-6-5	0	120	2	1.7	2	-
172-6-6	0	53	2	3.8	-	2
172-6-7	0	98	5	5.1	-	5
172-6-8	0	73	0	0	-	-
172-6-9	0	41	2	4.9	-	2
	Total	1070	16	1.5		
Rep III						
172-5- 7	0	174	1	0.6	-	1
172-5- 8	0	71	0	0	-	-
172-6- 10	0	243	0	0	-	-
172-6- 11	0	118	0	0	-	-
172-6- 12	0	34	0	0	-	-
172-6- 13	0	77	1	1.3	-	1
172-6- 14	0	37	0	0	-	-
	Total	754	2	0.3		

^{1/}One pod per flower and assuming an average of one seed per pod.

Table 7. F₂ progeny tests of three hybrids involving sterile cytoplasm and presumed restorer genes.

Pedigree	Presumed genotype of F ₁	% stainable pollen ^{1/}		
		100	50	0
--No. of F ₂ Plants--				
Tekoa/PI370630	$\begin{matrix} Ms_1 ms \\ S_1 \end{matrix}$	1	9	6
Tekoa/PI251786	$\begin{matrix} Ms_1 ms \\ S_1 \end{matrix}$	5	6	1
PI368648/ILL193	$\begin{matrix} Ms_2 ms \\ S_2 \end{matrix}$	$\frac{7}{13}$	$\frac{7}{22}$	$\frac{--}{7}$

^{1/}Approximate percentage of % stainable pollen.