14

77

BARD

FINAL REPORT

PROJECT NO. US-869-84

Antigenic and Biological Characterization of Potyviruses that Infect Cucurbits

RECEIVED
1989-01-17

D. E. Purcifull, B. Raccach, R. Yokomi, S. Cohen,
E. Hiebert

3

1989

30758

000000101431

Cover page
BARD
P.O.B. 6
Bet Dagan, Israel

BARD Research Project: US-869-84

Title: Antigenic and Biological Characterization of Potyviruses That Infect Cucurbits

Name and Address of Investigators:

Principal Investigator: D. E. Purcifull
Dept. Plant Pathology
University of Florida
Gainesville, FL 32611, USA

Co-Principal Investigators:

Dr. B. Raccach
Dept. Virology
Agricultural Research Org.
Volcani Center
P.O.B. 6
Bet Dagan, Israel

Dr. S. Cohen
Dept. Virology
Agricultural Research Org.
Volcani Center
P.O.B. 6
Bet Dagan, Israel

Dr. R. Yokomi
U.S. Department of Agriculture
2120 Camden Road
Orlando, FL 32803, USA

Dr. E. Hiebert
Department of Plant Pathology
University of Florida
Gainesville, FL 32611, USA

Name and Address of Affiliated Institutions:

Institute of Food & Agr. Sciences
University of Florida
Gainesville, FL 32611, USA

Agricultural Research Org.
Volcani Center
Bet Dagan, Israel

Project Commencement Date: September 14, 1985

Type of Report: Final

Signatures:

Principal Investigator

Alan E. Purcifull

Institution's Authorized
Official (Signature & Stamp)

Dillard C. Marshall

DILLARD MARSHALL
ASSISTANT DIRECTOR OF RESEARCH

630.72, 635.61; 576.23; 632.8
BAR/PUR
and copy

B. Table of Contents

C. Abstract	page 3
D. Objectives of the original research proposal	page 4
E. Research report	page 4
F. Description of cooperation	page 76
G. Evaluation of research achievements with respect to aims of the original proposal	page 76
H. List of publications, including Appendix I	page 78

C. Abstract

Polyclonal antisera were prepared in rabbits to purified viruses, capsid proteins, and cylindrical inclusion proteins associated with three potyviruses that infect cucurbits, papaya ringspot virus type W (PRSV-W), zucchini yellow mosaic virus (ZYMV), and watermelon mosaic virus 2 (WMV-2). These antisera were used in studies of virus relationships, epidemiology, host range, *in vitro* translation, diagnosis, and assessment of disease control measures. Polyclonal antisera also were prepared to ZYMV helper protein and to PRSV-W amorphous inclusion protein (AIP). Monoclonal antibodies were produced to PRSV-W and its capsid protein, to ZYMV, WMV-2, and PRSV-W AIP. Serological tests showed that of 549 cucurbit samples collected in various parts of Florida in 1987 and 1988, 44.6% were infected with PRSV-W, 37.1% with WMV-2, and 9.2% with ZYMV. A filamentous virus serologically related to a potexvirus was detected only in samples of *Coccinea* and *Trichosanthes*. Antigenic variation in isolates of PRSV-W was detected by immunodiffusion tests with polyclonal antisera and with monoclonal antibodies in enzyme-linked immunosorbent assays. Biological variability was detected in PRSV-W and ZYMV by host range studies. Aphid-transmissible (AT) and non-transmissible (NAT) isolates of ZYMV were found in Israel. An NAT isolate produced active helper component which mediated transmission of an AT isolate. In reciprocal tests, potato virus Y (PVY) helper component readily assisted transmission of ZYMV but ZYMV helper inefficiently assisted PVY. The ZYMV is widespread in Israel and is the most important potyvirus of cucurbits there. Epidemiological studies provided evidence that colonizing aphids such as *Aphis gossypii* possibly contribute significantly to secondary spread of ZYMV, although other aphids, such as *A. citricola* may be important in primary infections. A combination spray of an oil (Virol) and a pyrethroid (Maverick) increased the yield of marketable melons. The spray treatments are now recommended for control of ZYMV in Israel. The inheritance of resistance to ZYMV in the interspecific cross of *Cucurbita maxima* X *C. ecuadorensis* was studied.

D. Objectives of the original research proposal.

1. To develop, through the production of monoclonal antibodies and conventional antisera, serodiagnostic tests that will distinguish ZYMV, PRSV-W and WMV-2. These antisera and tests will be used in epidemiological studies.

2. To develop immunological probes for virus-specified, nonstructural proteins associated with infections caused by WMV-1, WMV-2, and ZYMV (e.g. cylindrical inclusion proteins, and aphid transmission helper component proteins), to be used in in vitro translation studies for mapping the genes of these potyviruses, and also for serodiagnostic purposes.

3. To determine wild hosts and alternative crop hosts of principal potyviruses that infect cucurbits in Israel and Florida and to assess the suitability of these hosts as natural virus sources using aphid vectors.

4. To assess the variability of ZYMV, PRSV-W, and WMV-2 strains with regard to host range, host reaction and antigenicity.

5. To determine the major vector aphid species which are responsible for PRSV-W, WMV-2, and ZYMV transmission using the methods described by Raccah (1983). These major vectors would then be used to test the transmissibility of different isolates of the three viruses, using purified virus and helper component protein.

The overall objective of this project was to provide information that would reduce the number and intensities of ZYMV, PRSV-W, and WMV-2 epidemics in cucurbits.

E. Research report

Introduction

The potyvirus group is comprised of some 75–100 plant viruses, each of which has filamentous particles 720–900 nm long; all potyviruses that have been examined induce characteristic cytoplasmic, cylindrical (pinwheel), proteinaceous inclusion bodies in their hosts (Edwardson, 1974; Hollings and Brunt, 1981). Most members of the group are transmitted by aphids. Several distinct potyviruses cause major diseases of cucurbit crops (Lovisolo, 1980). Papaya ringspot virus type W (PRSV-W), formerly named watermelon mosaic virus 1 (Purcifull et al., 1984a), and watermelon mosaic virus 2 (WMV-2) (Purcifull et al., 1984b) have been recognized as important pathogens of cucurbits throughout the world for many years (Anderson, 1954; Grogan et al., 1959; Van Regenmortel, et al., 1962; Webb et al., 1965; Greber, 1978; Lovisolo, 1980; Makkouk and Lesemann, 1980; Purcifull et al., 1984b, 1984c). Zucchini yellow mosaic virus (ZYMV) (Lisa and Lecoq, 1984), which was originally described in Italy (Lisa et al., 1981), has recently been found to cause serious diseases of cucurbits in France and other Mediterranean areas, including Israel and Lebanon (Lecoq et al., 1983; Lesemann et al., 1983) and the U.S.A. (Purcifull, et al., 1984a; Provvidenti, et al., 1984; Nameth, et al., 1984). All three potyviruses induce diseases on squash, cantaloupe, cucumber, and watermelon. Losses caused by these viruses result from reduced production and reduced fruit quality (Provvidenti and Schroeder, 1970; Adlerz et al., 1983; Provvidenti et al., 1984). PRSV-W, WMV-2, and ZYMV are not known to be seed-borne in cucurbits (Grogan et al., 1959; Hein, 1977; Lecoq et al., 1981), and therefore it is assumed that the viruses are brought into the crop by incoming aphids which acquire the viruses from alternative crops or from infected wild plants. Adlerz (1977) has shown a correlation between WMV-2 infection and occurrence of *A. citricola* in Florida. This same aphid species is common in Israel and it was found to be the most important vector for the transmission of PVY to peppers in 1980 and in 1981 (Raccah, 1983; Raccah et al., 1985). The determination of aphid species which are involved in the spread of these viruses in Israel and in Florida is needed in order to evaluate the transmissibility of each of the viruses in the laboratory and to relate it to isolates in the field.

The epidemiology of PRSV-W, WMV-2, and ZYMV in Israel and Florida is poorly

understood. Breeding programs designed to produce resistant varieties are affected by variations in the virulence of virus strains and it is important to characterize natural virus variants and to use them in evaluating potential sources of resistance. Weed and noncucurbit crop hosts in which the viruses may survive from season to season, and factors affecting aphid transmissibility of the viruses are not well known. Such information will be useful in the implementation of disease control programs.

Serological tests have provided one of the chief means of identifying potyviruses of cucurbits (Lovisolo, 1980). The serological relationships among PRSV-W, WMV-2, and ZYMV have been somewhat controversial. Webb and Scott (1965) reported the PRSV-W and WMV-2 were antigenically distinct and had distinctive host ranges. Milne and Grogan (1969) reported that PRSV-W and WMV-2, and papaya ringspot virus type P (PRSV-P) are all closely related serologically. There is now considerable evidence, however, that PRSV-W and WMV-2 are antigenically, biologically, and cytologically distinct (Purcifull and Hiebert, 1979; Russo, *et al.*, 1979). PRSV-W and PRSV-P are regarded to be very similar with respect to serological properties (Purcifull and Hiebert, 1979; Gonsalves and Ishii, 1980; Yeh, 1984). PRSV-W has been classified as a strain of PRSV, type W (infects cucurbits, but not papaya), whereas type P strains of PRSV infect both cucurbits and papaya (Purcifull *et al.*, 1984). ZYMV is serologically related to WMV-2, but demonstration of these relationships depends on the antiserum used and the type of test employed (Lisa, *et al.*, 1981; Purcifull, *et al.*, 1984b; 1984c; Davis and Yilmaz, 1984). Definitive serological tests are needed for virus diagnosis in order to proceed effectively with epidemiological studies and breeding programs designed to evaluate resistance of cucurbit cultivars.

Materials and Methods

Purification of immunogen and production of polyclonal antisera. In Florida, PRSV-W, Adlerz Florida isolate, its amorphous inclusions, and its cylindrical inclusion body proteins were purified by previously published methods (Purcifull *et al.*, 1984a). WMV-2 and its cylindrical inclusions also were purified as previously described (Purcifull *et al.*, 1984b). Zucchini yellow mosaic virus (ZYMV) was purified as follows from systemically infected squash leaves collected 3 weeks after inoculation. Homogenize 100 g tissue in 200 ml 0.02 M Hepes buffer containing 10 mM sodium EDTA and 0.2 g sodium sulfite. Emulsify the extract with 100 ml each of carbon tetrachloride (CCl₄) and chloroform, respectively. Centrifuge at 2500 RPM (Servall GSA rotor) for 5 minutes. Filter the supernatant through 5 layers of cheesecloth. Resuspend the pellets in 50 ml of 0.02M Hepes buffer, and repeat filtration and centrifugation. Combine the supernatants, and centrifuge at 8000 RPM for 14 min (GSA rotor). At this step, save both the supernatant and the pellets. Resuspend the pellets for the purification of cylindrical inclusions (CI). To the supernatant, add 1% Triton X-100, 6% polyethylene glycol (PEG), and 100mM sodium chloride, and stir at 4C for 30 min (Servall SS34). Centrifuge at 15000 RPM for 10 min, and resuspend the pellet. After low speed centrifugation, layer the supernatant on cesium sulfate gradients (10 g cesium sulfate + 27 ml of .02M Hepes buffer, pH 8.2). Centrifuge at 32k RPM overnight (SW 41 Beckman rotor). Collect the band containing virus and add PEG to the collected material to precipitate the virus. Finally, resuspend the virus in 0.02M Hepes containing 10 mM EDTA, pH 8.2 or 0.02 M Tris buffer, pH 8.2, containing 10mM EDTA. To continue the purification of CI, add 5% Triton X-100, and stir at 4C for one hour. Centrifuge at 12k RPM for 10 min (SS34 rotor), and resuspend the pellets in 0.02M Hepes buffer, pH 7.5, containing 0.1% 2-mercaptoethanol. Centrifuge and resuspend in this manner three times. Finally resuspend the pellets in 0.02 M Tris buffer, pH 8.2. The CI, virus, and amorphous inclusion (AI) preparation were stored frozen. The CI proteins, AI protein, and capsid proteins were prepared and purified from frozen preparations by polyacrylamide gel electrophoresis as described by Hiebert *et al.* (1984).

A filamentous virus (TV) isolated in 1986 (FC-1860) from the cucurbit *Trichosanthes* in

Florida was purified for preparation of antisera. The FC-1860 isolate was cultured in small sugar pumpkin by mechanical inoculation. One hundred grams of tissue, 18 days after inoculation, was homogenized in 200 ml buffer (pH 7.5) consisting of 0.5 M potassium phosphate, 10 mM EDTA, and 1 gm sodium sulfite. The homogenate was expressed through cheesecloth. The expressed homogenate was centrifuged at 1020 g (max) for 5 min. The supernatant was homogenized with 150 ml chloroform and 150 ml *n*-butanol. The homogenate was centrifuged at 1020 g as before. The virions were precipitated from the supernatant by adding 6% PEG (final concn) (MW6000), sodium chloride to 0.1 M, and Triton X-100 to 1% concn, stirring for 1 hr, and centrifuging at 11,700 g for 10 min. The pellets were resuspended in 15 ml of resuspension buffer (RB) (pH 8.0) containing 20 mM potassium phosphate and 10 mM EDTA. The resuspended material was centrifuged at 7,700 g for 10 min. The birefringent supernatant was layered onto 30% CsCl in RB (5 ml supernatant/5 ml CsCl in a Beckman SW 41 rotor tube) and centrifuged at 140,000 g for 16 hr (5° C). The fraction layered 12 to 19 mm from the bottom of the tube was collected by droplet fractionation. The collected material was diluted 1:1 with RB and centrifuged 12,100 g for 10 min. The virions were concentrated from the supernatant by precipitation with 6% PEG and centrifugation at 7,700 g for 5 min. The pellet was resuspended in 2 ml RB. The resuspended material had a 260/280 ratio of 1.37 and was estimated to contain 35 mg of virus.

In Israel, purification procedures were developed for ZYMV, which is the most common potyvirus found in cucurbits in Israel. The isolate chosen was a wild type collected in the field, found to be highly transmissible and having a host range similar to that of isolate C-4. Later, all the other isolates were purified by the same method. C-13 yielded the highest. The potyviruses were raised in zucchini plants, and tissue was harvested between 2 and 3 weeks after inoculation. Several purification procedures were attempted, but the method used was that of Moghal and Francki (1976). The method was modified by excluding the polyethylene glycol step (PEG). This was found to be of advantage when the virus was purified from zucchini plants, where PEG tended to increase the accumulation of a slime-like pellet which increased aggregation of virus and caused great loss in virus yield. Tissue was homogenized in buffer, thioglycolic acid, CCl_4 , and chloroform. After the separation of the crude leaf material and organic solvent layer, the aqueous fraction was further clarified by a second low speed centrifugation followed by a high speed one on a cushion of 20% sucrose. The pellet was resuspended overnight by gentle stirring in the cold, and layered on a 10-40% sucrose gradient the next day. The virus band obtained was concentrated and tested for purity by spectrophotometry. If the band was not pure enough, the sucrose gradient step was repeated.

Antisera were prepared to Florida isolates by immunizing rabbits and collecting antisera using procedures similar to those described elsewhere (Purcifull and Batchelor, 1977). Immunogen consisted of purified virus, gel-purified proteins (Hiebert *et al.*, 1984), or SDS-treated virus (Table 1). In Israel, antiserum was prepared to one ZYMV isolate and to isolate MC.

An antiserum also was prepared in Israel to helper protein of ZYMV, which was purified as follows:

Extract 1 g zucchini leaves in 3 ml in 0.3 M potassium phosphate buffer pH 9.0. Follow by low speed centrifugation at 5000 g for 10 min to remove debris, and the virions were removed by ultracentrifugation at 100,000 g for 2 h. Only the top 1/3 of the supernate volume was used for further purification (after taking an aliquot for testing aphid transmissibility). Then 30% ammonium sulphate was added to the supernate and the solution was allowed to stand for 60 min in the cold. The pellet was collected after 8000g centrifugation for 10 min. The pellet was resuspended in 1/10 of the original supernatant volume, using an all glass homogenizer, and allowed to resuspend further in the cold for 30 min. The solution was

centrifuged again at 10000g for 10 min, and sucrose was dissolved in it to 20% (w/v). Helper was loaded on each lane of 5–15% gradient of polyacrylamide gel. The HC fraction was resolved by SDS PAGE stained in 0.2% coomassie brilliant blue and the band of HC about 54K was composed of the band obtained by immunoprecipitation with antibodies prepared and kindly provided by Drs. T.P. Pirone and D.W. Thornbury. The bands from 6–8 lanes of gel were excised and extensively washed in distilled water 3 consecutive times in 10 ml each for 4 h. The washed gel bands were homogenized in 1 ml PBS (phosphate saline buffer) and emulsified with an equal volume of Freund incomplete adjuvant. The emulsion was injected subcutaneously into a rabbit along the back in at least in six places. This procedure was repeated four times at biweekly intervals and then a fifth booster injection was administered. The rabbit was bled and the serum fractionated from blood. The serum was kept at -20°C and used for immunoprecipitation and biological assay.

Preparation of monoclonal antibodies to ZYMV and WMV-2.

In vivo immunization:

In all cases, 4 Balb/c mice were immunized on day 1 with a subcutaneous (SC) injection of 50 μg purified virus in 0.5 ml distilled water, emulsified 1:1 in Freund's complete adjuvant (FCA). The subsequent injections also were administered intraperitoneally (IP), using 50 μg emulsified 1:1 in Freund's incomplete adjuvant (FIA). One mouse was selected for a fusion based on positive results from a test bleeding. The final injection, 0.25 μg in 0.5 ml water without adjuvant, was given directly into the tail vein (intravenous, IV). If the entire volume could not be injected into the tail vein, the remainder was injected IP. Three to five days after the IV injection the mouse was sacrificed for a fusion.

In vitro immunization:

Two types of *in vitro* fusion procedures were used as modified from that outlined by Boss (1986), Wieggers *et al.* (1986), and S. Zam (personal comm.). Two mice were either (a) previously immunized as described above or (b) not previously immunized. In either case, the final IV immunization was administered by *in vitro* culture to 1 or 2 combined excised spleens on the day the mice were sacrificed. The fusion was performed 3–5 days later.

For ZYMV 1119 (poorly aphid transmissible, PAT) the immunization schedule from which the best McAbs were obtained was as follows:

- Day 1 – SC; 50 μg ; FCA
- Day 21 – IP; 50 μg ; FIA
- Day 28 – IV & IP (in vivo) or as an *in vitro* boost; 25 μg ; no adjuvant
- Day 31 – Fusion

For WMV-2, although the above schedule was used, no virus-specific clones were isolated. The immunization schedule which produced virus-specific McAbs was as follows:

- Day 1 – SC; 50 μg ; FCA
- Day 21 – IP; 50 μg ; FIA
- Day 83 – IP; 50 μg ; FIA
- Day 147 – IP; 50 μg ; FIA
- Day 167 – IV & IP; 25 μg ; no adjuvant
- Day 170 – Fusion

For ZYMV 1119 AT, both whole virus (ZYMV/AT) and capsid protein (ZYCP/AT) were used in parallel immunizations and fusion protocol. The immunization protocol which produced virus-specific McAbs was as follows:

- Day 1 – SC; 50 μg ; FCA
- Day 30 – IP; 50 μg ; FIA
- Day 37 – IV & IP (in vivo) or *in vitro* boost, 25 μg ; no adjuvant
- Day 40 – Fusion

In vivo fusion protocol:

The mouse was sacrificed, and the spleen was excised and gently mashed through an 80-mesh sieve, and placed in 5 mls of serum-free regular growth medium (DMEM). The SP2/0 myeloma cell line, obtained from the Hybridoma lab, U. of Florida, used as the fusion partner, was maintained in a logarithmic growth phase in DMEM-F, supplemented with 10% fetal calf serum, 1 mM L-glutamine, 1% penicillin/streptomycin, and 1 μ g/ml fungizone. Both spleen cells and SP2/0 cells were washed twice by centrifugation at 2,000 rpm for 7 min in serum free DMEM, and were combined at a ratio of 1:2 SP2/0:spleen cells, respectively. They were centrifuged together one final time and were resuspended in 50% 1500 MW PEG in 75 mM Hepes over a period of 15 sec. The cell suspension was swirled in a 37C water bath for 30 sec., incubated 90 sec. at 37C. One ml of serum-free DMEM was then added over 30 sec, swirled 30 sec, followed by the addition of 10 ml DMEM over 30 sec and incubation for 5 min in a 37C water bath. The solution was then centrifuged and the pellet resuspended in HAT selective media containing 20% fetal calf serum to give a final concentration of 5×10^5 cells/ml. These samples were plated out at 100 μ l/well in 96-well plates, and placed in a 37C incubator with an atmosphere of 6% CO₂ and 90% relative humidity. Cultures were fed with 50 μ l/well of HT medium 7 days after fusion and then individually as colonies developed. They were handled individually and screened for antibody production as they developed. Hybridoma cells were maintained on HT medium.

In vitro fusion protocol:

On the same day that mice for the *in vivo* fusion received an IV & IP injection, two mice, either previously immunized or not, were sacrificed, their spleens combined, and placed into a culture medium containing DMEM with 5mM 2-ME, 50 μ g pure virus, and 20 μ g/ml of the adjuvant N-acetylmuramyl-L-alanyl-D-isoglutamine (Calbiochem, Inc.), and 25 μ g of the immunogen. After three days in culture, the fusion was performed as previously outlined.

Screening Procedures:

All cultures were initially screened by both indirect and modified indirect ELISA (Halk, 1986; Van Regenmortel and Burckard, 1980) using purified virus preparations extracts from healthy plant tissues, and extracts from tissues infected with the respective virus. The modified indirect ELISA consisted of the following steps:

- (1) Rabbit IgG to the respective virus, at 1/1000 dilution in coating buffer, pH 9.6, was placed in ELISA wells at 75 or 100 μ g/well. Plates were incubated at 4C overnight or at 37C for 1 hr. This was followed by 3 washes with phosphate buffer saline containing Tween-20 PBS-T.
- (2) Antigen was added at 50 or 75 μ l/well, diluted in PBS-T at 1/1000 dilution (for pure virus) or 1/10 (for plant tissues). The tissues were triturated in a mortar and pestle, and expressed through Miracloth (Calbiochem, Inc.). Plates were incubated at 37C, 1 hr, followed by three washes.
- (3) Tissue culture supernate was added, 50 μ l/well, incubated at 37C, 1 hr. This step was also followed by three washes.
- (4) The enzyme conjugate, goat anti-mouse IgG and IgM alkaline phosphatase was added at 1/1000 dilution each in conjugate buffer (PBS-T with 2% PVP-40 and 0.2% ovalbumin) at 50 μ l/well. After incubation at 37C for 1 hr plates were washed three times.
- (5) The substrate, p-nitrophenylphosphate (PNPP) was added at 1 mg/ml using 50 μ l/well. Plates were read with a microplate autoreader EL309 (Bio-Tek Instr.) as reactions developed.

The indirect ELISA consisted of the same steps as above, omitting the first step and with the antigen dilutions made in coating buffer.

Cloning, Isotyping, and Ascites Production:

The most promising cell lines were selected for cloning, increased in large volume, and

used for ascites production. Each culture was cloned two to three times by limiting dilution. Cultures were serially diluted to a point where a single 96-well containing 30–50 viable cells could be identified. That well (approx. 100 μ l) was diluted in 10 mls of HT & plated out at 100 μ l well. Single cell colonies were selected and retested for virus specificity and similarity to the primary hybridoma, and increased. Isotyping was done by using a Zymed Monoclonal antibody ID kit using subclass-specific antisera for mouse IgG1, IgG2a, IgG2b, IgG3, IgM and kappa and lambda light chains. Ascites were produced by injecting at least 1×10^6 cells into Balb/C mice which had been primed 10–14 days earlier with pristane (0.5 ml, IP). Ascites were collected from five days to three weeks later using an 18-gauge needle, and clarified using Sure-Sep II serum separators (Organon Teknika).

Preparation of monoclonal antibodies to PRSV-W and its amorphous inclusion protein.

To prepare monoclonal antibody to capsid protein, hybridoma cells were obtained using preimmunized mice (Wiegers et al. 1986) and an *in vitro* technique modified from that described by Boss (1986) (S. Zam, personal communication). Balb/C mice were injected subcutaneously with 50 μ g of capsid protein in Freund's complete adjuvant, followed in two weeks by an intraperitoneal injection with 50 μ g of capsid protein in incomplete adjuvant. One month later and three days prior to the fusion the mouse spleen was removed and the spleen cells were suspended in an *in vitro* medium containing 25 μ g of antigen (capsid protein of PRSV-N, Adlerz isolate). The spleen cells were incubated in a CO₂ incubator at 37 C. On the day of the fusion the cells were centrifuged, put into 5 mls of DMEM, counted using a hemocytometer, and fused as described in the previous section. The cells were plated out (5×10^5 cells/ml, 100 μ l/well) into 96 well plates in HAT selection media [DMEM-F, containing 20% fetal calf serum, and 1 ml HAT/50 ml of media (hypoxanthine, aminopterin and thymidine, Cat. no. H 0262, Sigma Chemical Co., St. Louis, MO)], and placed in a CO₂ incubator. As growing colonies of hybridoma cells developed they were fed with HT media and their supernatants were tested for antibody production against purified antigen (similar to that used for immunization) in indirect ELISA. Supernatants of cells that were positive in this preliminary test were retested against extracts of 'Small Sugar' pumpkin (ssp) infected with PRSV-W, ZYMV, WMV-2, or zucchini yellow fleck virus (ZYFV) and healthy pumpkin. Hybridomas which were positive in this test were isotyped using a MonoAb EIA Kit (ZYMED LAB, Inc., San Francisco, CA) and frozen in liquid nitrogen. Primary hybridoma F3C-C10 (IgG1) was removed from liquid nitrogen after storage for 6 months, thawed, added to HT media, centrifuged, resuspended in HT media, and plated out. Supernatants from surviving cells were retested and the cells were cloned twice. Cells from a secondary clone of F3C-C10 were increased, counted, centrifuged, resuspended in sterile PBS and injected (1×10^7 cells) into pristane primed mice for ascites production.

With regard to PRSV-W amorphous inclusion protein, two of four *in vivo* fusions yielded antibody producing hybridomas. The Balb/C mouse, whose spleen was used in fusion F10, was injected four times with 50 μ g of the amorphous inclusion preparation. For the first injection, antigen was emulsified in Freund's complete adjuvant and injected SC. For the second injection, antigen was emulsified in incomplete adjuvant and was administered IP 16 days later, followed by a third IP injection 25 days later, and a fourth two months later. One month after the fourth injection and three days before the fusion the mouse was injected again both IV and IP with amorphous inclusion protein suspended in phosphate buffered saline (PBS). On the day of the fusion the mouse was euthanized and the spleen removed. Spleen cells were expressed the a 80 mesh sieve, counted, and fused with SP2/0 myeloma cells in a 2 to 1 ratio with the aid of polyethylene glycol. Cells were plated out in HAT media (5×10^5 cells/well) and placed in a CO₂ incubator for 7 days. The plates were fed with 50 μ l of HT media after 7 days. Those wells in which hybridomas were growing were fed as needed and 200 μ l of the supernatant was removed and tested in both plate-trapped and antibody-

trapped indirect ELISA against extracts from PRSV-W infected and healthy ssp. Hybridoma F10E-A5 was isotyped, cloned twice by limiting dilution and retested. Cells from clone F10E-A5-E5-H7 were increased, centrifuged, counted, resuspended in PBS, and injected (1×10^7 cells/mouse) into 4 pristane primed Balb/C mice for the production of ascites fluid. The fluid obtained was centrifuged, pooled, dispensed into vials at 1 ml per vial, and stored at -17°C .

Fusion F19 was done in a similar manner. The mouse used in this fusion was initially injected with $50\ \mu\text{g}$ of AI protein in complete adjuvant. Part of the injection was given SC and the remainder IP. A second IP injection in incomplete adjuvant was given one month later and a third 23 days later, part IP and part intramuscularly. A final IP injection was given 3 months later and the fusion was done 3 days after the last injection. The McAB obtained from this fusion was cloned twice and used for the production of ascites fluid as above.

Serological tests.

The SDS immunodiffusion tests were as described by Purcifull and Batchelor (1977) and Purcifull *et al.*, (1984c).

The enzyme-linked immunosorbent assays (ELISA) were indirect and modified indirect forms (Yeh and Gonsalves, 1984; Halk, 1986; Van Regenmortel and Burckard, 1980) as described above in the screening procedures for WMV-2 and ZYMV monoclonal antibodies. The procedures for western blotting were as reported (Burnett, 1981; Towbin *et al.*, 1979). For ELISA testing of isolates in Israel, the double antibody sandwich direct techniques of Clark and Adam (1977) was used. Immunosorbent electron microscopy was as reported by (Milne and Luisoni, (1977); Milne, (1984).

In vitro translation and analysis of cucurbit potyviral genomes.

The RNAs of selected ZYMV isolates were isolated from purified virus and translated *in vitro* in the rabbit reticulocyte lysate and wheat germ systems as previously described (Dougherty and Hiebert, 1980; deMejia *et al.*, 1985; Xiong *et al.*, 1988). Analysis of the *in vitro* translation products were conducted based on the immunoprecipitation patterns of components by various antisera to capsid and nonstructural proteins.

Host range, disease surveys and assessment of variability of cucurbit potyviruses.

Host range and host reactions of selected isolates in Israel and Florida were conducted by inoculating greenhouse grown plants mechanically or with viruliferous aphids. Field samples were collected and analyzed serologically. In Florida, samples that exhibited differences in ELISA reactions with monoclonal or polyclonal antibodies, that gave evidence of variation in SDS immunodiffusion with polyclonal antisera, that showed evidence of cytological differences by light microscopy (Christie and Edwardson, 1977, 1986; Purcifull *et al.*, 1984a), or that showed different symptoms in the indicator plant *Cucurbita pepo* 'Small Sugar Pumpkin' (ssp.) were maintained for additional studies.

Preparation and assessment of helper factor

In some of the initial experiments, a crude preparation of helper component was used (for experiments presented in Table 23). The helper component was obtained from zucchini leaves infected with ZYMV (a transmissible isolate of ZYMV) and from C-13 which was found to be non-aphid transmissible. Transmission tests were carried out as described elsewhere (Pirone, 1981; Raccach and Pirone, 1984); virus was mixed with a sample of the HC (20% sucrose was included for aphid phagostimulation). Groups of *M. persicae* previously fasted for 1 hr were allowed to acquire the virus-HC mixture for 10 min. The aphids were then confined in groups of 10 aphids per test plant on cucumbers. Inoculation access feeding was maintained overnight (16hr), then the aphids were killed with an aphicide and the symptoms were read

1–2 weeks thereafter.

For experiments presented in Table 24, helper component was purified as described in the previous section on the preparation of polyclonal antisera.

A procedure for immunoabsorption chromatography of ZYMV helper component (HC) was developed by D.W. Thornbury (Univ. Kentucky) at Bet Dagan in cooperation with B. Raccach. Protein A sepharose is equilibrated in buffer (.3 M KP04, pH 9.0); powder is suspended in approximately 10–20 volumes of buffer and allowed to swell for 30 to 60 minutes with occasional agitation. Columns are prepared in 3 or 5 cc syringes. A bed support is prepared by cutting a circle of either miracloth or GFA (glass fiber filter paper) paper to fit the inside diameter of the syringe. Connections to the UV monitor and fraction collector are made with thin PTFE or polyethylene tubing and 18 ga. needles. The syringe is filled with buffer and care is taken to make sure that there is no blockage of flow by air bubbles. The sepharose is suspended in buffer and poured into the column. The buffer is allowed to flow in order to pack the column to a height of one ml. Then the column is washed with buffer until the base line measured in the UV monitor (A 280) is flat and stable. About 100 to 200 μ l of antiserum are bound to each column by diluting it in the buffer above the top of the protein A gel. This allows maximum binding time without stopping the flow of the column. The column is washed with buffer until the baseline is again flat (until all of the unbound serum proteins have passed through the column). Buffer flow is stopped, and 50 to 75 μ l of HC in sucrose are loaded to the top of the column. The sample is allowed to enter the column; this takes approximately 5 to 7 drops, then the flow is stopped for 30 min to allow the binding of HC to IgG. The buffer flow is resumed to collect the unbound OD. In order to keep the dilution of activity to a minimum, the delay volume is predetermined (the number of drops between the monitor and the outlet to the collection tube). The peak is collected in two fractions; the first contains the maximum OD of the peak and the second the remainder of the peak. This maintains the concentration of activity of the first fraction at a 1/50 dilution or less. Because of this it is necessary that the HC extract used in the experiment be able to take this dilution and still have detectable activity.

For assay of samples, HC is tested against 3 sera: anti-ZYMV–HC, anti-PVY–HC, and normal serum. The volume of each fraction to be assayed is measured and the dilution factor calculated:

$$\frac{\text{volume of sample collected}}{\text{volume of sample applied}} = \text{dilution factor}$$

For each sample, untreated HC is diluted to the same dilution and tested for reduction of activity due to dilution rather than immunoabsorption. Recovery of bound protein and concentration for gels is accomplished as follows: 1. Elute protein and IgG from protein A with either .5M glycine–HCl, pH 3.5, or acetic acid/NaCl. 2. Concentrate all fractions for SDS PAGE by adding an equal volume of saturated ammonium sulfate and collecting the precipitate after incubation at 5 C overnight. Resuspend in the appropriate volume of SDS sample buffer, heat and store frozen until needed. 3. After all of the bound protein has been eluted from the columns, re-equilibrate the columns with buffer, check the pH of the eluant and stop flow. If the columns are to be stored for longer than a week, add .02% azide to the storage buffer.

Results and Discussion

Reactivity of polyclonal antisera.

Antisera to capsid proteins of PRSV–W (#1125), WMV–2 (1117 or 1132), and ZYMV (#1133) collected within c. 5–6 weeks after immunization were useful for detection and distinction of the viruses in SDS–immunodiffusion tests (Fig. 1). Later collections of the ZYMV

antiserum usually cross-reacted with WMV-2, but showed spur formation. Antiserum to PRSV-W capsid protein collected after re-immunization several months after the first immunization also cross-reacted with other potyviruses. These antisera also could be used to distinguish these potyviruses from other viruses in cucurbits, i.e., cucumber mosaic virus (Kuwite and Purcifull, 1982), squash mosaic virus (Purcifull *et al.*, 1981), and the unidentified virus (TV) from *Trichosanthes*, isolate #1860 (Fig. 1). Antisera to cylindrical inclusion (CI) proteins indicated that the CI proteins of the potyviruses are related, but different (Fig. 2).

Zucchini yellow fleck virus obtained from Italy (Vovlas *et al.*, 1979) was shown to be serologically related to PRSV-W, although they were not identical. Antisera to PRSV-W amorphous inclusion protein (1136) and PRSV-W cylindrical inclusion protein (1130) also cross-reacted with ZYFV-induced antigens. In immunofluorescence tests, the amorphous inclusions induced by ZYFV cross-reacted with antiserum to PRSV-W amorphous inclusion protein (Baker and Purcifull, 1987).

In reciprocal SDS-immunodiffusion tests, the antisera to capsid proteins of ZYMV (Florida isolate 1119) and the Reunion Island isolate of ZYMV, showed that these viruses are serologically related (H. Lecoq and D. Purcifull, unpublished data). However, the presence of spurs at the intersection of precipitin lines indicated that these isolates are not serologically identical.

In indirect ELISA, polyclonal antiserum 1136 to PRSV-W AIP reacted with all the PRSV isolates, WMV-M, ZYMV, and WMV-2. It did not react with SqMV, TV, CMV or healthy plant extracts. In SDS immunodiffusion tests, all the PRSV-W isolates and PRSV-P gave reactions of identity (no spur formation) with polyclonal antiserum 1136. Reactions of partial identity (spur formation) were seen with ZYMV, and WMV-2.

Using antiserum prepared to the ZYMV-W isolate, ELISA tests and serologically specific electron microscopy were conducted in Israel. These tests indicated that the MV and 13 isolates were serologically closely related to the ZYMV-W isolate, whereas the MC isolate was antigenically very different (Tables 2 and 3).

Experiments were conducted in Orlando to determine limits of antigen detection of ZYMV in polystyrene microtiter plates by ELISA and nitrocellulose membranes by dot blot immunobinding assay (DBIA) when the conjugate was alkaline phosphatase and a fluorogenic compound, 4-methylumbelliferyl phosphate. Fluorescence of 4-methylumbelliferyl phosphate was measured at wavelengths of excitation and emission of 360 and 450 nm, respectively, with a Sim Aminco spectrofluorometer, model SPF-500 C, whereas the optical density of *p*-nitrophenyl phosphate was measured at a wavelength of 405nm in a BioTek plate reader, Model 309. Detection levels of ZYMV in infected tissues were comparable with both substrates by indirect ELISA. Detection of log dilutions of ZYMV antigens resulted in coefficients of correlation of $r^2 = .99$ for both cases (Fig. 3). Detectable levels, however, did not exceed an antigen dilution of greater than 1/10,000 of the original dilution. In DBIA tests on nitrocellulose, NBT/BCIP was slightly more sensitive than 4-methylumbelliferyl in detecting low levels of ZYMV. Fluorescence was useful in DBIB on nitrocellulose because it clearly differentiated between a positive serological reaction and spots caused by plant pigments. This is important when the pigments can not be readily eliminated from the sample.

Characterization of Monoclonal Antibodies.

Reactivity of monoclonal antibodies to ZYMV and WMV-2. With the monoclonal antibody to poorly aphid transmissible ZYMV, no virus specific clones were obtained in the *in vivo* fusion, however the two *in vitro* fusions yielded 20%, or 12 virus-specific hybridomas (not previously immunized) and 14%, or 6 virus-specific hybridomas with an 80% fusion efficiency. Three of the most reactive primary hybridomas were selected for cloning, isotyping, and ascites production. All 20 clones produced were of the IgM isotype and reacted in the modified-indirect ELISA (antibody-trapped) only. With the exception of mild isolate 1994, all clones

were reactive with all Florida ZYMV isolates tested, but varied in reaction strength (Table 4). It is not known if the failure of the mild isolate 1994 to react is due to its low titer or to its inability to react with the monoclonal antibodies. All clones also reacted with the Lecoq isolates of ZYMV, with the exception that clone FG3-C5-C4 failed to react distinctly with the Reunion Island isolate. One clone, (BB2-A2-D4) which had a titer of 1024 for the tissue culture supernate in ELISA, was selected for ascites production. The final titer for ascites was 1/100,000 although 1/10,000 was the best working dilution for routine assays.

The efficiencies of the *in vivo* fusions were 37% and 50% for ZYMV/AT and ZYCP/AT, respectively. The fusion efficiency for the *in vitro* fusions was 40% & 52% for ZYMV/AT and ZYCP/AT. After final selections have been made, 8 stable primary hybridomas have been selected for ZYMV/AT and 11 for ZYCP/AT. To date, all of the selected hybridomas have been cloned at least once, with two selections injected for ascites production. Preliminary isotyping tests indicate that most of these may be of an IgG subclass. For each treatment, whether whole virus or capsid protein, there are clones which react in either modified-indirect, indirect ELISA, or in both. One primary hybridoma in particular reacted with all potyvirus isolates tested including ZYMV, WMV-2, PRSV-W, and ZYFV.

With WMV-2, the only fusion from which virus-specific hybridomas were obtained was the long-term (6 month) *in vivo* fusion. Of the 23 initial virus specific primary hybridomas, only two maintained their antibody production after successive subcultures. This loss of antibody-production is often characteristic of hybridomas in their early stages of growth (Halk, 1986). One strongly reactive hybridoma was chosen for ascites production and further screening. This clone, FD1-B5-B5, reacted in both modified-indirect and indirect ELISA, but with differing reactivity patterns. The modified-indirect ELISA was more useful for routine diagnosis of WMV-2 infections, and the indirect ELISA was better for distinguishing differences among WMV-2 isolates. Table 5 provides an example of the differences between the two types of ELISA tests with monoclonal antibody in comparison to polyclonal antisera both in ELISA and immunodiffusion. Overall, there was agreement between modified indirect ELISA (monoclonal) and immunodiffusion (polyclonal) with 65 of the 66 samples tested. One sample reacted very weakly in immunodiffusion with polyclonal antiserum but was negative in ELISA tests (FC-2359); the reason for this is not yet known. In contrast, there was agreement between indirect ELISA (monoclonal) and immunodiffusion (polyclonal) of only 37 of 49 samples.

Preliminary results from tests with the McAbs developed to ZYMV and WMV-2 during this study suggest that they may be useful for further characterization and distinction of the many potyvirus isolates which have been collected from the field and from cooperators. The McAbs to the poorly aphid transmissible strain of ZYMV react only in modified-indirect ELISA, indicating that they recognized only epitopes which are exposed on the surface of the intact virus particle. Correspondingly, these McAbs react specifically with only ZYMV isolates, and not with the other potyviruses tested. In contrast, the McAb to WMV-2 which reacts in both modified-indirect and indirect ELISA reacts with nearly all WMV-2 isolates in the former test and only with 76% of the isolates tested in the latter test. The reason for this differential reactivity within one clone has not been determined. One possibility is that certain virus isolates may be more susceptible to degradation than others, thus some isolates would react in one or both types of ELISA. It has been suggested that some ZYMV isolates are more susceptible to degradation than others (H. Lecoq, personal comm.). Another possible explanation is that the location of a particular epitope may differ between various isolates, thus being recognized in both tests, or in only one. A third possibility is that the virus titer being recognized is highly variable among different virus isolates, even under identical cultural conditions, thus giving variable reactivities in ELISA. This would not however, explain the complete nonreactivity of some isolates.

Characterization of the McAbs for both the whole virus and coat protein of the highly

aphid transmissible isolate of ZYMV are still in progress. Preliminary tests indicate that some clones may be reactive in one or both modified-indirect and indirect ELISA, therefore providing a wide range of reactive antibodies for strain characterization and epitope analysis.

Characteristics of monoclonal antibodies to PRSV-W. A monoclonal antibody to PRSV-W capsid protein (F3C-C10) was selected and used for ascites production. This monoclonal antibody was of the subclass IgG1 with kappa light chains. When tested in plate-trapped, indirect ELISA, it reacted with PRSV-W (Adlerz isolate) but not with sap from healthy pumpkin leaves. The F3C-C10 monoclonal antibody did not react with all PRSV-W isolates tested, however, as indicated in Table 6. These results are discussed more fully in the section on antigenic and biological variation.

McAB F3C-C10, which reacts with some PRSV isolates in both indirect and modified indirect ELISA, also reacts with seven other potyviruses in indirect ELISA. It reacts with bean yellow mosaic (204-1) (BYMV), bidens mottle (BiMV), peanut mottle, peanut stripe, potato virus Y (PVY), WMV-2, and soybean mosaic viruses. It did not react with cowpea aphid-borne mosaic, clover yellow vein (CyVV), pepper mottle (PeMV), tobacco vein mottling (TVMV), ZYMV, ZYFV, or wheat streak mosaic viruses. Polyclonal antiserum 1125 reacts in plate-trapped indirect ELISA with PeMV, PVY, CYVV, BYMV, WMV-2, tobacco vein mottling, bidens mottle, peanut stripe, peanut mottle, soybean mosaic, ZYMV, and ZYFV. It does not react with PeMV, cowpea aphid-borne mosaic, wheat streak mosaic, CMV, SqMV, TV or healthy tissue extracts. In SDS immunodiffusion tests, reactions of partial identity (spur formation) were seen with extracts from plants infected with WMV-2, TVMV, soybean mosaic, BiMV, PVY, peanut mottle, PeMV and bean yellow mosaic viruses. No reactions were seen with CMV, cowpea aphid borne, peanut stripe, ZYMV, CYVV, wheat streak mosaic, or healthy tissue.

Apparently PRSV-W shares common epitopes with at least 12 other potyviruses. Since McAB F3C-C10 recognizes only 7 of the 12 potyviruses recognized by polyclonal antiserum 1125 there are at least two different epitopes PRSV-W shares with other potyviruses.

Besides McAB F3C-C10, which was prepared to PRSV-W capsid protein, four McAB to the whole virus of PRSV-W (Adlerz isolate) have recently been produced (F21 was an *in vivo* fusion, F22 was an *in vitro* fusion). Their tissue culture supernatants have been tested in modified indirect ELISA with extracts of the same 14 PRSV-W isolates used to test McAB F3C-C10, and with PRSV-P, PRSV-T, WMV-M, ZYMV, ZYFV, WMV-2, SqMV, CMV, TV, and healthy pumpkin (Hssp). Between them these five McABs define at least six different epitopes of the Adlerz isolate of PRSV-W.

First, none of the McABs reacted with WMV-M indicating that the capsid protein of PRSV-W has at least 5 or 6 epitopes not present on the capsid protein of WMV-M. Two of the five McABs (F3C-C10 and F21D-E10) reacted with PRSV-T, confirming data using polyclonal antiserum that WMV-M is more distantly related to PRSV-W than PRSV-T. All five McAB reacted with PRSV-P, confirming its close relationship with PRSV-W.

McAB F21D-E10, and IgG1,k which only reacted in modified indirect ELISA, reacted with all the PRSV isolates tested so far. It appears to be an antibody to a conserved epitope specific to PRSV. This McAB did not react with WMV-2, WMV-M, ZYMV or ZYFV. It did not react with CMV, SqMV, TV, or Hssp.

McAB F21C-E4, an IgG2a,k which reacted in both indirect and modified indirect ELISA, reacted with all but two of the PRSV-W isolates (2207 and 2169). It also reacted with PRSV-P but it did not react with PRSV-T, WMV-M, WMV-2, ZYMV, ZYFV, SqMV, CMV, TV, or Hssp. Like McAB F3C-C10, this McAB appears to recognize an epitope that is not present on the capsid protein of all PRSV-W isolates. In SDS immunodiffusion tests using an early bleeding date of polyclonal antiserum 1142, which was made to the untreated virus of PRSV-W (Adlerz), the homologous antigen spurred over 2207 and 2169. The spur was weak but visible and the precipitin bands with these viruses were strong indicating a small

antigenic difference, possibly the same one detected with this McAB. No spurs were seen when 2207 and 2169 were tested against polyclonal antiserum 1125 to PRSV-W capsid protein. These two isolates were tested with the same bleeding date of 1125 that shows reactions of partial identity with the New York isolate and isolate 2201.

McAB F22A-B9, an IgM,k, reacted with all the PRSV-W isolates and PRSV-P. It did not react to PRSV-T, WMV-M, WMV-2, ZYMV, ZYFV, CMV, SqMV, TV, or Hssp. This McAB appears to recognize an epitope of PRSV-W and PRSV-P which is not present on the capsid protein of PRSV-T.

McAB F22A-C8, an IgM,k, reacted only with the homologous antigen in antibody trapped indirect ELISA. Preliminary work with the aphid transmissibility of the PRSV isolates used in this study has shown that the Adlerz isolate is very poorly aphid transmissible and that the capsid protein is responsible for this lack of transmissibility. All the other isolates were readily aphid transmissible. Perhaps this McAB is detecting an epitope on the capsid protein of PRSV-W (Adlerz) that is associated with this difference in aphid transmissibility.

Ascites fluid production to the four new McABs to the whole virus of PRSV-W is in progress. The ascites will be tested against the same PRSV isolates and the other squash viruses to confirm the data obtained with tissue culture fluid.

Reactivity of a monoclonal antibody to PRSV-W amorphous inclusion protein.

A single hybridoma (FIDE-A5) was found that tested positively with PRSV-W and negatively with healthy ssp. It reacted in plate-trapped and antibody-trapped indirect ELISA and was isotypized as an IgG1 with kappa light chains. While the original screening procedure showed that McAB F10E-A5 reacts with infected tissue and not healthy tissue in ELISA, Western blots (Fig. 4) confirmed this and showed that this monoclonal was reacting to a protein in infected tissue that migrated in electrophoresis at the approximate molecular weight of the amorphous inclusion protein (51K-56K daltons). The monoclonal antibody reacted like the polyclonal antiserum and there was no reaction to SP2/0 ascites or to preimmunized rabbit serum. There also was no reaction in the lane with infected tissue at molecular weights for the cylindrical inclusion (66K-75K daltons) or the capsid protein (30K-45K daltons). Polyclonal antiserum to these two proteins did react with the appropriate molecular weight bands (data not shown). This monoclonal antibody also reacted only to purified amorphous inclusion protein and not to purified cylindrical inclusion or capsid protein in an modified indirect ELISA where each protein was trapped by its homologous antiserum (Table 7).

In indirect ELISA, MCA F10E-A5 reacted with all of the PRSV isolates tested (Table 8). It also reacted with WMV-M, ZYMV (Table 9) and with WMV-2 (Table 10). It did not react with squash mosaic, cucumber mosaic, or healthy ssp. (Table 8). In addition, it reacted with soybean mosaic, peanut stripe, peanut mottle, and ZYFV. It did not react with extracts from plants infected with tobacco vein mottling (TVMV), tobacco etch (TEV), clover yellow vein (CYVV), (PeMV), bidens mottle (BiMV), wheat streak mosaic, TV or with extracts from healthy plants. Reactivity of this MCA with PVY, BYMV, and cowpea aphid-borne mosaic virus is not clear at this time as some tests showed a weak reaction and others showed a negative reaction.

Characterization of viruses by *in vitro* translation

Several zucchini yellow mosaic virus (ZYMV) isolates were compared by *in vitro* translation analyses. The isolates were distinguished on the basis of host reaction, aphid transmissibility, and coat protein serology. The main difference between ZYMV Reunion isolate (ZYMV-R) and the other ZYMV isolates was the apparent efficient processing of the helper factor protein (51kD protein) in the translation of the ZYMV-R in the rabbit reticulocyte lysate (RRL) system (Fig. 5). In the other ZYMV isolates and for all other potyviral translations to date with the RRL system, the helper factor appears as a polyprotein (in the case of ZYMV, as 88kD

protein) (Hiebert and Dougherty, 1988). The translation analysis of the ZYMV-R in the wheat germ system (Fig. 5) appeared similar to that found for other potyviral RNAs, i.e. the predominant product did not react with any of the available antisera. No translational product differences were detected between ZYMV E15 (aphid transmissible) versus ZYMVPAT (poorly aphid transmissible) and between ZYMV VC1119 versus ZYMV FC 1119B.

Since the helper factor appeared to be processed in the RRL translations of ZYMV-R, attempts were made to assay the translation products for helper factor activity. No activity was detected in limited experiments.

The RNA from the cucurbit virus 1860 from *Trichosanthes* (TV) (suspected to be a potexvirus) was translated in the RRL system. The predominant product was about 180kD and no product immunoprecipitable by coat protein antiserum was detected. This result is consistent with translations reported for most members of the potexvirus group (Dougherty and Hiebert, 1985).

Natural and Experimental hosts of cucurbit potyviruses.

Host range studies in Israel. Isolates were collected on the basis of differences in their symptom reaction on cucurbits. The isolates used in this study are designated by letters or numbers. The capital letter represents the source plant, the number or small letter is a serial marker given in the process of collection.

Two isolates derived from cucumbers (C-4 and C-13), two isolates derived from melons (MV and M-c) and one isolate derived from watermelon (W-1) were studied. C-4 and C-13 have shown symptoms typical of ZYMV. MV differed from the isolates collected from cucumbers and showed symptoms similar to WMV-2.

Of the five isolates of potyviruses tested, a series of host plants was found which allows a reliable distinctive reaction: C-13 and C-4 differ one from the other by their reaction on *Cucumis ficifolius* (CFC), on the other hand, neither infects peas, a characteristic which separates them from the other three potyviruses (Table 11). MV differs from M-c and W-1 by its reaction on *Cucumis africanus* (CAF). Finally, M-c is the only isolate which is symptomless on *Cucumis sativus* cv. Shimshon (CS) (the rest induce symptoms). The seven test plants used in Table 11 provide a reliable tool to separate these 5 potyviruses. Table 12 shows the reaction of several cucurbit hosts to the virus isolates tested. Immune reaction was recorded to C-13 among a long list of non-cucurbit plant species, as shown in Table 13.

Serological detection of viruses in cucurbits in Florida. In 1987 and 1988, antisera to PRSV-W, WMV-2, ZYMV, cucumber mosaic virus (CMV), squash mosaic virus (SMV), and the virus from *Trichosanthes* (TV) were used to detect the corresponding viruses in field samples by SDS-immunodiffusion tests. The samples represented different crops, seasons, and locations in the state. These samples were either collected in disease surveys or were submitted for diagnosis, and they were collected either in commercial fields or in experimental plots that had become infected naturally. Of 549 samples, the percentages of samples that assayed positively were as follows: PRSV-W (44.6%), WMV-2 (37.1%), ZYMV (9.2%), CMV (2.3%), SMV (0%) and TV (4.0%).

The occurrences of the viruses varied in some instances depending on the crop (Table 14), on location in the state (Tables 15 and 16), and on the time of year that the samples were collected (Table 16). Samples that reacted with two different antisera were common and were determined to be the result of plants that were doubly infected. These samples included those that reacted with antisera to PRSV-W and WMV-2, WMV-2 and ZYMV, PRSV-W and ZYMV, WMV-2 and CMV, and PRSV-W and TV. Triply infected plants also were detected, including those that reacted with antisera to PRSV-W, ZYMV, and WMV-2, or one that reacted with antisera to PRSV-W, ZYMV, and CMV. All samples were tabulated individually for their reactivity to each virus antiserum. In order to preserve virus isolates for further study,

leaves from infected plants (either original material or material resulting from greenhouse inoculations) were dried under vacuum and stored over desiccant at 4°C. At least 214 isolates were preserved in this manner.

The number of samples that did not react with any of the six antisera constituted 17.3% of the total (Table 2). These samples either were not infected with any of the six viruses or the virus titer was not high enough to be detected in this test system. Most of these samples either had no characteristic virus-induced mosaics or distortion symptoms, or had symptoms that may have been due to other causes. There is a possibility that some samples were infected with viruses that would not have been detectable with the antisera used.

PRSV-W is one of the most prevalent viruses in the state (Tables 14,15,16). Although it occurs in South Florida in the spring and fall (Table 16), its occurrence in north and central Florida seems primarily to be in the fall crops. PRSV-W was detected in all the major cucurbit crops and in *Coccinea grandis*, but not in *Trichosanthes dioica* (Table 14). The pattern of occurrence of PRSV-W in Florida was similar to that reported previously by Adlerz and others (Adlerz, 1969, 1981; Adlerz, *et al.*, 1983).

WMV-2 is an important pathogen of squash, watermelon, and cantaloupe, and it also was detected in cucumber (Table 14). WMV-2 was detected in both north and central Florida in the spring and fall (Table 16). It was not detected at all in the south Florida counties of Collier, Dade, and Palm Beach. Adlerz and others obtained similar results with WMV-2 (Adlerz, 1969; Adlerz *et al.*, 1983). Interestingly, Webb and Scott (1965) noted that only WMV-2 was detected in Collier County in 1962-63, but only PRSV-W (WMV-1) was detected there in 1964-65.

In 1981, ZYMV was found for the first time in Florida (Adlerz *et al.*, 1983; Purcifull *et al.*, 1984c), and its severe effects on several cucurbits were the cause for concern because ZYMV-resistant cultivars were not available. In the 1987-1988 seasons, ZYMV was detected in various parts of the state, but most of the ZYMV-infected cucurbits were detected in central Florida (Tables 15 and 16). Although ZYMV was detected in fewer samples than either PRSV-W or WMV-2, it is still regarded as an important pathogen in cucurbit production in Florida because of its severity, its occurrence in 9.2% of the cucurbits tested, and its distribution. The sources of inoculum of ZYMV in the state are not well known, although natural infection of *Melothria pendula* in central Florida has been reported (Adlerz *et al.*, 1983).

The TV was detected only in *Trichosanthes dioica* and *Coccinea grandis* (Table 14) in Dade Co. (Table 15), the only location where these two species were collected for assay. Although squash is susceptible to the TV, this virus was not detected in any of the more than 70 squash samples collected in Dade Co. The pathological effects of TV on *Coccinea* and *Trichosanthes* are unknown at this time. Furthermore, the characterization of TV is insufficient to permit its classification with respect to other plant viruses. However, preliminary tests indicate it is filamentous and that it may be serologically related to papaya mosaic virus, a known potexvirus (Purcifull, unpublished data).

Cucumber mosaic virus has been known to occur in cucurbitaceous crops in Florida for many years (Anderson, 1952), and it has been detected in commelinaceous weeds in northern, central and southern Florida (Adlerz, 1981; Baker and Zettler, 1988). In our study, CMV was detected only in a small number of the cucurbit samples collected in north and central Florida (Table 16). It was not detected in any of the 186 cucurbit samples collected in south Florida.

Squash mosaic virus (SMV) was not detected in this study, although it has been reported previously in cantaloupe in Florida (Anderson, 1954; Purcifull *et al.*, 1981). The number of cantaloupe samples assayed during the 1987-1988 seasons, however, was not large (36 samples). Anderson reported more than 30 years ago that SMV (then called muskmelon mosaic virus) was uncommon in central Florida. Two cantaloupe isolates previously collected in Florida reacted serologically (Purcifull *et al.*, 1981) with the same SMV antiserum used in the present study.

Mosaic virus diseases continue to constitute significant problems in the production of cucurbits in Florida. This is particularly true of the diseases caused by the aphid-borne potyviruses, PRSV-W, WMV-2, and ZYMV. The *Trichosanthes* virus (TV) represents a pathogen that apparently has not previously been described in Florida (Alfieri, *et al.*, 1984). There are no indications at this time that TV occurs to any significant extent, if at all, in major cucurbit crops in Florida. However, this situation might change in the future. Because of the large number and diverse types of viruses that infect cucurbits world-wide (Lovisolo, 1980), it seems likely that more viruses will be found in Florida as more surveys are conducted, antisera to additional viruses are used, and more diverse detection techniques are employed. More extensive application of serological and other indexing techniques could also help to obtain a better understanding of the epidemiology of the viruses affecting cucurbits in this state.

Seed transmissibility of cucurbit potyviruses in squash. Transmission of WMV-1, WMV-2 and ZYMV from 'President' or 'Multipik' seeds collected from a potyvirus-infected squash field in central Florida was not obtained (0/1300 and 0/850, respectively for the two cultivars). However, when a portion of President seed was used to mechanically inoculate Small Sugar receptor plants, transmission resulted from normal seed (3/1300 or 0.23%) and "paperthin" aborted seeds (15/450 or 1.76%), whereas no transmission occurred with normal or paperthin seeds (0/850 and 0/350, respectively) of 'Multipik', a resistant cultivar (Table 17). ZYMV was detected most frequently, but WMV-2 and PRSV-W were also detected (Table 17).

Biological and Antigenic Variation of Cucurbit Potyviruses.

Biological variation of ZYMV. In Leesburg and Orlando, biological and antigenic characteristics of 4 ZYMV isolates, 86-1, 86-4, 86-8, and 86-16, which were obtained from aborted seed coats of ZYMV-infected zucchini squash var. 'President' were compared with 2 known ZYMV isolates, 81-25 and FC1119. No serological differences were detected between these 6 isolates by SDS immunodiffusion tests with polyclonal antiserum to ZYMV (1133). The isolates were mechanically inoculated in a host range (Table 18). Blister size and color in small sugar pumpkin differentiated the isolates into 2 groups: 1) light green leaves with raised dark green blisters which included 81-25, 86-1, and 86-4; and 2) predominantly green leaves with smaller areas or streaks of chlorosis and small blisters or distortions which included 86-8, 86-16, and FC1119. Lack of systemic infection in Alaska pea and no evidence of local lesions in 633-3 cantaloupe and vein clearing symptoms in *Melothria* indicated that the isolates tested were not coinfecting with PRSV-W or WMV-2. Minor isolate differences were observed in *Luffa*. None of the isolates caused symptoms in *Ranunculus sardous*, but each isolate infected this species, which is a known host of ZYMV (Lisa and Lecoq, 1984).

Laboratory vector tests with 3 aphid species also were conducted on the 6 isolates. The most efficient vector of ZYMV was *Myzus persicae* followed by *Aphis gossypii* and *Aphis citricola* (Table 19). Transmission efficiency of FC1119 was low. This isolate was originally highly aphid transmissible but had been maintained by mechanical transmission for several years. An isolate of FC1119 obtained by aphid transmission was found to be much more efficiently vectored and was called FC1119AT. High transmission rates were obtained from 3 of the aborted seed ZYMV isolates, 86-1, 86-4, and 86-8, whereas transmissibility of 86-16 was lower. Transmission of 81-25 was intermediate by comparison. None of the isolates tested were nontransmissible. Differences in vector transmission were also observed between biotypes of *A. gossypii*. A Leesburg colony of *A. gossypii* (AG-L) was much more efficient in transmission than an Orlando biotype (AG-O) of the aphid. These data support both the observation that transmissibility is a property of the virus and an attribute of the vector.

In Gainesville, selected pathotypes of Florida isolates of ZYMV differing in geographical origin, symptomatology in pumpkin, and aphid transmissibility were defined based on their

behavior in inoculated melon lines (Table 20) (H. Lecoq and D. Purcifull, unpublished data). Three isolates (FC 1994, 2000, and 2050) belong to the pathotype F (i.e., inducing a wilting reaction on melon lines possessing the Fn gene) while three isolates (FC 1119GH, 1119AT, and 2154) belong to the pathotype NF (i.e., inducing mosaic but not wilting). The proportion of the pathotypes seems to be similar in Florida and Europe (H. Lecoq, personal comm.), suggesting a similar pattern of variation for this virus under different ecological situations. None of the 6 isolates systemically infected melon plants homozygous for the dominant resistance gene Zym. Therefore, these 6 isolates belong to pathotype O. This suggests the potential of this gene for breeding programs in Florida. However, inoculation of FC 1119 AT to F1 hybrid plants of susceptible and resistant parents led to a low percentage of systemic infections. Subcultures (FC-1119-A, FC-1119B) obtained from 2 of these plants consistently induced mosaic and leaf deformation in the resistant parent, indicating that they overcame the resistance conferred by Zym. Therefore, they belong to pathotype 2 (Table 20). No serological differences have been found between these isolates in SDS immunodiffusion using antisera to the coat protein or cylindrical inclusions. The FC-1994 was a unique, mild isolate from a nonsymptomatic squash plant in the field. The 1994 showed mild mosaic symptoms in pumpkin. Mild isolates might have some potential as a cross-protecting agent.

Biological and antigenic variation of PRSV-W isolates from Florida. The Adlerz isolate of PRSV-W (PRSV-WA) and various other isolates of PRSV-W and PRSV isolates were used in this study. An isolate from Greece and an isolate from Jordan came from G. Martelli, an isolate from California came from R. Webb, an isolate from New York came from R. Providenti, one isolate was obtained from the American Type Culture Collection (ATCC PV-23), and the remaining isolates were collected in Florida. The Florida isolates, 2052 (in zucchini) and 2040 (in yellow squash) were collected in Jan. 1987 from Dade County and isolates 2030 (in zucchini) was collected from Collier County at the same time. In addition, an isolate of PRSV-T from Guadeloupe from L. Quiot-Douine (Quiot-Douine et al. 1986), a PRSV-P (Hawaii) from D. Gonsalves, and an isolate of watermelon mosaic virus from Morocco (WMV-M) originally from B. Lockhart, were used in this study. Isolates 2201 and 2207 (both in pumpkin) and 2169 (in watermelon) were collected in Alachua County in the fall of 1987. Isolates 1870 (in yellow squash) and 1637 (in zucchini) were collected in Dade County in April 1986 and January 1985, respectively.

Because of the results found with the aforementioned isolates in both serological (see below) and biological tests (above), 38 cucurbit samples collected in the spring of 1988 for a continuing survey of cucurbit viruses (Tables 14,15,16) were also tested for reactivity to the monoclonal antibody F3C-C10. Samples 2246 to 2276 were collected from Dade County and samples 2277 to 2283 were collected from Collier County. Eight of these samples (2246,2247,2252,2262,2263,2266,2267, and 2269) were inoculated to 4 melon lines after initial inoculation to ssp.

In SDS-immunodiffusion, isolate 2201 From Florida, a New York isolate, and isolates PRSV-T and WMV-M (Morocco) were spurred over by the homologous antigen, PRSV-WA, with polyclonal antiserum 1125. The precipitin lines of the other isolates coalesced with the homologous antigen. All the PRSV-W isolates and PRSV-P tested reacted with lines of identity with antiserum 1130 to cylindrical inclusion protein. WMV-M and PRSV-T reacted with partial fusion to PRSV-WA with this antiserum. With antiserum 1136 to amorphous inclusion protein, the precipitin lines of PRSV-P and all the PRSV-W isolates coalesced with PRSV-WA. The precipitin lines of PRSV-T and WMV-M were spurred over by the homologous antigen. Thirty-six additional field samples from Florida were infected with PRSV based on positive reactions to the polyclonal antiserum 1125 in SDS immunodiffusion.

In plate-trapped indirect ELISA, isolates 2040, 2052, 2030, WMV-M, and healthy ssp did not react with the monoclonal antibody F3C-C10. The remaining isolates reacted positively

with this antibody. None of the antigens reacted with the ascites of SP2/0 cells. All isolates reacted with polyclonal antiserum 1125 but not with normal serum. Healthy ssp did not react with either serum. Table 6 gives the absorbance (405 nm) readings one hour after the addition of substrate. Readings are an average of two wells.

In the ELISA test of the inoculated papayas, only the positive controls (PRSV-WA and PRSV-P in ssp) and the papaya inoculated with PRSV-P reacted with antiserum 1125. None reacted with normal serum.

While all 36 of the isolates 2246 to 2283 reacted positively to polyclonal antiserum 1125 in plate-trapped indirect ELISA, only thirty of the PRSV-infected samples reacted positively to the monoclonal antibody F3C-C10. Eight samples (2246, 2247, 2252, 2262, 2263, 2266, 2267, 2269), or approximately 20% of the infected samples, did not react with the monoclonal antibody.

A summary of the types of symptoms caused on the melon lines inoculated is contained in Table 21. In general, all isolates gave resistant reactions on VAC, Edisto 47, and PI 161375 while the rest of the isolates, including 2030, gave susceptible reactions on these melon lines. In addition, 2040, PRSV-T, and WMV-M gave resistant reactions on VED, while the remaining isolates, including 2052 and 2030, gave susceptible reactions. Isolates 2246, 2247, 2252, 2262, 2263, 2266, 2267 and 2269 also gave a susceptible reaction (mosaic on Supermarket and a resistant reaction, local lesions, on WRN-29). Like isolate 2040, isolates 2246, 2247, 2252, and 2266 gave resistant reactions on both Edisto 47 and VED. The reaction of 2246 on Edisto 47 consisted of local lesions (LL) and systemic necrotic spots (SNS) while the reactions of 2040 and 2247 consisted of LL, SNS, and top necrosis (TN) followed by rapid plant death. Isolates 2252 and 2266 gave LL only on Edisto 47 and SNS and/or TN on VED. Like isolate 2052, isolates 2262, 2267, and 2269 gave a resistant reaction on Edisto 47 and a susceptible reaction on VED. However, unlike isolate 2052, which caused TN and quick plant death on Edisto 47, these isolates caused a severe stunting with necrosis and death of the lower leaves and SNS on the newer leaves. These plants did not die and had one or two very small chlorotic leaves with SNS at the end of the test. Like isolate 2030, 2263 gave a susceptible reaction on both Edisto 47 and VED.

Two important aspects of plant virus control strategy are virus identification and the breeding of virus resistant or tolerant cultivars. In order to do these efficiently an understanding of the serological and biological variability of the virus in question is essential. This study has shown the existence of both serological and biological variability heretofore not known in Florida isolates of PRSV-W.

In SDS immunodiffusion, antiserum 1125 detected epitope variation in the capsid protein of isolates 2201 which was not correlated with any of the other differences found among Florida isolates. This isolate and the New York isolate, which was also spurred over by PRSV-WA, reacted essentially the same as PRSV-WA on the melon lines except perhaps for a subjective observation of slightly less severe symptoms. This result indicates the need to use other types of tests (i.e. biological) and to include antigens and antisera of other cucurbit viruses or antisera to other PRSV-W proteins in serological tests in order to differentiate epitope variation in the capsid protein of PRSV-W from antiserum cross-reactivity of PRSV-W antisera with other potyviruses.

Another variation in the capsid protein of PRSV-W isolates was recognized by the lack of reactivity of approximately 20% of the isolates tested with the monoclonal antibody F3C-C10. Although the correlation was not 100%, this serological difference appeared to be related to the differential biological reactions seen in the melon line inoculation study. Only two of the 12 isolates which did not react with the monoclonal antibody gave susceptible reactions in Edisto 47 and VED. Only one isolate gave resistant reactions in these cultivars and also reacted with the monoclonal antibody.

It is interesting that three types of biological reactions were found among the Florida

isolates that did not react to monoclonal antibody F3C-C10. However, an epitope of a protein is estimated to consist of 5-7 amino acid residues and it has been shown that a single amino acid change can modify the reactivity of a monoclonal antibody (Moudallal et al., 1982). Perhaps the differential biological reactions seen with isolates like 2040, 2052, and 2030 represent different amino acid changes in the epitope recognized by F3C-C10. Some changes only modify the reactivity to the monoclonal antibody (i.e. 2030). Other changes modify the reactivity of the monoclonal antibody and the virus-plant interaction in certain melon lines (i.e. 2040 and 2052). Knorr and Dawson (1988) found that a single nucleotide change in the 3' region of the capsid protein gene of TMV, leading to a single amino acid change, could convert a susceptible reaction in *Nicotiana sylvestris* (N' gene) to a resistant reaction. It is possible that something similar occurs in potyviruses.

Knorr and Dawson (1988) also reported that one TMV mutant which induced a resistant reaction in *N. sylvestris* (N' gene) did not contain the nucleotide change found in the other mutants. We also found one isolate, PRSV-T, which reacted to our monoclonal but still gave a resistant reaction on VAC, Edisto 47, PI 161375, and VED. This suggests that either the apparent correlation of biological and serological differences found in this study are coincidental or, as suggested by Knorr and Dawson (1988), more than one gene product or sequence alteration can affect virus-host compatibility.

In any case, it is certain that some isolates of PRSV-W in Florida induce resistant type reactions in cultivars not known to possess resistance genes to standard PRSV-W isolates (i.e. FC-2040 in VED, VAC, and Edisto 47). This indicates that these isolates have different levels of virulence in melons and that these cultivars have "weak" resistance genes.

Aphid transmissibility of cucurbit potyviruses and partial characterization of helper component. Studies with isolates from Israel. The transmissibility of four isolates was tested using two major well known vector species of these viruses, namely the green peach aphid, *Myzus persicae*, and the melon aphid, *Aphis gossypii*. The results are shown in Table 22; two isolates were found to be non-transmissible by aphids, namely MV and C-13. The other two isolates were transmitted at varying efficiency. *M. persicae* was more efficient in transmission of C-4 than of the M-c isolate while *A. gossypii* has shown the reverse, being more efficient in transmission of M-c than C-4.

The helper component (HC) used in the stages of the study was a crude one. The helper component was obtained from zucchini leaves infected with ZYMV-T (a transmissible isolate of ZYMV) and from C-13 which was found to be non-aphid transmissible. Inoculation access feeding was maintained overnight (16 hr), then the aphids were killed with an aphicide and the symptoms were read 1-2 weeks thereafter. The results are presented in Table 23. No aphid-transmission was obtained with aphids which acquired C-13 in mixture with ZYMV-TC HC, with Potato virus Y-HC or with C-13 HC. On the other hand, HC from the non-transmissible C-13 was capable of mediating the transmission of purified ZYMV-T (Table 23). This finding implies that the lack of transmission of C-13 is related to its coat protein rather than to the lack of helper protection.

Helper components prepared from transmissible and non-transmissible isolates were all active in assisting transmission of purified transmissible isolates. The helper components were purified as follows:

Extraction in 0.1 M phosphate buffer pH 9.0.

Sedimentation of virus by ultracentrifugation ca at 100,000g.

Collection and evaluation of the helper capacity of the supernatant (SN).

Concentration of helper from the SN using 30% ammonium sulphate.

This purification resulted in a good quality helper component which ensured transmission when used at 10% of the transmission volume with 100 µg/ml virus (Table 24).

Four HC preps were prepared for two distinct transmissible isolates (ZYMV-W and

ZYMV-MC) and two non-transmissible isolates (ZYMV-13 and ZYMV-MV). Homologous and heterologous mixtures of HCs and viruses were tested by aphid transmission experiments. All four HC were found able to assist transmission of purified virus preparations from transmissible virus isolates.

The use of ammonium sulfate was imperative for concentrating the ZYMV-HC that was dissociated in SDS and purified by preparative polyacrylamide gel electrophoresis prior to use as immunogen. The antiserum to SDS-dissociated protein extracted from polyacrylamide gels of ZYMV-HC preparations was tested for neutralization of HC activity using immunoabsorption chromatography. Protein A agarose-IgG columns were prepared using anti-ZYMV-HC and anti-PVY-HC antiserum. It had been previously found that the PVY-HC antiserum reacted with ZYMV-HC and neutralized its activity in immunoprecipitation experiments. A highly active preparation of ZYMV-HC was passed through each column and allowed to react with the respective antiserum. The unbound protein from the ZYMV-HC preparations was then tested for activity in the aphid bioassay. Activity of the column extracts was compared with untreated ZYMV-HC diluted to a volume equivalent to that of the eluants from each column. Both antisera completely absorbed the HC activity while the diluted control showed 40% transmission of ZYMV. This provides the first evidence that the protein extracted from polyacrylamide gels to which the antiserum was made is related to ZYMV-HC activity. Since the return of Dr. Thornbury to Kentucky immunoabsorption experiments using the ZYMV-HC antiserum against PVY-HC has shown that it also absorbs PVY-HC activity. These reciprocal experiments provide strong evidence we identified the protein responsible for ZYMV-HC activity.

Studies with isolates from Florida. The propensity of aphids to transmit WMV-2 and ZYMV was tested using field collected alate aphids prior to virus symptom appearance in the field. Aphids were collected by suction and yellow water-pan traps and were immediately placed on either a WMV-2 - or ZYMV-infected 'Small Sugar' pumpkin plant for a 2-min virus acquisition period and then placed on Dixie squash receptor plants. *Aphis middletonii* and *Myzus persicae* were the most frequently trapped aphids and had the highest percent transmission of WMV-2 and ZYMV of 25% and 28%, respectively (Table 25). Several other aphids appeared to have high transmission efficiency (i.e. *Lipaphis erysimi*) but were collected in low numbers.

Studies were initiated to determine if serological titer of the virus or the helper component was related to vector transmissibility. Polyclonal antisera to capsid protein (CP) of ZYMV FC1119 (1133-1) and to amorphous inclusion (AI) protein to PRSV-W (AI 1042) were used in these tests. The tests were conducted by indirect ELISA using goat anti-rabbit conjugated to alkaline phosphatase. The antigen was diluted 1/10 in 20 mM sodium carbonate buffer, pH 9.6. Preliminary data have indicated no correlation of observed vector transmission of measured levels of CP and AI (Table 26). However, the poorly transmitted FC1119 had a much lower titer of CP and AI than the other isolates tested. In comparison, an isolate identified as FC1119AT, which was obtained by aphid transmission (AT) of isolate FC1119, was found to be more readily transmissible than the parent isolate. It had higher serological titer of CP and AI in infected donor plants than the original isolate. The lower antigen level of FC1119 may have been an indication that the virus titer of this isolate was below the threshold necessary to obtain higher rates of vector transmission.

Preliminary tests were conducted in Florida with helper component purified and concentrated by high-speed centrifugation and ammonium sulfate precipitation. Helper components from four ZYMV isolates were tested by *in vitro* transmission tests with *Myzus persicae*. Isolates 81-25 and 86-8 had active HC with purified ZYMV from isolate 81-25; whereas no transmission resulted with HC from FC1119 and FC1119AT with the same virus source. The highest activity of HC from ZYMV 81-25 (based on the number of infected

plants) was from clarified sap following centrifugation at 150,000 g (r_{av}) for 2 hr (Table 27). The HC was successfully concentrated by density gradient centrifugation but lost its biological activity. However, it should be noted that the purification of active ZYMV–HC is often affected by unknown factors, and that different results can be obtained with the same purification procedure. Similar results have been observed with PVY–HC. The activity of ZYMV–HC observed in the high–speed supernatant of plant sap was typical of those reported by Sako and Ogata (1981) and Lecoq and Pitrat (1985) who found that crude sap yielded higher aphid transmission rates than other HC extraction methods.

Epidemiological Studies with Cucurbit Potyviruses.

Epidemiology of ZYMV in Israel. In the spring of 1986, a 1/2 acre of commercial zucchini field was used for the spring season epidemiological study. The plot was sown on March 15, and aphid collection started upon germination. Ten yellow water pans and 10 green tile traps were exposed in the field. Aphids were collected twice a week. As symptoms appeared, the infection was monitored, and the peaks of infection were related to the peaks of aphid species trapped. The results showed that the peaks of alates of *Aphis citricola* and *A. gossypii* were at a time when increase of virus disease was minor. Exponential increase of virus disease was only noted after the peaks of flight subsided. Large populations of aphids, particularly *A. citricola*, were present in late March and the beginning of April. However, at that time and almost three weeks later than the last occurrence of aphids in flight, fast increase in virus spread occurred. Also, the spread in the field as seen from the layout of infection maps shows a clear case of spread from plant to plant. Indeed, prior to the fast increase in spread, there was an intensive colonization of melon aphids (*A. gossypii*).

Additional epidemiological studies were carried out in two locations which were characteristic for commercial cultivation of melons, watermelons and squash: Gilgal in the Jordan Valley and Masuot Itzhak in Lachish. In Gilgal, virus spread was recorded in the Fall season of 1986 (Fig. 6). Aphid activity data were taken both in the fall of 1986 and in the spring of 1987 (Table 28 and Figures 7 and 8). In Masuot Itschak, the monitoring of virus spread started in May and ended in August 1987.

In Gilgal a 1/2 acre plot of Spaghetti squash (cv. Gugeti) which was sown in the Experimental Station in mid September 1986, germinated in early October and virus spread data was taken since then. The only virus ascertained in the field was ZYMV. Distribution of infection was recorded once or twice a week. Follow up of infection was maintained for 4 weeks when an early winter and low temperature caused a premature termination of the experiment. Aphid incidence was recorded in 10 yellow traps distributed in the Spaghetti squash plot. When the number of aphids landing in traps was very large, a sample was taken from each of the 10 yellow water traps. Lacking electrical power supply in the region, we were unable to use suction traps. Instead, aphids were collected alive and tested for their infectivity using the method suggested by and coordinated with the late Dr. Warren Adlerz (Adlerz, 1987). As seen in Fig. 6, a notable secondary spread of ZYMV was discerned. Aphid composition of species in the fall of 1986 and in the spring of 1987 is presented in Table 28. The build up of aphid population during the fall of 1986 and the spring of 1987 is presented in Fig. 7 and 8. None of the aphids trapped in the water pans were found capable of transmitting ZYMV to test plants. The composition of species greatly differed between the fall and the spring: In the fall three species were found most frequently: *Penatlonia nigronervosa* (which derives from adjacent banana plantations), *Acyrtosiphon pisum* (which derive from leguminosae) and *A. brunae* (from unknown source). The potential of these species in transmission of ZYMV is now under simulated field conditions. A completely different composition of species was ascertained in the spring, where three familiar species comprised more than 50% of the total aphid population: *M. persicae*, *A. gossypii* and *R. maidis*. All three species are well known to be capable of transmitting the virus. It is to be noted that few

aphids only were recorded landing during the brief periods in the yellow pans. Of those, none were found to transmit the virus to test plants.

In Masuot Itzhak two melon (cv. Galia) commercial plots were used for monitoring virus spread. The first was shown two weeks earlier than the second. In the first plot, there was an initial infection of virus when monitoring started, while in the younger plot no virus was seen in the first inspection. The plot was in the vicinity of cotton, which was infested with the melon aphid at the stage of melon germination. Here too, secondary spread of virus was recorded, particularly before the increase in summer heat.

The Arava valley, on the way to Eilat, is a location where cucurbits and particularly melons can be grown even in the winter, during the mild so called "eternal summer". The place is distant from most major growing regions in Israel. A question arose as for the source of aphids in this location. For this purpose, Dr. Eastop from the British Museum, came to Israel to collaborate on the search of aphids in the desert. The outcome of this work is summarized in the attached manuscript (Eastop and Raccach, 1988, Appendix 1). The abstract of this report is as follows: A survey of aphid species which colonize or reproduce on wild and cultivated plants in the Arava Valley of Israel, was carried out at the end of winter 1986. Thirty-three aphid species were collected on wild plants and 23 in green tile or yellow traps exposed in cultivated plots. Aphids were recorded from 78 plant species in 22 botanical families. *Myzus persicae*, *Aphis craccivora*, *Brachyunguis harmalae*, *Acyrtosiphon gossypii* and *Aphis gossypii*, were present on plants in 19 botanical families. *M. persicae* was the most frequently trapped aphid species in both the green tile or yellow (Moericke) traps. The possible origin of the aphid species found in the desert was considered in view of the ecology and geography of the region. The role of aphid species found in the Arava Valley, in the epidemiology of potato virus Y (PVY) and zucchini yellow mosaic virus (ZYMV), is discussed.

A major enigma for ZYMV, was our inability to discover the possible winter host for the virus in Israel. A few wild cucurbits were found here and there in the vicinity of the crop. However, of those tested so far for presence of virus, none were found to be infected. This is particularly important since most cultivated cucurbits are very sensitive to cold temperatures and the virus should disappear along after the host with the first frost. However, virus was found in winter in umbelliferous wild plants in the Arava. The virus recovered readily infected cucurbits and is related to ZYMV.

Epidemiological studies of cucurbit potyviruses in Florida. Natural infectivity of aphids collected by suction and yellow water-pan traps from a WMV-2- and ZYMV-infected zucchini field was tested in the spring. Aphids most frequently collected were *A. middletonii*, followed by *L. erysimi*, *A. citricola*, and *M. persicae* (Table 36). *A. middletonii* and *A. citricola* had the highest percent transmission of WMV-2 and ZYMV. *L. erysimi* also transmitted these viruses, and based on these data, may be a noteworthy vector for field spread.

Control of Cucurbit Potyviruses

Efforts to develop control measures for ZYMV were undertaken in Israel. These efforts involved the development of a combination mineral oil-pyrethroid spray combination and a study of the inheritance of resistance in an interspecific cross of *Cucurbita*.

Control of ZYMV in Israel with a combination of pyrethroid and oil sprays. In 1986, an experimental spray program on spaghetti squash in Israel was conducted as follows. Five treatments were each carried out in two replicates. Each replicate consisted of 3 seedling beds. On each bed, there was one row, plants were sown in pairs, ca. 50 cm away from each other. Spaghetti squash cv. Gugetti was sown in the plot on September 20, 1986. The first spray was conducted on October 1st, when plants had an average of 2 true leaves. The first two sprays were conducted using a ULV carry-on sprayer. Later, we used a high volume and pressure sprayer which also used a stream of air to disperse the droplets. The

sprayer nozzles were stainless steel TX 5., and the pressure was at least 200 psi. Counts were taken from one row which was the central row in each treatment as soon as first symptoms appeared in the field. Results were analyzed using the CHI SQUARE TEST (Table 29). As seen in Table 29, the strongest decrease in the rate of infection was recorded in the combination of oil and pyrethroid. Records took into account virus infection, as recorded by visual inspection. The effect on the plants in general and on the yield in particular was even more remarkable, as the plants found infected in the treatments, showed symptoms only at the very end of the branches, with little if any effect on the quality or quantity of the fruit. An additional experiment was conducted in Talme Yaffe (Lachish), but the rate of infection in the control was so low that it did not allow valid comparison.

In 1988, an additional experiment was conducted to evaluate the effects of sprays on control of ZYMV in Israel. The experimental plot was at Zafaria village which specialized in growing melons. The plot was sown on April 5, 1988, and was the first which was sown in the open (being sensitive to low temperatures, melons are sown under a plastic cover in earlier sowings). Because of technical problems, sprayings started only on the second week after germination, and were maintained at a one week interval up to June 6, 1988. The first three sprays were done with a high volume and pressure sprayer; those applied later were done with a sleeve sprayer combined with a strong air stream. The plot was part of a large commercial plot, which was surrounded with melon plots, and there were citrus groves in the region. The section designated for the experiment was of a 1/2 acre (2,000 m²) in size. In order to reduce margin effects, the treatments were applied to a large section composed of 500 m² each. The direction was such that the control plot was most extreme to the east, farthest from the prevailing wind, to expose it to the least effect of incoming alates. The experimental plot included 7 rows for each treatment, and data were taken from the three central rows, to avoid possible spray drift effects. In mid May, a severe heat wave prevailed for three days followed by unusually high temperature for more than a week. In general temperatures in the summer of 1988 were higher than the multiannual temperature records for the regions. There were no trappings of aphids in green or yellow traps after that, nor were aphids found on vegetation. Green tile and yellow traps were exposed in each of the treatments, and aphids were collected 2–3 times a week, and kept for identification. The distance and lack of electricity did not allow live trapping of aphids. Infection was monitored once or twice a week, first by visual inspection, and later, after discovering the presence of some CMV in the plot, all the plants were also tested by ELISA for ZYMV. Absence of other cucurbit potyviruses was ascertained by periodical SSEM, Ouchterlony and ELISA tests. At the end of the experiments, the total fruit from 50 plants were picked, segregated to marketable (reticulated nice looking melons) and nonmarketable (virus infected, distorted and green and uneven fruits), and the yield from each treatment was counted and weighed. The composition of aphid species in the Zafaria village conforms with that described by Raccach et al. (1985) for Bet-Dagan (which is 8 km South to Zafaria). Two species, *A. citricola* and *A. gossypii*, comprised more than 70% of the total population trapped in green tiles and almost 98% of the total population trapped in yellow traps (Table 30). Of the two aphid species, *A. citricola* appeared at the beginning of the season, but its peak did not coincide and came earlier to the logarithmic increase of the infection in plot (Figs. 9–12). On the other hand, *A. gossypii* and other *Aphis* spp. (combined) were present at the logarithmic increase stage of the infection (Figs. 9–12). For two weeks, after the heat wave, and up to the picking stage, no aphids were trapped either in yellow traps or in green tile traps. Also, no aphids were recorded on the vegetation. However, an apparent spread of the virus continued, which no vector could be accounted for. In order to explain this discrepancy, we studied the possibility that the extraordinarily high temperatures affected the symptom expression which resulted in gradual appearance of symptoms, looking as if virus was still spreading. The following experiment was planned to explain the phenomenon: Cucumber test

plants (*Cucumis sativum* L. cv. *Beth-Alfa*) at the cotyledon stage were used for testing the effect of temperature on symptom expression. Virus was acquired by *M. persicae* from ZYMV infected zucchini. Each of the test plants was inoculated with 3 viruliferous aphids. Then, the test plants were divided into two groups, one kept in growth chambers at 35 to 40 C and another at 22 to 25 C. First symptoms appeared after a week among the plants which were kept at 25 C and eventually all the test plants became infected. Those kept at 35 C showed a delayed development, and only after about two weeks, certain atypical symptoms started to appear in 2/3 of the plants. Symptoms became apparent after about three weeks in almost all the test plants, but even at this stage, the symptoms were retarded and atypical. Lacking data on live trapping, we cannot relate the infection to particular species. However, judging from the time of appearance of major aphid species, it is logical to assume that *A. citricola* could only contribute to primary infection and some minor secondary spread. The presence of *A. gossypii* was related to the build up of virus in the field, and related well with clustered spread of the virus.

The effect of the treatment was seen in the rate of infection in the field, as well as in the time when the infection occurred. This resulted in an increased number of marketable fruit and in higher yield (see Fig. 13 and Tables 31 and 32).

As seen in Fig. 13 and in Tables 31 and 32, it is apparent that both oil and the combination of oil and Pyrethroid, yielded more fruit than for the unprotected plants. Following these and the previous findings, these treatments are now the recommended practice in many melon fields in Israel.

An experiment was conducted on the effect of pyrethroids on aphid behavior, using alate aphids. Leaves of cucumbers were dipped in water, in Telstar 0.15% and in Maverick 0.15% solution. The leaves were allowed to dry for about two hours before being tested. Groups of 10–12 alates of *A. gossypii* and *M. persicae* were put on the leaves, and the number of alates taking off from the leaves were monitored for a period of 10 min. There was a significant difference between Telstar treated leaves and untreated even 2 min after aphid exposure. The proportion of alates departing from the leaves is shown in Fig. 14. Although ZYMV as a typical nonpersistent virus that can be inoculated in a matter of seconds, the likelihood of infection increases with time during the first minutes. The fact that an effect was noticed immediately after the first minutes explains the difference between the efficiency of pyrethroids and other insecticides for prevention of virus transmission (Pirone et al. 1988).

Inheritance of resistance to ZYMV in Cucurbita.

The inheritance of resistance to zucchini yellow mosaic virus (ZYMV) in the interspecific cross *Cucurbita maxima* X *C. ecuadorensis* was studied in generations P1, P2, F1, F2, BCPI and BCP2 (Tables 33,34,35). The continuous variation presented in these generations indicates a quantitative inheritance. The resistance of *C. ecuadorensis* is partially dominant over the sensitivity of *C. maxima*. A few genes with major effects along with genes with small effects appear to control ZYMV resistance. Both the dominant and the additive contribution to resistance were found significant. Their interaction was significant, too. However, the major contribution was due to the additive rather than the dominance effects. Narrow sense heritability is 0.91. There is a positive correlation ($r=0.76$) between virus titer as measured by ELISA and intensity of visual symptoms in the plant.

Literature Cited

Adlerz. 1979 Proc. Fla. State Hort. Soc. 82:161.

- Adlerz. 1976. J. Econ. Entomol. 69:495.
- Adlerz. 1981. in "Pests, Pathogens, and Vegetation". (J.M. Thresh, ed.). Pitman, London.
- Adlerz. 1987. J. Econ. Entomol. 80:87.
- Adlerz, Purcifull, Simone, & Hiebert. 1983. Proc. Fla. State Hort. Soc. 96:72.
- Alfieri, Wehlburg, Landon, & Kimbrough. 1984. Fla. Dept. Agric. & Consumer Services Bulletin
No. 11. 389 p.
- Anderson. 1952. Plant Dis. 36:377.
- Anderson. 1954. Phytopathology 44:198.
- Baker & Purcifull. 1987. Phytopathology 77:1722 (Abstr.).
- Baker & Zettler. 1988. Plant Dis. 72:513.
- Boss. 1986. Methods in Enzymol. 121:27.
- Burnette. 1981. Anal. Biochem. 112:195.
- Christie & Edwardson. 1977. Fla. Agric. Exp. Stn. Monogr. Ser. IX. 150 pp.
- Christie & Edwardson. 1986. Plant Dis. 70:273.
- Clark & Adams. 1977. J. Gen. Virol. 34:475.
- Davis & Yilmaz. 1984. Phytopathology 74:848 (Abstr.).
- DeMejia, Hiebert, & Purcifull. 1985. Virology 142:24.
- DeMejia, Hiebert, Purcifull, Thornbury, & Pirone. 1985. Virology 142:34.
- Dougherty & Hiebert. 1980. Virology 101:466.
- Dougherty & Hiebert. 1985. In "Molecular Plant Virology" (J.W. Davis, ed.), Vol. II. CRC
Press, Boca Raton, Fla.
- Eastop & Racciah. 1988. Phytoparasitica 16:23.
- Edwardson. 1974. Fla. Agric. Exp. Stn. Monogr. Ser. IV. 398 pp.
- Gonsalves & Ishii. 1980. Phytopathology 70:1028.
- Greber. 1978. Austr. J. Agric. Res. 29:1235.

- Grogan, Hall, & Kimble. 1959. *Phytopathology* 49:366.
- Halk. 1986. *Methods in Enzymol.* 118:766.
- Hein. 1977. *Phytopathol. Zeit.* 89:221.
- Hiebert & Dougherty. 1988. In "The Filamentous Plant Viruses" (R.G. Milne, ed.) Plenum Press, N.Y.
- Hiebert, Purcifull & Christie. 1984. In *Methods in Plant Virology*, Vol. 8 (K. Maramorosch & H. Koprowski, eds.). Academic Press, Orlando, FL.
- Hollings & Brunt. 1981. CMI/AAB Descriptions of Plant Viruses No. 245.
- Knorr & Dawson. 1988. *Proc. Nat. Acad. Sci. (U.S.A.)* 85:170.
- Kuwite & Purcifull. 1982. *Plant Dis.* 66:1071.
- Lecoq, Lisa & Dellavalle. 1983. *Plant Dis.* 67:824.
- Lecoq & Pitrat. 1985. *Phytopathology* 75:890.
- Lecoq, Pitrat, & Clement. 1981. *Agronomie* 1:827.
- Lesemann, Makkouk, Koenig, & Samman. 1983. *Phytopathol. Z.* 108:304.
- Lisa, Boccardo, d'Agostino, Dellavalle & d'Aquilio. 1981. *Phytopathology* 71:667.
- Lisa & Lecoq. 1984. CMI/AAB Descriptions of Plant Viruses No. 282.
- Lovisolo. 1980. *Acta Horticulture* 88:33.
- Makkouk & Lesemann. 1980. *Plant Dis.* 64:799.
- Milne. 1984. In *Methods in Virology*, Vol. 7. (K. Maramorosch & H. Koprowski eds.). Academic Press, Orlando, FL.
- Milne & Grogan. 1969. *Phytopathology* 59:809.
- Milne & Luisoni. 1977. In *Methods in Virology*, Vol. 6. (K. Maramorosch & H. Koprowski, eds.). Academic Press, Orlando, FL.
- Moghal & Francki. 1976. *Virology* 73:350.
- Moudallal, Briand, & Van Regenmortel. 1982. *EMBO J.* 1:1005.

- Nameth, Dodds, & Paulus. 1984. *Phytopathology* 74:794 (Abstr.).
- Pirone. 1981. *Phytopathology* 71:922.
- Pirone, Raccach, & Madden. 1988. *Plant Dis.* 72:350.
- Providenti, Gonsalves & Humaydan, 1984. *Plant Dis.* 68:443.
- Providenti & Schroeder. 1970. *Plant Dis. Repr.* 54:744.
- Purcifull & Batchelor. 1977. *Fla. Agric. Exp. Sta. Bulletin* No. 788.
- Purcifull, Christie & Lima. 1981. *Phytopathology* 71:1221.
- Purcifull & Hiebert. 1979. *Phytopathology* 69:112.
- Purcifull, Edwardson, Hiebert & Gonsalves. 1984a. CMI/AAB Description of Plant Viruses No. 292.
- Purcifull, Hiebert & Edwardson. 1984b. CMI/AAB Description of Plant Viruses No. 293.
- Purcifull, Adlerz, Simone, Hiebert & Christie. 1984c. *Plant Dis.* 68:230.
- Quiot-Douine, Purcifull, Hiebert, & DeMejia. 1986. *Phytopathology* 76:346-351.
- Raccach. 1983. In "Plant Virus Epidemiology", (R.T. Plumb & J.M. Thresh, eds.). Blackwell Scientific Publications, Oxford, England.
- Raccach, Gal-On, & Eastop. 1985. *Ann. Appl. Biol.* 106:451-460.
- Raccach & Pirone. 1984. *Phytopathology* 74:305.
- Russo, Martelli, Vovlas, & Ragozzino. 1979. *Phytopathol. Medit.* 18:94.
- Sako & Ogata. 1981. *Virology* 112:762.
- Towbin, Staehlin, & Gordon. 1979. *Proc. Nat. Acad. Sci. (U.S.A.)* 76:4350.
- Van Regenmortel, Brandes, & Berko. 1962. *Phytopathol. Z.* 45:205.
- Van Regenmortel & Burckard. 1980. *Virology* 106:327.
- Webb, Bohn & Scott. 1965. *Plant Dis. Rept.* 49:532.
- Webb & Scott. 1965. *Phytopathology* 55:895.
- Wieggers, Uhlig & Dernick. 1986. *J. Gen. Virol.* 67:2053.

Xiong, Hiebert & Purcifull. 1988. *Phytopathology* 78:1128.

Yeh. 1984. Ph.D. Diss; Cornell Univ., Ithaca, N.Y. 98 pp.

Yeh and Gonsalves. 1984. *Phytopathology* 74:1273.

Table 1. Polyclonal antisera prepared in rabbits to cucurbit potyviruses and their potyvirus-specified proteins, and to a virus isolated from *Trichosanthes*.

Virus*	Immunogen	Designation and Source of Isolate	Rabbit #s
PRSV-W	virus	Adlerz; Florida	1142
PRSV-W	capsid protein	Adlerz; Florida	1125
PRSV-W	CI protein	Adlerz; Florida	1130
PRSV-W	AI protein	Adlerz; Florida	1136
ZYMV	virus	W; Israel	6/12/85
ZYMV	virus	1119; Florida	1160
ZYMV	capsid protein	1119; Florida	1133
ZYMV	capsid protein†	Lecoq; Reunion Is.	1141
ZYMV	CI protein	1119; Florida	1135
ZYMV	helper protein	W; Israel	32/1/87
WMV-2	virus	1656; Florida	1134
WMV-2	capsid protein	1656; Florida	1117, 1132
WMV-2	CI protein	1656; Florida	1115
MC	virus	MC; Israel	22/5/88
TV	virus	1860; Florida	1124
TV	SDS-treated virus	1860; Florida	1129

*PRSV-W = papaya ringspot virus type W; ZYMV = zucchini yellow mosaic virus; WMV-2 = watermelon mosaic virus; MC = isolate from Israel which is serologically related to the Florida isolate of WMV-2; TV = *Trichosanthes* virus (possible potexvirus).

Table 2. Absorbance (a405) of the ELISA reaction of ZYMV-W antiserum against four cucurbit potyviruses.

IgG (μ g/ml)	Conjugate Dilution	Virus Isolate			
		MC	MV	13	ZY-W
1	1:500	0.0176	0.9458	1.4362	1.4307
1	1:1000	0	0.4712	0.7792	0.7782
2	1:1000	0.0247	0.9217	1.35	1.4742

The results represent means of 8 replicates for each treatment.

Table 3. Mean number of virus particles per electron microscope field at the same magnification as an indication of the trapping capacity of antiserum to ZYMV for four cucurbit potyviruses.

Virus Isolate	Mean Number : SD
MC	0.75 : 1.69
MV	58.95 : 9.32
ZY	40.6 : 6.70
13	92.6 : 10.78

The means represent fields (4 fields in each of 5 grids).

A dilution of 1:2000 of the ZYMV antiserum in 0.1M phosphate buffer pH 7, was used for coating the grid, prior to trapping the virus particles.

Table 4. Reaction of ZYMV Monoclonal Antibodies with Cucurbit Potyviruses in Modified Indirect ELISA.

Monoclonal antibody	<u>Purcifull Florida ZYMV Isolates</u>						Healthy Pumpkin
	1119GH	1119AT	Mild 1994	2000	2050	2154	
BB2-A2-D4 ^a	0.903	1.472	0.016	1.476	0.902	1.562	0.054
FG3-C5-C4 ^b	0.681	1.40	0.033	1.607	0.593	1.679	0.054
AD4-H11-E7 ^b	0.178	0.521	0.044	0.418	0.096	0.543	0.053

Monoclonal antibody	<u>Yokomi and Adlerz Florida ZYMV Isolates</u>					Healthy Pumpkin
	81-25	86-4	86-1	86-8	86-16	
BB2-A2-D4 ^b	1.899	1.349	1.726	1.372	1.683	0.016
FG3-C5-C4 ^b	1.323	0.606	1.038	0.757	1.172	0.011
AD4-H11-E7 ^b	0.879	0.229	0.773	0.317	0.756	0.025

Monoclonal antibody	<u>Lecoq French ZYMV Isolates Other Cucurbit Potyviruses</u>						Healthy Pumpkin
	<u>Reunion</u>	<u>PAT^d</u>	<u>E-15</u>	<u>PRSV-W^c</u>	<u>WMV-2</u>	<u>ZYFV^e</u>	
BB2-A2-D4 ^a	0.276	1.191	0.923	0.050	0.044	0.018	0.011
FG3-C5-C4 ^b	0.042	1.069	0.892	0.034	0.031	0.023	0.001
AD4-H11-E7 ^b	0.705	1.540	1.588	0.061	0.075	0.065	0.021

^aAscites diluted 1/1000 in TBS^bTissue culture supernate diluted 1/10 in TBS^cValues are an average of two readings at 405 nm^dPoorly aphid transmissible isolate^eSame primary hybridoma used for these plant samples; different subclone

Table 5. Reaction of a Monoclonal Antibody^a for WMV-2 with Field Collected Cucurbit Viruses in ELISA and Immunodiffusion Tests.

<u>Sample No.</u>	Modified- Indirect ELISA	Indirect ELISA	Polyclonal- Indirect ELISA ^b	Immuno- diffusion; polyclonal ^c
1656	0.459 ^c	0.249	0.239 ^d	+
2289	0.261	0.037	0.565	+
2291	0.127	0.069	0.430	+
2292	0.015	0.018	0.042	-
2293	0.198	0.021	0.437	+
2294	0.200	0.029	0.352	+
2295	0.210	0.879	0.410	+ (strong)
2296	0.282	0.819	0.307	+ (strong)
Healthy pumpkin	0.008	0.021	0.027	-
ZYMV 1119	0.016	0.039	0.053	-
Squash mosaic virus	0.002	0.005	0.048	-

^aWMV-2 clone FD1-B5-B5 ascites at 1/1000 dilution in (TRIS buffered saline).

^bSerum from rabbit no. 868 diluted 1/1000 in (TBS).

^cValues are an average of two readings read at 405 nm.

^dValues are based on single samples read at 405 nm.

^eBased on reaction with rabbit polyclonal antiserum no. 1134.

Table 6. Detection of antigenic variation in PRSV isolates and WMV-M by the monoclonal antibody F3C-C10^b.

<u>Virus isolate</u>	<u>F3C-C10</u>	<u>1125^c</u>
WMV-1	1.16	.502
FC-1637	.927	.585
FC-2169	1.025	.512
FC-2040 ^a	.008	.428
FC-2201	.748	.390
FC-1870	.663	.383
FC-2030 ^a	.010	.474
FC-2052 ^a	.004	.395
PRSV-P	.191	.221
Greece	.830	.409
Jordan	.989	.434
New York	.640	.409
ATCC	.578	.355
Calif.	.606	.362
Tigre	.439	.250
WMV-M ^a	.008	.200
H.ssp	.004	.004

^aIsolates which did not react with monoclonal antibody F3C-C10.

^bPlate-trapped indirect ELISA readings (A-405) 1 hr after addition of substrate.

^cNo. 1125 represents polyclonal antiserum to PRSV-W capsid protein.

Values represent the average readings of two wells. Any value over 0.10 was considered a positive reaction. Values lower than .02 were considered nonreactive. Absorbancy values for SP2/0 ascites ranged from .002-.018. Absorbancy values for normal serum ranged from .001-.009.

Table 7. Reactivity of MCA F10E-A5 to AIP in antibody-trapped (modified indirect) ELISA to three purified proteins of PRSV-W*.

PROTEIN	ABSORBANCE READING (A-405)**
Amorphous inclusion	0.298
Cylindrical inclusion	0.003
Capsid	0.007

*AIP = Amorphous inclusion protein of papaya ringspot virus type W.

**Each protein was trapped by its homologous polyclonal antiserum. Readings are an average of two wells.

Table 8. Serological reactivity of MCA F10E-A5 to AIP with various papaya ringspot virus isolates infecting pumpkin (ssp)*.

VIRUS	ABSORBANCE (A-405 nm)**	
	PLATE-TRAPPED	ANTIBODY-TRAPPED
PRSV-W		
WMV-1A	0.152	0.378
ATCC	0.433	0.396
Florida (9)	0.253-0.455	0.327-0.536
Jordan	0.261	0.414
Greece	0.289	0.474
New York	0.296	0.431
California	0.317	0.394
PRSV-P	0.290	0.426
PRSV-T	0.415	0.407
Hssp	0.007	0.013

*AIP = Amorphous inclusion protein of papaya ringspot virus type W.

**Readings were taken at 2 hr for the plate-trapped test, at 1 hr for the antibody-trapped test, and are an average of two wells

Table 9. Serological reactivity of MCA F10E-A5 to AIP and polyclonal AIP antiserum in plate-trapped ELISA with several other viruses that infect pumpkin.

VIRUS	ABSORBANCE (A-405 nm)**	
	MCA	POLYCLONAL
PRSV-W	0.152	0.273
WMV-M	0.335	0.364
ZYMV	0.138	0.329
WMV-2	0.036	0.155
CMV	0.017	0.028
SqMV	0.011	0.033
Hssp	0.007	0.021

*AIP= Amorphous inclusion protein of papaya ringspot virus type W.

**Readings were taken at 2 hr for the MCA, at 30 min for the polyclonal antiserum, and are an average of two wells.

Table 10. Reactivity of MCA F10E-A5 and polyclonal AIP antiserum in plate-trapped ELISA with extracts of *Nicotiana benthamiana* (Nb) infected with watermelon mosaic virus-2 and cucumber mosaic virus*

ANTIGEN	ABSORBANCE READINGS (A-405)**	
	MCA	POLYCLONAL
WMV-2	0.415	0.485
CMV	0.020	0.016
HNb	0.017	0.021

*AIP=Amorphous inclusion protein of papaya ringspot virus type W.

**Readings were taken at 1 hr for the MCA, at 10 min for the polyclonal antiserum, and are an average of two wells.

Table 11. The reaction of seven hosts to five cucurbit potyviruses

Virus	Host Reaction		
	Immune	Symptomless	Sensitive
C-13	Pea	ZR, CFG	CS, CBA, CAF, CFC
C-4	Pea	ZR, CFG, CFC	CS, CBA, CAF
W-1		ZR, CFG, CAF	Pea, CS, CBA
MV		ZR, CFG	Pea, CS, CBA, CAF
M-c	CFG, CFC	CS, ZR, CAF	Pea, CBA

a. The potyviruses enclosed herein are designated on the basis of the source host. Only after further serological and biological characterization will they be classified into conventional groups of cucurbit potyviruses (such as ZYMV, WMV2 or PRSV-W).

ZR = Cucumis sativus cultivar showing symptomless reaction to ZYMV.

CBA = C. sativus cv. Bet-Alpha, a sensitive cultivar to most potyviruses tested so far.

CS = C. sativus cv. Shimshon.

CAF = Cucumis africanus.

CFG = Cucumis figarej.

CFC = Cucumis ficifolius.

Table 12. Host range for several differential cucurbits

Test Plant	C-13	C-4	W-1	MV	M-c
Royal Slice	SL	SL	SL	SL	SL
ZYMV-R					
<u>C. metuliflora</u>	S	S	S	S	S
Havat Shalem	S	S	S	S	S
<u>C. figarei</u>	SL	SL	SL	SL	SL
<u>C. longipees</u>	S	S	S	S	S
<u>C. proliphetarum</u>	S	S	S	S	S
<u>C. zeypheri</u>	S	S	S	S	S
<u>C. ficifolius</u>	S	SL			I
<u>C. ariguria</u>	S				S
<u>C. hookeri</u>	S				S
<u>C. dipsaceus</u>	S				S

(Reaction was recorded as sensitive, causing visible symptoms designated as "S"), symptomless (where virus could be ascertained in the plant but no or almost no symptoms were discernible, designated as "SL"), and immune (where no virus was present in the host, designated as "I").

Table 13. List of species found immune to the Israel isolates of potyviruses tested in the present study.

<u>Plant species</u>	<u>Botanical family</u>
<i>Calotropis procera</i>	Asclepiadaceae
<i>Solanum vilosum</i>	Solanaceae
<i>Nicotiana glauca</i>	Solanaceae
<i>Hyoscyamus desertorum</i>	Solanaceae
<i>Ipomea</i> spp.	Convolvulaceae
<i>Malva nicaensis</i>	Malvaceae
<i>Bidens tripartita</i>	Compositae
<i>Malvella Sherardiana</i>	Malvaceae
<i>Solanum melogena</i>	Solanaceae
<i>Lavatera cretica</i>	Malvaceae
<i>Vigna sinensis</i>	Leguminosae
<i>Sonchus oleraceous</i>	Compositae
<i>Datura metel</i>	Solanaceae
<i>Brassica pekinensis</i>	Cruciferae
<i>Nicotiana tabacum</i>	Solanaceae
<i>Nicotiana clevelandii</i>	Solanaceae
<i>Capsicum annuum</i> (cv. laor)	Solanaceae
<i>Nicotiana glutinosa</i>	Solanaceae
<i>Solanum lycopersicum</i>	Solanaceae
<i>Hibiscus esculentum</i>	Malvaceae

Table 14. Results of serological assays for six viruses in various cucurbit crops in Florida, 1987–1988.

Crop ^y	# of samples assayed	# of samples positive for ^z						N
		PRSV-W	WMV-2	ZYMV	CMV	SMV	TV	
Squash	305	177	89	27	6	0	0	39
Watermelon	130	34	89	15	0	0	0	17
Cucumber	39	19	2	7	3	0	0	16
Cantaloupe	36	4	23	2	3	0	0	8
Coccinea	15	11	0	0	0	0	14	1
Trichosanthes	11	0	0	0	0	0	8	3
Miscellaneous	13	0	1	0	1	0	0	11
Totals	549	245	204	51	13	0	22	95

^zData indicate the number of samples that were identified based on their reactivity to papaya ringspot virus type W (PRSV), watermelon mosaic virus-2 (WMV-2), zucchini yellow mosaic virus (ZYMV), cucumber mosaic virus (CMV), squash mosaic virus (SMV), and the Trichosanthes virus (TV).

N=the number of samples that did not react with any of the six antisera.

^ySquash primarily consisted of yellow summer squash or zucchini, but includes all samples of the genus Cucurbita. The miscellaneous category includes unidentified Cucumis samples, (one reacted with WMV-2 antiserum and one with cucumber mosaic virus antiserum), and 4 samples each of Luffa and Momordica (all of which failed to react with any of the antisera).

Table 15. Results of serological assays for six viruses in cucurbits in Florida in 1987–1988, tabulated by county.

County	Region of state	# of samples assayed	Assay Results					
			# of samples positive for :					
			PRSV-W	WMV-2	ZYMV	CMV	SMV	TV
Alachua	NF	79	27	30	1	4	0	0
Citrus	CF	8	0	8	0	0	0	0
Collier	SF	32	28	0	2	0	0	0
Dade	SF	104	73	0	1	0	0	22
Dixie	NF	8	0	8	0	0	0	0
Duval	NF	3	0	0	0	2	0	0
Gadsden	NF	13	2	9	0	0	0	0
Gilchrist	NF	8	1	8	0	0	0	0
Hillsborough	CF	5	4	0	5	1	0	0
Jackson	NF	38	3	17	0	0	0	0
Jefferson	NF	33	15	7	0	0	0	0
Lafayette	NF	1	0	1	0	0	0	0
Lake	CF	43	35	18	14	0	0	0
Levy	CF	18	1	17	0	0	0	0
Marion	CF	43	8	38	6	0	0	0
Palm Beach	SF	50	42	0	3	0	0	0
Pasco	CF	29	3	22	12	4	0	0
St. Lucie	CF	2	2	0	0	1	0	0
Sumter	CF	16	0	16	1	1	0	0
Suwanee	NF	7	1	3	2	0	0	0
Union	NF	9	0	2	4	0	0	0

Number of samples that were identified based on their reaction with the designated antisera. See footnotes of Table 1 for explanation of abbreviations.

Region of state: NF=Northern Florida; CF=Central Florida; SF=Southern Florida.

Table 16. Results of serological assays for six viruses in cucurbits tabulated by region and time of year^z.

Region/time of year ^x	Total # assayed	PRSV-W	# of samples ^y		CMV	SMV	TV
			WMV-2	ZYMV			
NF/Jan.-July	97	3	52	0	6	0	0
NF/Aug.-Nov.	102	46	33	7	0	0	0
CF/Jan.-July	111	4	101	13	5	0	0
CF/Aug.-Nov.	53	49	18	25	2	0	0
SF/Jan.-July	112	83	0	4	0	0	19
SF/Aug.-Nov.	74	60	0	2	0	0	3

^zCombined results of assays conducted in 1987 and 1988.

^yNumber of samples that were identified based on their reactivity with the designated antisera. See footnote of Table 14 for explanation of abbreviations.

^xRegion of state where samples were collected:

NF=Northern Florida; CF=Central Florida; and SF=Southern Florida (see table 15 for counties included in each region). Time of year indicates that samples were collected from January to July or from August to November.

Table 17. Detection of infectious potyviruses in seeds of two squash cultivars collected from a potyvirus--infected field in Fall 1985. Leesburg, FL.

Variety	Number Infected/Number Tested		
	Germinated Seedling ¹	Normal Seed ²	"Paperthin" Seed (aborted) ²
President	0/1300	3/1300 ³	15/450 ⁴
Multipik	0/850	0/850	0/350

¹ Seed transmission.

² Seedcoat ground in buffer and used to mechanically inoculate Small Sugar pumpkin. All infections were identified by subsequent host range tests.

³ All 3 WMV2.

⁴ 12 ZYMV; 2 WMV2 + ZYMV; 1 PRSV-W + ZYMV

Table 18. Host range of ZYMV isolates identified at Leesburg, FL.

Isolate	SS	AP	633-3LA/(SS)*	M	R/(SS)*
81-25	Dk B, Pale	NS	Syst Mild syst mottle, (+)	VC, B	NS/(+)
86-1	SB, Pale	NS	Syst Mild mottle? (-)	Strg VC	NS/(+)
86-4	SB, Pale	NS	Syst Mild syst mottle, (+)	VC, B	NS/(+)
86-8	Dr grn distor	NS	Syst Strg syst mottle, (+)	VC	NS/(+)
86-16	Dk grn distor	NS	Syst Syst mottle (+)	Strg VC, Mottle, B	NS/(+)
FC1119	Dk grn	NS	** Mild syst mottle? (-)	VC	NS/(+)

SS=Cucurbita pepo, 'Small Sugar'; AP=Pisum sativum, 'Alaska'; 633=3 Cucumis melo, 633-3 of P.I. 180280; LA=Luffa acutangula; M=Melothria pendula; R=Ranunculus sardous

*Secondary inoculation to SS from symptomless primary host; +=virus recovered; - =virus not recovered.

Dk=dark, B=blister, SB=speck blister, Pale = pale color, Syst=systemic, VC=vein clearing, Strg=strong, Distor=distortion, grn=green, NS=no symptoms.

**not done.

Table 19. Aphid transmissibility of ZYMV isolates identified at Leesburg, FL.

Isolate	Percent Transmission*				Avg.
	MP	AC	AG-O	AG-L	
81-25	56	41	32	64	46
FC1119	7	0	0	40	12
86-1	81	48	28	74	54
86-4	93	20	25	77	49
86-8	93	47	49	87	66
86-16	13	33	16	77	35
FC1119AT	**	13	20	80	33

* Tests conducted with 3-5 aphids/plant; Donor host=small sugar pumpkin (SS); Receptor host=SS. MP=Myzus persicae; AC=Aphis citricola; AG-O=Aphis gossypii, Orlando biotype; AG-L=Aphis gossypii, Leesburg biotype. Percent transmission is the average of 2 experiments each with 15-20 replications.

** not done.

Table 20. Pathotypes of Florida isolates of ZYMV based on their reactions in *Cucumis melo* 'Doublon' and PI 414723.

^a Possible Reactions in 'Doublon' (homozygous for <u>Fn</u>)	ZYMV pathotype	^b Possible reactions in PI 414723 (homozygous for <u>Zym</u>)	ZYMV pathotype	Florida ZYMV isolates and their pathotypes
Wilting	F	No systemic infection	0	1994,2000,2050=F,0
		Systemic necrotic spots	1	
		Mosaic, leaf deformation	2	
Mosaic, yellowing, vein clearing, leaf deformation, no wilting	NF	No systemic infection	0	1119GH,1119AT,2154=NF,0
		Systemic necrotic spots	1	1119A,1119B = NF,2
		Mosaic, leaf deformation	2	

^{a,b}Possible reactions based on those noted for European isolates.

Table 21. Reaction of nine melon lines to 16 PRSV isolates and WMV-M**

Florida isolates									
melon lines	PRSV-WA	1637	1870	2169	2201	2207	2030*	2052*	2040*
Ouzbeque	VB	VC	VC	VC	VB	VC	VC	VC	VC
Supermarket	VB	VC	VC	VC	VB	VC	VC	VC	VC
Vedrantaïs	VB	VC	VC	VC	VB	VC	VC	VC	TN
VAC	VB	VC	VC	VC	VB	VC	VC	TN	TN
Edisto	VB	VC	VC	VC	VB	VC	VC	TN	TN
PI161375	M	MO	MO	MO	M	M	MO	M	TN
PI414753	TN	TN	TN	TN	TN	TN	TN	LL/SNS	LL
72025	SNS/TN	TN	O/SNS	TN	TN	TN	SNS/TN	SNS	LL
WMR 29	O	LL/SNS	O	O	O	LL	LL/SNS	LL	LL

Other isolates								
melon lines	ATCC	Calif.	Greece	Jordan	NY	PRSV-P	PRSV-T	WMV-M*
Ouzbeque	VC	VC	VC	VB	VC	VC	VC	VC
Supermarket	VC	VC	VC	VB	VC	VC	VC	VC
Vedrantaïs	VC	VC	VC	VB	VC	M	SNS	TN
VAC	VC	VC	VC	VB	VC	M	TN	TN
Edisto	VC	VC	VC	VB	VC	M	O/SNS	TN
PI161375	M	MO	M	MO	MO	M	SNS	TN
PI414753	TN	TN	TN	TN	TN	TN	LL	TN
72025	TN	TN	LL/SNS	TN	TN	LL/SNS	LL	SNS
WMR 29	LL	LL	O	SNS/TN	LL	LL	LL	O

*isolates which did not react with monoclonal antibody F3C-C10

**Abbreviations

susceptible reactions

VC: vein clearing, yellowing, green blister

VB: vein banding

MO: mosaic

M: mottle

resistant reactions

TN: top necrosis

SNS: systemic necrotic spot

LL: local lesions

O: no symptoms

Table 22. Transmission of potyvirus isolates in Israel by *Myzus persicae* and *Aphis gossypii*.

Aphid species	Tests	C-13	C-4	MV	M-c
<i>M. persicae</i>	B	0/60	34/55	0/60	23/68
		0	61.8	0	33.82
<i>A. gossypii</i>	B	0/65	17/66	0/65	42/62
		0	25.75	0	64.57

3 aphids were confined on each test plant.

Table 23. Transmission of purified transmissible and nontransmissible ZYMV isolates mediated by homologous and heterologous helper components.

Helper* source	Virus** isolate	Rate of transmission
ZYMV-13	ZYMV-13	0/13; 0/13; 0/10; 0/10
ZYMV-13	ZYMV	7/11; 6/11; 4/10; 5/10
ZYMV	ZYMV-13	0/11; 0/12; 0/10; 0/11
ZYMV	ZYMV	5/10; 5/10; 6/11; 6/11
ZYMV	PVY	0/10; 0/12; 1/12
PVY	PVY	7/11; 7/12; 6/13; 8/12
PVY	ZYMV-13	0/11; 0/11; 0/10; 0/10
PVY	ZYMV	6/13; 7/13; 5/13; 5/11

Transmissions were carried out as described in Raccach and Pirone (1984).

*Helper extract for ZYMV isolates was obtained from the supernates of infected leaves extracts in 0.1M phosphate buffer pH 7.5 and DIECA. Centrifugation was carried out at 123000g for 2 hrs.

**Virus was presented at a concentration of 100 µg/ml in mixture with the helper solution. The ZYMV isolates are from Israel.

Table 24. Transmission rate of ZYMV-W acquired by aphids from membranes.

1. Supernate (90 μ l) + TSM (10:1)	- - -> 0/12
2. Supernate (90 μ l) + ZYMV-W (10:1)	- - -> 4/12
3. PEG fraction (10 μ l) + ZYMV-W (10:1)	- - -> 3/12
4. AS fraction (10 μ l) + ZYMV-W (10:1) + TSM (80:1)	- - -> 13/14

TSM= 0.1M Tris sulfate pH 7.5 + MgSO₄.

PEG= 8% Polyethylene glycol 6000 fraction obtained from high speed supernate.

AS= 30% ammonium sulfate fraction obtained from high speed supernate.

ZYMV-W= Zucchini yellow mosaic virus (wild type), purified and adjusted to 1 mg/ml.

Aphids were allowed to acquire the solution from parafilm membranes for 10 min, then transferred for an inoculation access feeding for 18 h on virus free zucchini seedlings.

Table 25. Propensity of alate aphids collected in a zucchini field to transmit an isolate of WMV2 and ZYMV in Spring, 1986, Leesburg, FL.

Aphid species ^{1,2}	No. transmitting/no. tested (%)		Total % trans.
	WMV-2	ZYMV	
<i>Myzus persicae</i>	2/22 (9)	13/31 (42)	28
<i>Aphis middletonii</i>	9/49 (18)	15/48 (31)	25
<i>Lipaphis erysimi</i>	0/3 (0)	2/6 (33)	22
<i>Rhopalosiphum rufiabdominalis</i>	0/3 (0)	1/2 (50)	20
<i>Uroleucon</i>	0/1 (0)	1/1 (100)	50
<i>Aphis craccivora</i>	1/5 (20)	0/2 (0)	14
<i>Aphis nerii</i>	1/1 (100)	0/0 (0)	50
<i>Rhopalosiphum spp.</i>	1/1 (100)	0/0 (0)	50

¹Data in table based only on species that transmitted WMV-2 and/or ZYMV in this test. Alate aphids collected by suction and yellow water-pan trap before zucchini field showed symptoms of potyvirus infections.

²Aphids placed on small sugar pumpkin infected with either WMV-2 or ZYMV for a 2-min virus acquisition period and then transferred to Dixie squash receptor plant.

Table 26. Preliminary data comparing serological titer of viral coat protein¹ and helper component² to aphid transmission with several ZYMV isolates and different aphid vectors.

Isolate	OD ₄₀₅ ³	<u>No. infected/no. tested</u>				
		Aphids ⁴				
	CP	AI	MP	AC	AG-O	AG-L
FC1119	.38	.13	0/12	1/12	0/12	3/12
FC1119AT	.83	.34	9/12	2/12	5/12	10/12
86-4	.62	.33	9/12	3/12	5/12	11/12
86-1	.64	.35	—	5/15	—	11/15
86-8	.54	.41	—	7/15	—	15/15
86-16	.61	.36	—	5/15	—	9/15
81-25	.61	.38	—	3/15	—	6/15

¹ Anti ZYMV 113-1 to coat protein (CP)

² Anti PRSV-W amorphous inclusion protein AI 1042 (AI)

³ Indirect ELISA using goat anti-rabbit conjugated with alkaline phosphatase. Plant samples for titer measurement of CP and AI were taken at the acquisition feeding site of the vectors tested.

⁴ 3-5 aphids/plant; donor host=small sugar pumpkin (SS); receptor host=SS.

— = not done; MP=*Myzus persicae*; AC=*Aphis citricola*; AG-O=*Aphis gossypii* Orlando biotype; AG-L=*Aphis gossypii* Leesburg biotype.

Table 27. Transmission of helper component-mediated purified ZYMV (Florida isolates 81–25) by *Myzus persicae*^a.

Virus ^b	Helper component ^b	Transmission ^c
ZYMV	None	0/12
None	High speed supernatant (SN)	0/12
ZYMV	High speed SN	8/12
ZYMV	(NH ₄) ₂ SO ₄ (AS) of SN	1/12
ZYMV	Peak 1 280 nm sucrose grad. AS of SN	0/24
ZYMV	Peak 2 280 nm sucrose grad. AS of SN	0/24

^a This aphid transmission experiment was conducted during Dr. Raccach's visit to Orlando

^b ZYMV isolate 81–25 concentration was 1.4 mg/ml and was used at a ratio of 10 μ l to 90 μ l of helper component (from ZYMV isolate 81–25).

^c Aphids were reared on pepper and starved for 2–4 hr; acquisition access period was 5 min; inoculation access period was overnight. Ten aphids per small sugar pumpkin receptor plant were used.

Table 28. Aphid species caught in yellow water traps during the fall of 1986 and the spring of 1987, at Gilgal in the Jordan Valley, in a spaghetti squash plot.

Aphid species	Fall N	1986 Percent	Spring N	1987 Percent
<u>Acyrtosiphon pisum</u> Harris	116	25.95	7	2.01
<u>Aphis brunea</u>	61	13.65	31	8.91
<u>Aphis citricola</u> v.d. Goot	1	0.22	4	1.15
<u>Aphis craccivora</u> Koch	2	0.45	8	2.30
<u>Aphis fabae</u>	1	0.22	19	5.46
<u>Aphis gossypii</u> Glover	11	2.46	41	11.78
<u>Dacytnotus</u> spp.	1	0.22	0	0.00
<u>Eriosoma lanigerum</u> Hausmann	7	1.57	0	0.00
<u>Macrosiphum euphorbiae</u> Thomas	0	0.00	3	0.86
<u>Myzus persicae</u> Sulzer	15	3.36	169	48.56
<u>Penatlonia nigronervosa</u> Cocq.	150	33.56	6	1.72
<u>Rhopalosiphoninus staphyleae</u> Koch	0	0.00	1	0.29
<u>Rhopalosiphum maidis</u> Fitch	43	9.62	28	8.05
<u>Tetraneura africana</u> v.d. Goot	11	2.46	0	0.00
Not identified	28	6.26	31	8.91
Total	447	100.00	348	100.00

N=Number of aphids trapped in yellow water traps

Percent = Percent of each species in the total aphid population

Table 29. Effect of a combination of mineral oil (Virol) and a pyrethroid (Maverick) on the rate of ZYMV infection in spaghetti squash.

Treatment	Number of Plants	Rate of Infection
Virol 1.5%	132	31.1b
Virol 1.5%+ Maverick 0.05%	127	14.9c
Maverick 0.05%	162	35.2b
O-P (Rogor 40)	164	35.4b
Control	129	46.51a

Treatments followed by the same letter are not significantly different ($p < 0.05$), based on the Chi square test.

Table 30. Proportion of aphid species of the total population trapped in green tiles at Zafaria in Bet-Dagan from April 10 up to May 15, 1988.

	Control	Mav+Virol 0.15%+1.5%	Maverick 0.15%	Virol 1.5%
<u>Aphids</u>				
<u>Aphis citricola</u>	47.895	61.446	29.714	47.340
<u>Hyperomyzus lactucae</u>	2.105	0.482	0.571	2.128
<u>Capitophorus elaeagni</u>	10.526	5.301	9.714	9.574
<u>Macrosiphum euphorbiae</u>	3.684	2.169	6.286	2.128
<u>Aphis gossypii</u>	21.579	24.819	41.143	22.340
<u>Aphis spp.</u>	8.421	1.687	7.429	8.511
<u>Acyrtosiphon pisum</u>	0.000	1.687	1.143	0.000
<u>Myzus persicae</u>	1.053	0.482	1.143	0.532
<u>Uroleucon sonchi</u>	2.632	0.964	0.571	4.787
Other species	2.105	0.964	2.286	2.660

On May 15, there was an extreme heat wave that practically eliminated all the aphids in the vicinity of the experimental plot.

The plots were sprayed once a week from germination to June 6, 1988.

Table 31. Weight (in Kg) of the total melons picked from 50 plants in each treatment (presented in % of marketable fruit versus % of distorted/virus infected melons).

WEIGHT	CONTROL	MAV 0.15%	VIROL 1.5%	MAV+VIROL* 0.15%+1.5%	TOTALS
MARKETABLE	20.3	39	59.1	70.7	189.1
DISTORTED	79.7	61	40.9	29.3	210.9
Totals	100	100	100	100	400

Results of Chi-squared test:

Chi-squared value: 59.56

Degrees of freedom: 3

Table value (P=.05): 7.81

Table value (P=.01): 11.34

Conclusion: VERY SIGNIFICANT

*MAV=Maverick

Table 32. Number of melons picked from 50 plants in each treatment (presented in % of marketable fruit versus % of distorted/virus infected melons).

NUMBER	CONTROL	MAV* 0.15%	VIROL 1.5%	MAV+VIROL 0.15%+1.5%	Totals
MARKETABLE	36.3	58.3	76.5	87.9	259
DISTORTED	63.7	41.7	23.5	12.1	141
Totals	100	100	100	100	100

Results of Chi-squared test:

Chi-squared value: 66.81

Degrees of Freedom: 3

Table value (P=.05): 7.81

Table value (P=.01): 11.34

Conclusion: VERY SIGNIFICANT

*MAV=Maverick

Table 33. No. of plants, mean values of ELISA (absorbance at 405 nm), standard deviations, coefficients of variance (CV), and range of values for zucchini yellow mosaic virus (ZYMV) titer in the cross *Cucurbita maxima* X *C. ecuadorensis*.

GENOTYPE	NO. OF PLANTS	MEAN + SD	CV	RANGE
P1	70	1.157 + 0.159	0.137	0.730-1.4
P2	61	0.046 + 0.019	0.413	0.015-0.08
F1	52	0.206 + 0.103	0.5	0.080-0.47
F2 107	210	0.363 + 0.354	0.975	0.030-1.26
F2 105	224	0.316 + 0.281	0.889	0.050-1.25
P1 X 107	114	0.642 + 0.364	0.566	0.110-1.50
107 X P1	170	0.645 + 0.281	0.435	0.100-1.30
107 X P2	120	0.069 + 0.041	0.594	0.028-0.21
105 X P2	133	0.064 + 0.044	0.687	0.028-0.22
F2 S. Mama	240	0.385 + 0.31	0.805	0.020-1.20

Note. P1=C. maxima; P2=C. ecuadorensis; 107=C. maxima as a female parent; 105=C. ecuadorensis

Table 34. Heterogeneity test (X) for ZYMV titer to reciprocal generations in the interspecific cross *Cucurbita maxima* X *C. ecuadorensis*.

GENERATION A	GENERATION B	MEAN A	MEAN B	P
F2 105	F2 107	0.316	0.363	0.5<P<0.7
107 X P1	P1 X 107	0.645	0.642	0.2<P<0.3
107 X P2	105 X P2	0.069	0.064	0.2<P<0.3
F2 S. Mama	F2 107	0.385	0.363	0.3<P<0.5

a—Each population was divided into three classes as follows: a plant was defined as P1 if it was within two standard deviations from the mean of P1, and the same for P2 and F1. 105=*C. ecuadorensis* as a female parent; 107=*C. maxima* (Golden Nugget) as a female parent; S. Mama=*C. maxima* (Sweet Mama) as a female parent.

Table 35. Estimates of gene effects, standard errors, and t tests for ZYMV titer in the interspecific cross *Cucurbita maxima* X *C. ecuadorensis*.

Gene Effect	Estimate of gene effect	t
m	0.337 + 0.026	12.9**
a	0.608 + 0.02	30.4**
d	-0.255 + 0.117	2.17*
aa	0.140 + 0.113	1.23
ad	0.052 + 0.026	2.01*
dd	0.013 + 0.144	0.09

** Significant at 0.01

* Significant at 0.05

Table 36. Natural WMV-2 and ZYMV infectivity of alate aphids collected in a zucchini field in Spring, 1986, Leesburg, Florida.

Aphid sp.	No. collected	No. infective	% infective ¹	% alate pop. ¹
<i>Aphis middletonii</i>	339	11	0.99	31
<i>Aphis citricola</i>	279	10	0.90	25
<i>Aphis nerii</i>	27	4	0.36	2
<i>Aphis craccivora</i>	33	3	0.27	3
<i>Myzus persicae</i>	117	2	0.18	11
<i>Lipaphis erysimi</i>	311	2	0.18	28
Unknown	2	2	0.18	0.2
Totals ¹	1108	34 ²	3.07	

¹Total no. aphids based only on species that transmitted potyviruses in this test. Alate aphids collected by suction and yellow water-pan trap.

²24 WMV-2; 7 ZYMV; 3 ZYMV + WMV-2.

Figure Legends

- Fig. 1. Immunodiffusion test for serological detection of six viruses that infect cucurbits. The following undiluted antisera prepared to sodium dodecyl sulfate-treated immunogens (Table 1) were added to the central wells: P=papaya ringspot virus type W; W=watermelon mosaic virus-2; Z=zucchini yellow mosaic virus; C=cucumber mosaic virus; S=squash mosaic virus; and T=Trichosanthes virus. Sodium dodecyl sulfate treated leaf extracts from pumpkin plants infected with the following viruses were added to the outer wells: 1=papaya ringspot virus type W; 2=watermelon mosaic virus-2; 3=zucchini yellow mosaic virus; 4=cucumber mosaic virus; 5=squash mosaic virus; and 6=Trichosanthes virus.
- Fig. 2. SDS-immunodiffusion tests with antisera to cucurbit potyviral cylindrical inclusion proteins and extracts from virus-infected plants. Center wells contain cylindrical inclusion (CI) protein antisera collected c.6 weeks after the first immunization; P-antiserum to PRSV-W (WMV-1) CI protein; W=antiserum to WMV-2 CI protein; and Z antiserum to ZYMV CI protein. Outer wells contain SDS-treated extracts from pumpkin: 1= Healthy control; 2= Trichosanthes virus, FC-1860; 3 and 6 = ZYMV; 4= PRSV-W; and 5=WMV-2.
- Fig. 3. Comparison of the sensitivity of ZYMV detection in infected tissue by indirect ELISA using 4-methylumbelliferyl phosphate (fluorescence) and p-nitrophenyl phosphate (optical density) as substrates for alkaline phosphatase conjugate.
- Fig. 4. Western blot analysis of the reactivity of monoclonal antibody MCA F10EA5 (prepared to PRSV-W amorphous inclusion protein) with healthy pumpkin (H), PRSV-W infected pumpkin (I) and purified amorphous inclusion protein (P). After protein transfer, nitrocellulose membranes were incubated in dilutions of 1) Ascites fluid of MCA F10E-A5; 2) Ascites fluid of non-fused SP2/O cells; 3) polyclonal antisera to the amorphous inclusion protein of PRSV-W (1136). Molecular weights on the left side of the figure represent the following marker proteins: phosphorylase b (94k), bovine serum albumin (67k), glutamic dehydrogenase (53k) and carbonic anhydrase (29k).
- Fig. 5. Analysis of *in vitro* translation products of zucchini yellow mosaic virus Reunion isolate (ZYMV-R) by immunoprecipitation and SDS-polyacrylamide gel electrophoresis. RRL (left) represents translation done in the rabbit reticulocyte lysate. Lanes from left to right are: TP, total products of translation in the RRL system; RRL translation products immunoprecipitated by antiserum to PRSV-WAI (papaya ringspot virus type W) amorphous inclusion protein; to CI, ZYMV cylindrical inclusion protein; to TEVNI49K, tobacco etch virus (TEV) nuclear inclusion (NI) 49k protein; to TEVNI54K; to CP, ZYMV-R coat protein (CP); NS, normal serum. WG (right) represents translation done in the wheat germ system. Lanes from left to right are TP, total products of translation in the WG system; WG translation products immunoprecipitated by antiserum to PRSV-W AI, ZYMV CI, TEVNI49K, TEVNI54K, ZYMV-R CP, and NS, respectively. The molecular weights of the major translation products are given on the left of Fig. 5. The translation conditions and the analyses of products were as outlined in Xiong *et al.*, 1988, *Phytopathology* 78:1128-1134.
- Fig. 6. Weekly distribution maps for ZYMV in a spaghetti squash plot in the Gilgal Experimental Station, in the fall of 1986.

- Fig. 7. The incidence of aphid species trapped in yellow waterpans, in a spaghetti squash plot in the Jordan Valley, from Sept. 30 to Dec. 9, 1986.
- Fig. 8. The incidence of aphid species trapped in yellow waterpans, in a spaghetti squash plot in the Jordan Valley, from Mar. 31 to May 20, 1987.
- Fig. 9. Weekly ZYMV infection and incidence of aphid species in green tiles in untreated melon plot.
- Fig. 10. Weekly ZYMV infection and incidence of aphid species in green tiles in melon plot sprayed with pyrethroid (0.15% Maverick).
- Fig. 11. Weekly ZYMV infection and incidence of aphid species in green tiles in melon plot sprayed with oil (1.5% Virol).
- Fig. 12. Weekly ZYMV infection and incidence of aphid species in green tiles in melon plot sprayed with pyrethroid and oil (0.15% Maverick and 1.5% Virol).
- Fig. 13. Effects of pyrethroid (0.15% Maverick) and oil (1.5% Virol) sprays on production of melons. For each treatment, the marketable melons are represented in the pile on the left, and the unmarketable melons are represented in the pile on the right. From top to bottom: P=pyrethroid alone; O=oil alone; PO=pyrethroid plus oil; and C=unsprayed melons.
- Fig. 14. Percent of *Aphis gossypii* alates taking off from treated leaves. The triangles represent Maverick at 0.15% in water and the empty rectangles represent 0.15% Telstar. The dark rectangles represent aphids exposed to leaves treated with water alone.

Fig. 1

Fig. 2

Fig. 3. Comparison of the sensitivity of ZYMV detection in infected tissue by indirect ELISA using 4-methylumbelliferyl phosphate (fluorescence) and p-nitrophenol phosphate (optical density) as substrates for alkaline phosphatase conjugate.

Fig. 4

Fig. 5

[illegible]

P	A	R	E	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X
1																								1
2																								0
3																								2
4																								1
5																								1
6																								0
7																								4
8																								1
9																								0
10																								0
11																								1
12																								1
13																								1
14																								0
15																								0
16																								0
17																								4
18																								2
19																								1
20																								2
21																								1
22																								1
23																								1
24																								0
25																								2
26																								1
27																								0
28																								0
29																								1
30																								0
31																								1
32																								1
33																								1
34																								0
35																								0
36																								1
37																								2
38																								1
39																								0
40																								1
41																								2
42																								3
43																								2
44																								2
45																								1
46																								0
47																								2
48																								1
49																								1
50																								0
51																								0
52																								0
53																								0
54																								1
55																								1
56																								2
57																								2
58																								2
59																								3
60																								1
61																								0
62																								0
63																								1
64																								1
65																								0
66																								0
67																								0
68																								0
69																								1
70																								2
71																								1
72																								1
73																								4
74																								0
75																								0
76																								0
77																								0
78																								0
79																								0
80																								0
81																								0
82																								0
83																								0
84																								0
85																								0
86																								0
87																								0
88																								0
89																								0
90																								0
91																								0
92																								0
93																								0
94																								0
95																								0
96																								0
97																								0
98																								0
99																								0
100																								0

Fig. 6
2nd WEEK

1 5 2 0 0 3 5 3 9 1 6 6 3 0 0 1 0 4 2 0 4 7 1 52

Fig. 6
3rd WEEK

	A	P	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V	W	X	Y	Z
1																										2
2																										2
3																										2
4																										1
5																										3
6																										1
7																										4
8																										1
9																										1
10																										1
11																										2
12																										3
13																										1
14																										6
15																										6
16																										1
17																										1
18																										3
19																										1
20																										2
21																										1
22																										3
23																										3
24																										3
25																										1
26																										2
27																										2
28																										1
29																										2
30																										2
31																										2
32																										3
33																										1
34																										3
35																										1
36																										4
37																										2
38																										1
39																										1
40																										2
41																										2
42																										2
43																										2
44																										3
45																										2
46																										2
47																										4
48																										2
49																										2
50																										4
51																										1
52																										1
53																										6
54																										2
55																										2
56																										5
57																										5
58																										2
59																										3
60																										2
61																										6
62																										2
63																										1
64																										4
65																										2
66																										3
67																										1
68																										3
69																										2
70																										2
71																										2
72																										2
73																										6
74																										1
75																										1
76																										4
77																										4
78																										4
79																										4
80																										4

Fig. 6
4th WEEK

Fig. 7

Fig. 8

Weekly ZYMV infection and incidence of aphid species in green tiles

Fig. 9

Weekly ZYMV infection and incidence of aphid species in green tiles

Fig. 10

Weekly ZYMV infection and incidence of aphid species in green tiles

Fig. 11

Weekly ZYMV infection and incidence of aphid species in green tiles

Fig. 12

Fig. 13

Percent of *A. gossypii* alates
taking-off from treated leaves

Fig. 14

F. Description of Cooperation.

Dr. B. Raccah visited in Leesburg and Gainesville, Florida twice for 2 days in 1986. He met with the late Dr. Warren Adlerz to coordinate epidemiological experimentation. He also visited Leesburg to coordinate the monitoring of aphids and virus infections with Dr. Ray Yokomi and Ms. Debra Howd. In 1988, Dr. Raccah visited the lab of Dr. Yokomi in Orlando and presented a seminar on his research. These visits provided opportunities for many of the project investigators to jointly discuss research progress and were very helpful.

Dr. R. Yokomi visited Israel in the fall of 1986 to work with Dr. Raccah on techniques of assessing helper component activity. Epidemiology was discussed and coordinated, and Dr. Yokomi joined Dr. Raccah in assessing the experiments on virus disease control in the Jordan Valley.

During his sabbatical leave in the Department of Plant Pathology, University of Florida from Aug. 1987 to May, 1988, Dr. Herve Lecoq of the INRA in France collaborated on research projects of cucurbit potyviruses and had discussions with the project investigators. Dr. Lecoq's collaboration and advice is gratefully acknowledged.

Polyclonal antisera to Israel and Florida isolates of ZYMV were exchanged. Polyclonal antisera to Florida isolates of cucurbit potyviruses and to PRSV-W amorphous inclusion protein were provided to R. Yokomi. Polyclonal antiserum to a Florida isolate of WMV-2 was tested in Israel with cucurbit potyvirus isolate Mc to confirm its similarity to WMV-2. Monoclonal antibody to ZYMV produced in Gainesville, Florida, was sent to Israel and it was reactive with ZYMV from Israel.

G. Evaluation of research achievements with respect to aims of the original proposal.

1. Production of polyclonal antisera and monoclonal antibodies for use in development of serological tests for detecting and distinguishing PRSV-W, ZYMV, and WMV-2.

Ten polyclonal antisera were produced in Florida and Israel to virus and capsid proteins associated with ZYMV, PRSV-W, and WMV-2. Two antisera also were produced to a possible potexvirus (TV) isolated from *Trichosanthes* in Florida.

Monoclonal antibodies were produced to PRSV-W capsid protein, PRSV-W, ZYMV, and WMV-2.

The polyclonal antisera were used in various tests depending on the purpose and location. Immunodiffusion tests in an SDS system were used for analysis of specimens collected in surveys and for assessment of relationships. ELISA, and SSEM also were used for studying relationships. ELISA tests were used with monoclonal antibodies.

Serological tests were used to assay samples collected in disease surveys in Florida. PRSV-W was detected in south Florida at various times of the year, but it did not become abundant in central or northern Florida until late summer and fall. The WMV-2 was the most abundant potyvirus of cucurbits in central and north Florida in the spring and also occurred there in the fall. WMV-2 was not detected in any of the three southern Florida counties surveyed. ZYMV was detected in various parts of the state, but was only about 1/5 to 1/4 as abundant as PRSV-W and WMV-2, respectively.

The antisera produced in this test have been used extensively in this project; they will be used in BARD project proposal No. U.S. 1390-87C, and will be useful for in the future for a variety of applied and basic research projects in both Florida and Israel.

2. To develop immunological probes for virus-specified, nonstructural proteins (e.g., cylindrical inclusion and helper component proteins).

Polyclonal antisera were produced to cylindrical inclusion proteins (CIP) and amorphous inclusion protein (AIP) associated with PRSV-W, CIP and helper component protein associated with ZYMV, and to CIP associated with WMV-2. Monoclonal antibody to PRSV-

W AIP also was produced.

The PRSV-W AIP polyclonal and monoclonal antibodies were shown to have a broad range of activity among PRSV-W isolates. A serological relationship of PRSV-W CIP and AIP to corresponding proteins induced by zucchini yellow fleck, another potyvirus, was established by ELISA and immunofluorescence.

Antiserum to helper protein has been useful in assessing relationships with helper component induced by other potyviruses and in tests involving neutralization of helper component activity. The antisera to nonstructural proteins were used for immunoprecipitation of in vitro translation experiments involving ZYMV.

Although no extensive tests have been conducted with a battery of virus isolates, preliminary tests indicate that the CIP antisera could be useful for virus identification.

3. Determine wild hosts and alternative crop hosts of principal potyviruses that infect cucurbits in Israel and Florida and to assess these hosts as natural sources of virus for aphid transmission.

Host ranges and aphid transmissibility of ZYMV isolates from various regions in Israel were studied. The ZYMV was detected in the Arava of Israel in umbelliferous hosts in the winter, and possibly these plants play a role as overwintering hosts.

Host range and aphid transmissibility of Florida ZYMV isolates also was determined. Seed transmission of ZYMV, PRSV-W, and WMV-2 in squash was not detected, although virus was detected in seed by mechanical transmission to indicator plants.

4. Assess variability of ZYMV, PRSV-W, and WMV-2 with regard to host range, host reaction and antigenicity.

Host range studies confirmed the biological variability of Israel and Florida isolates of ZYMV. Variation in the reactivity of melon lines to Florida isolates of PRSV-W was established. Antigenic variation in isolates of PRSV-W was detected by both polyclonal antisera and monoclonal antibodies. In Florida, more than 200 virus cultures were preserved in dried leaf tissue for further evaluation.

5. Determine major vector species and use these vectors to determine transmissibility of different isolates.

The composition of aphid species differed markedly by season from fall (1986) to spring (1987) in the Jordan Valley. Secondary spread of ZYMV was noted in two cucurbit fields in Israel. A survey of aphid species in the Arava Valley of Israel detected 33 species on wild plants and 23 in traps in cultivated plots. *Myzus persicae* was the most frequently trapped aphid in the Arava in winter 1986. A combination oil-pyrethroid spray and oil alone resulted in increased yields of marketable melons in Israel, and these programs have been recommended to growers as a means of reducing the effects of ZYMV.

Ammonium sulfate precipitation of ZYMV helper component was superior to precipitation with polyethylene glycol for preservation of helper component activity. Helper components prepared from both transmissible and nontransmissible isolates in Israel were all active in assisting transmission of purified transmissible isolates. A relationship between ZYMV and potato virus Y helper components was established by biological tests and by serology.

Myzus persicae was more efficient than *Aphis gossypii* and *A. citricola* for transmission of Florida isolates of ZYMV. Differences were noted in the vector efficiency of biotypes of *A. gossypii*. The most important vector species of WMV-2 and ZYMV at Leesburg were *Aphis middletonii*, *Myzus persicae* and *A. citricola*, based on abundance, transmission efficiency, and natural infectivity. *Lipaphis erysimi* also was a vector of WMV-2 and ZYMV.

Changes in the project. Dr. Warren Adlerz, an entomologist in Florida and one of the

original co-principal investigators of the project, died following a brief illness less than 10 months after this project began. Dr. Adlerz had already initiated various experiments relating to objectives 3,4, and 5. Dr. Ray Yokomi joined the project, and brought many of the experiments that Dr. Adlerz had begun to conclusion. Dr. Yokomi also traveled to Israel to confer with Dr. Raccah and other scientists there.

H. List of publications.

- Baker, C.A., and Purcifull, D.E. 1987. Serological relationships of three proteins of papaya ringspot virus type W to antigens of zucchini yellow fleck virus. *Phytopathology* 77:1722 (Abstr.).
- Baker, C.A., and Purcifull, D.E. 1986. Reactivity of a monoclonal antibody to the amorphous inclusion protein of papaya ringspot virus-type W (PRSV-W). *Phytopathology* 78 (in press) (Abstr.).
- Eastop, V.F., and Raccah, B. 1988. Aphid and host plant species in the Arava Valley of Israel: Epidemiological aspects. *Phytoparasitica* 16:23-32. (Appendix 1).
- Paran, I., Shifriss, C., and Raccah, B. 1989. Inheritance of resistance to zucchini yellow mosaic virus in the interspecific cross *Cucurbita maxima* X *C. ecuadorensis*. *Euphytica* (in press).
- Purcifull, D.E., Simone, G.W., Baker, C.A., and Hiebert, E. 1988. Immunodiffusion tests for six viruses that infect cucurbits in Florida. *Proc. Fla. State Hort. Soc.* 101 (Abstract was presented at annual meeting and accompanying paper submitted).
- Yokomi, R.K., Adlerz, W.C., and Howd, D.S. 1987. Biological characterization of several zucchini yellow mosaic virus isolates from central Florida. *Ent. Soc. America Ann. Mtg.* (Abstr.).