BARD

77

PROJECT NO. I - 1104 - 86

FINAL REPORT

**Molecular and Pathogenic Characterization of
the Hemangiosarcoma Inducing Virus of
Poultry**

M. Kotler, R.F. Smith, M. Malkinson

1991

B.

TABLE OF CONTENTS

A. Cover page	page	1
B. Table of contents	"	2
C. Abstract	"	3
D. Objectives	"	4
E. Detailed description of the research activities	"	5
F. Description of cooperation	"	
G. Evaluation of the research achievement with respect to the original proposal and its objective	"	
H. Conclusion and plan for the coming year	"	
I. List of publication	"	

C. ABSTRACT:**MOLECULAR AND PATHOGENIC CHARACTERIZATION OF THE HEMANGIOMA
INDUCING VIRUS**

Sporadic outbreaks of hemangioma in chickens (HA) have occurred in Israel since 1983. We have isolated a new field strain of avian hemangioma virus (AHV) from such a spontaneous tumors. This virus was biologically cloned and induces HA in about 35-45% of line 10 chickens injected on the day of hatching. AHV has a very wide host range. It propagates in cultures established from chicken, quail, duck, goose and turkey cells. The AHV was molecularly cloned in pBR322 and this clone was able to yield virus (AH-52) upon transfection of QT-6 cells. AH-52 has been defined as an A\C subgroup virus and induces 100% of lymphatic leukemia and 15-20% of hemangiomas in SC chickens. The AH-52 induces 25% AH and 25% lymphatic leukemia in line 10 chickens. The HAs are multifocal vascular neoplasms consisting of freely anastomosing vascular channels with areas resembling cavernous hemangioma.

AHV and AH-52 induce a cytopathic effect (CPE) on cultured avian and mammalian cells shortly (8-48h) after infection. The kinetics of killing were dependent on the m.o.i. and independent of virus replication (since UV irradiated virus led to cell death). We have shown that the env gp85 is responsible for the CPE by exchanging the env regions of AHV and RAV-1.

Since endothelial cells (EC) are involved in HA and these lesions are normally associated with blood clotting we have assayed the effect of the AHV on cultured endothelial cells. The env gp85 of AHV is responsible for the loss of thromboresistance of human and bovine EC. The induction of thrombogenicity is associated with a reduction in prostacyclin release and increased expression of tissue factor.

By sequencing the env of AH-52 we have identified elements that may be responsible for the pathogenicity of this virus. Using molecular recombination technique we are attempting to define the precise regions in env that are responsible for the induction of CPE, EC perturbation and pathogenicity.

D. OBJECTIVES

- a. Typing of the AHV.
- b. Defining the chicken breeds that are sensitive to AHV.
- c. Studying the pathogenic potential of AHV.
- d. Defining the genomic elements which confer the potential to induce hemangiosarcoma.

E. DETAILED DESCRIPTION OF THE RESEARCH ACTIVITIES AND RESULTS

1. Epidemiology and Pathogenicity

Outbreaks of hemangioma occurred sporadically on certain poultry farms in Israel during 1988 to 1990 and even at the present time this disease is continuing to threaten certain larger flocks in the country. Laying birds with skin lesions were collected and tumors which were present in the internal organs were removed, co-cultivated with QT-6 cell lines, and the new viruses were isolated for further characterizations. The new isolates were identical to the one isolated by us in 1983 by three criteria: 1) induction of cytopathic effect (CPE) on avian and mammalian cultured cells, 2) restriction enzyme map, and 3) the disease that was induced in experimentally infected chickens. Twelve hens were necropsied (in collaboration with Dr. Soffer, Department of Pathology, Hadassah Hospital) and the morphological features of the spontaneous, multifocal vascular neoplasms were described.

Histologically, the tumors were characterized by areas consisting of freely anastomosing vascular channels with a prominent capillary-like appearance and lined by bland-looking endothelial cells, which alternated with areas resembling cavernous hemangioma. Occasionally, solid areas composed of plump, pleomorphic cells were also present. Although there was no clear evidence for metastatic spread, the tumor was obviously invasive. Electron microscopy and immuno histochemistry confirmed the nature of the neoplastic cells, demonstrating in particular pinocytotic vesicles, well developed junctional complexes, fragmented basal lamina, occasional Weibel-Palade bodies and patchy factor VIII related antigen immunoactivity. C-type retrovirus particles were seen budding from the tumor cells. The morphology and histology feature of the hemangioma was described in detail by us (Soffer D., Resnick-Roquel, N., Eldor, A., and Kotler, M., Cancer Res. 50:4787-4793 (1990).

2. Neoplasms Induced by AHV

The AHV isolated from a layer hen with hemangioma was propagated on CEF. A biological clone of this virus was used to infect chicks on the day of hatching as well as embryonated eggs. About 50% of the birds that were exposed to the virus developed either cavernous hemangiomas or B cell lymphomas (see Table 1, 1st Annual Report). Our results clearly indicate that AHV is an oncogenic virus. The AHV belongs to the group of chronic oncoviruses, and retains its potential to induce hemangiomas after cloning and successive passage in CEF cultures.

The AHV₅₂ (molecular clone of AHV - see below) propagated in QT-6 cells were used for infection of line 10 chicks. This experiment was severely interrupted by Marek disease. However, 3-5 cavernous hemangiomas out of 35 birds appeared in the experimental group, but not in the mock-infected control group. This experiment will be repeated in the near future.

This information was collected in collaboration with Dr. Malkinson, Beit Dagan, Israel. The AHV₅₂ proviral clone was transferred to Dr. R.E. Smith, Department of Microbiology, Colorado State University at Ft. Collins. Dr. Smith then used this DNA to transfect CEF and harvested the virus progenies (AHV₅₂ virus). The AHV₅₂ virus (10^5 infectious units) was inoculated intravenously to

14-day-old embryonated eggs of the SC chickens. All the hatched chicks (15) developed B cell lymphomas between the ages of 3 to 21 weeks. In addition to the lymphomas, one of them also had a hemangioma and another developed a fibrosarcoma. The short latent period, together with the high frequency of lymphomas and the appearance of two different malignant diseases in a single chicken indicated that AHV₅₂ is a highly efficient oncogenic virus.

In the Colorado laboratory it was demonstrated that AHV₅₂ induced B-cell lymphomas; while in our experiments at The Hebrew University, the virus caused mostly hemangiomas. This can be explained by the different route of infection and by differences in lines of chickens used by the investigators in Israel and the U.S.A.

3. Mortality of Embryonated Eggs Caused by AHV

AHV, AHV₅₂, and RAV-1 were inoculated into the yolk sac of 4-day-old embryonated eggs belonging to seven lines of chickens (Table II).

Equal titers of viruses were used in all these experiments, as estimated by reverse transcriptase assay. The incubated eggs were examined on alternate days by illumination, and mortality of the chick embryos was determined (Table II). The virus inoculation and candling of eggs were carried out by Dr. M. Malkinson, Kimron Veterinary Research Institute, Bet Dagan. The results indicated that AHV₅₂ killed the chick embryos of lines Silver, White leghorn and New Hampshire at higher rates and in a shorter latent period than does RAV-1. Interestingly, White Leghorn and leghorn embryos were very susceptible to both AHV strains but not to RAV-1. The embryos of these lines died within 2 to 3 days post-inoculation. Embryos of the Silver line were more resistant to AHVs and RAV-1. These results should be of great help to chicken breeders and should be considered when planning the commercial layer lines to be used in Israel during the coming years.

These results, which demonstrate that AHVs killed embryos, correlate well with our previous findings on the CPE induced by AHV in avian and mammalian cultured cells. It also correlated well with our finding that AHVs perturbed endothelial cells. At present, we are studying causes for embryo deaths induced by AHVs.

4. Determination of AHV Subgroup

Subgroup of an avian leukemia virus is defined by its host range, interference and neutralization antibodies. The AHV has a very wide host range. It propagates well in chick, duck, goose, turkey and quail cells. Our experiments to type the AHV by interference experiments have failed probably because of the CPE it induced in culture. Experiments to prevent the CPE by interference also failed, probably because the CPE induced by AHV envelope is mediated via receptors other than those used by viruses of subgroup A, B, C, and D (these experiments were done in Jerusalem). Dr. Smith (Fort Collins, CO) has used his specific neutralization antibodies and chick host cells to type the AHV₅₂ cloned virus.

Typing of AHV₅₂ was conducted in two ways. A standard number of AHV₅₂ infectious units was incubated with chicken antisera specific for avian leukosis virus of subgroups A, B, C and D. Following incubation, virus/antiserum mixtures were added to monolayers of C/O chicken embryo fibroblasts, transferred once, and reverse transcriptase assays were performed on supernatant fluids after 6 days of incubation. The results indicated that antisera specific for subgroups A and C inhibited AHV₅₂ reverse transcriptase activity, while antisera for subgroups B and D were not inhibitory. In a different approach, AHV₅₂ was inoculated into 6-week old roosters, the inoculation was repeated one month later, and serum was obtained after an additional 30 days. Six sera were obtained, and all of them inhibited AHV₅₂ reverse transcriptase activity in a test conducted as outlined above. The two AHV₅₂ antisera showed the highest amount of inhibitory activity were used in a neutralization test of viruses of subgroups A, B, C and D. In this test, viruses of subgroups A, B, C, and D were incubated with AHV₅₂ antisera, the mixture was inoculated onto C/O fibroblasts, the cells were transferred once, and supernatant fluids were tested for reverse transcriptase activity after 6 days of incubation. The results showed that the anti-AHV₅₂ antisera neutralized viruses of subgroups A and C, but not viruses of B or D subgroups. These results suggest that the AHV₅₂ surface glycoprotein consists of a mosaic of determinants comprising antigens from both A and C subgroups.

5. Biological Characterization of AHV

A. CPE induced by AHV. The field strain AHV causes CPE on cultured avian and mammalian cells. The CPE is induced by AHV shortly after infection. The kinetics of the cell killing depend on the multiplicity of infection. The cytopathic effect on avian and mammalian cells is independent of virus replication, since UV-irradiated virus causes cell death as well. Biochemical and genetic experiments indicate that the AHV *env* gene products are responsible for the CPE: i) Partially purified AHV envelope glycoproteins (gp85), but not those of Rous sarcoma virus, Prague-C strain (Pr-C), induce CPE. ii) Rous associated virus-1 (RAV-1) in which the *env* region was replaced by the AHV gp85 region, induces CPE on avian and mammalian cultured cells.

Therefore, we suggest that CPE is induced by AHV via an interaction between the viral gp85 and cell membrane components (receptors). This mode of CPE would be unique among the avian sarcoma leukemia viruses (details of this work presented in the enclosed manuscript (Resnick-Roguel et al., J. Virol. 63:4325-4330, 1989)).

B. Perturbation of endothelial cells by AHV. Since large thrombi are associated with spontaneous and induced hemangiomas in chickens, we have investigated the effect of AHVs on certain specialized endothelial cell functions. Unfortunately, all our attempts to cultivate avian endothelial cells failed, so we have used human and bovine endothelial cell (EC) lines and the following results have been achieved:

1) Bovine endothelial cells (BAEC) subconfluentlly plated on endothelial extracellular matrix (ECM) coated dishes prevent platelet aggregation. BAEC infected with AHV lost their capacity to prevent platelet aggregation as early as 5 hrs post infection (p.i.), prior to any apparent damage; 2) The rate of

prostacyclin (PGI₂) secretion by AHVs treated human and bovine EC is reduced 2 to 3-fold, as compared with the untreated cells. The viral inhibitory effect is evident as early as 3 hrs p.i.; 3) Infection of BAEC and human EC with AHV caused 60-140-fold increased amounts of tissue factor (TF) activity over that produced by mock-infected cells; 4) The perturbation of EC is a unique characteristic of AHV because infection of the same cultured cells with RAV-1, another member of the avian leukosis virus group whose genomic sequences are closely related to AHV, did not impair the capacity of BAEC to prevent platelet aggregation; and 5) the perturbation of EC is caused by AHV envelope and virus propagation is not essential for this effect because: a) ultraviolet-inactivated AHV induced perturbation of EC; b) RAV-1, a closely related virus, could not affect EC cultures; c) recombinant viruses containing the AHV *env* induce platelet aggregation, but viruses containing RAV-1 were unable to induce this effect. (Details of this study are presented in the enclosed manuscript: Resnick-Rogoul et al., J. Virol. 64:4029-4032, 1990.)

6. Sequencing of AHV₅₂ env to LTR region

In an attempt to define the genetic elements which account for the unique biological features of AHV, we determined the nucleotide sequences of the *env* gene, the 3' non-coding region and the LTR of AHV₅₂. Computer comparison analyses with known sequences of other subgroup A thru E ALVs revealed unique sequences in the gp85 coding region of the *env* gene and the U₃ region in the LTR of AHV₅₂.

Sequencing the amino acid in the gp85 coding region has shown that AHV₅₂ differs from RAV-1 by six amino acids. Three of these amino acids are unique to AHV₅₂ since they are not found at the same location in other members of the ASLV family. One of these amino acids, an *Asn* residue at the N-terminus of gp85, creates a new potential site (*Asn-Arg-Thr*) for glycosylation. The other two amino acids unique to AHV₅₂ are *Leu* and *Gly* residues, located in the hr-1 region.

Sequence comparison at the 3' non-coding region of AHV₅₂ was possible only with that of RAV-2, and this has shown the two viruses to be identical.

In the U₃ region of the LTR (position 2253-2280), a novel tandem repeat composed of two stretches, each of 12 nucleotides, was identified. This element could not be found in RAV-0, RAV-2, SR-A and PR-C. Additional direct repeats were observed in the U₃ region of the AHV LTR at position 2365-2380. These repeats are composed of two stretches, 14 base pairs each, overlapping each other by GTG. This element, which is common both to AHV and RAV-2, is not found in RAV-0, where both stretches are part of a 30-base pair deletion. In SR-A and PR-C, only one of the two stretches could be found in this region. The rest of the AHV LTR is very conserved and cannot be distinguished from the LTRs of the other ASLVs. The details of this study are presented in the enclosed manuscript: Burstein et al., Virology 179:512-516, 1990).

7. Defining the Genomic Elements Which Confer the Potential to Induce Hemangioma

During the first two years of this study we have demonstrated that the AHV *env* glycoprotein gp70 is responsible for the induction of CPE and endothelial cell perturbation. The sequence analysis of the *env* gene revealed unique regions that might be responsible for the biological characteristics of this virus. In order to define these genomic elements, the N' or the C' termini of the AHV *env* gene were replaced with the homologous part of the RAV-1 gene. These heterologous *env* genes were introduced either to RAV-1 or AHV viral DNAs. QT-6 cells were transfected with the four recombinant viral DNA constructs and viruses were collected to assay their biological effects on avian and mammalian cells. These experiments are being conducted at the present time in the Jerusalem laboratory.

8. Production of Transgenic Chickens Containing the AHV Envelope Gene

AHV and RAV-1 construct DNAs deleted the *Bgl*III-*Bgl*III fragment which contained portions of the *gag* and *pol* genes, were used to transfect fresh roosters' sperm cells. PBR322 DNA was also used as a control. White leghorn chickens were artificially assimilated with the transfected sperm cells and several batches of embryonated eggs were collected. These experiments are carried out in collaboration with Drs. A. Feinsode and Y. Gruenbaum of the Hebrew University, Jerusalem. The presence of the deleted proviruses (Δ AHV and Δ RAV-1) in embryos or chickens is determined by Southern blots using a *Bam*HI 1900-bp fragment, and *Bgl*III-*Bgl*III 2600 bp as probes.

Our very preliminary experiments showed that 0-20% of the embryonated eggs contain Δ AHV or Δ RAV-1 proviruses. In addition, 2/11 chickens hatched from eggs laid by mothers assimilated with Δ AHV transfected sperm cells, had hematomas on their skin. None of the RAV-1 or mock-transfected embryos showed this syndrome. However, in another experiment where 21 Δ RAV-1 transfected embryonated eggs were incubated, most of the embryos (18) died before hatching.

At the present time we are concentrating on improving the transfection methodology and developing new vectors expressing the *env* gene of AHV and RAV-1, which may reduce the probability of recombination with the endogenous viruses.

F. DESCRIPTION OF COOPERATION

The AHV₅₂ was used for infection of embryonated eggs. The malignant diseases developed in the hatched chicks were described by Dr. R.E. Smith, Colorado State University. The typing of AHV₅₂ was also carried out by Dr. R.E. Smith using specific antibodies and susceptible chick embryonic cells. The AHV and AHV₅₂ preparations were used to study the susceptibility of chick embryos belonging to several lines. These experiments were carried out by Dr. M. Malkinson, Kimron Veterinary Research Institute.

There was a constant connection among the three laboratories. Dr. Kotler and Dr. Malkinson have met with each other almost weekly, and often speak on the telephone. Communication between Israel and the American laboratory was carried out via the Bitnet system, telephone calls, and mail. Dr. R.E. Smith visited Israel during the first year of this proposal and spent a week in

Jerusalem and two days in the Kimron laboratories. Dr. Malkinson visited the Fort Collins laboratories twice during the second and third year of this proposal, and Dr. M. Kotler stayed for a few days at Fort Collins in the third year.

The connections among the three laboratories are very tied and all three principal investigators are willing to continue this research in the coming years.

G. EVALUATION OF THE RESEARCH ACHIEVEMENT WITH RESPECT TO THE ORIGINAL PROPOSAL AND ITS OBJECTIVE

- 1) Typing of AHV was done in the U.S. laboratory.
- 2) Defining the chicken breeds that are sensitive to AHV has been done in the Kimron Institute, In Jerusalem and in the U.S. laboratories (see the report from year 2).
- 3) Studying the pathogenic potential of AHV has been done in Jerusalem by studying the CPE and endothelial perturbation. In the U.S.A. and in the Kimron laboratories, the pathogenic potential of the virus has been arranged by infection of embryonated eggs. Morphological and histological studies of the hemangioma has been carried out in Jerusalem.
- 4) Defining the genomic elements which confer the potential to induce hemangioma was studied in Jerusalem by: a) sequencing of the AHV envelope and LTR regions and comparing them to those of other ALVs; b) constructing recombinant viruses in which the *env* was replaced with that of RAV-1; c) constructing recombinant virus in which specific regions of the *env* gene were replaced with that of RAV-1; d) assaying these recombinants for their capacity to induce CPE of avian and mammalian perturbation of endothelial cells.
- 5) Defining the breeder lines which are sensitive to AHV.
- 6) The AHV *env* gene is being introduced into the chicken germ line to study the effect of this gene on the development of chick embryos and to define its pathogenic effect.

H. CONCLUSIONS AND PLANS FOR THE COMING YEAR

During the three years we worked on this proposal, we characterized the AHV by biological and biochemical methods:

- 1) We cloned, mapped and sequenced a major part of the AHV genome.
- 2) We defined additional biological characteristics of AHV. AHV induced endothelial cell perturbation. This characteristic is in line with the blood coagulation associated with the tumors in chickens.
- 3) The *env* gene of the virus was defined as the determinant for CPE and endothelial perturbation.

- 4) The regions inside the *env* gene responsible for the biological characteristic of the virus, namely, oncogenic potential, CPE induction, and endothelial perturbation are being defined.
- 5) We have found that AHV induced B-lymphomas on i.v. injection of all the 14-day-old chick embryos belonging to the SC line. Thus, AHV has been proven a very efficient oncogenic virus which can induce various malignant diseases in the lines of chickens used.
- 6) We have defined the susceptibility of chicken embryos to AHVs. This study will be extended to determine the genetic components in the AHV genome that are responsible for causing embryonic death. We will also study the histological effects causing the death.

The three principal investigators have applied for a BARD grant to continue this project.

I. LIST OF PUBLICATIONS

- Resnick-Roguel, N., Burstein, H., Hamburger, J., Panet, A., Eldor, A., Vlodavsky, I. and Kotler, M. Cytocidal effect caused by the envelope glycoprotein of a newly isolated avian hemangioma-inducing retrovirus. *J. Virol.* 63k 4325-4330 (1989).
- Resnick-Roguel, N., Eldor, A., Burstein, H., Hy-am, E., Vlodavsky, I., Panet, A. and Kotler, M. The envelope glycoprotein of avian hemangioma retrovirus induces a thrombogenic surface on human and bovine endothelial cells. *J. Virol.* 64, 4029-4032 (1990).
- Soffer, D., Resnick-Roguel, N., Eldor, A. and Kotler, M. Multifocal vascular tumors in fowl induced by newly isolated retrovirus. *Cancer Res.* 50, 4787-4793 (1990).
- Burstein, H., Resnick-Roguel, N., Hamburger, J., Arad, G., Malkinson, M. and Kotler, M. Unique sequences in the *env* of avian hemangioma retrovirus are responsible for cytotoxicity and endothelial cells perturbation. *Virology* 179, 512-516 (1990).