77

BARD

FINAL REPORT

PROJECT NO. US-2196-92

**Sustained Suppression of Pythium Disease:
Interactions between Compost Maturity and
Nutritional Requirements of Biocontrol Agents**

H.A.J. Hoitink, L.V. Madden, Y. Hadar, Y. Chen

1997

DATE: January 15, 1997BARD project No. US-2196-92

Title:

Sustained Suppression of Pythium Diseases: Interactions Between CompostMaturity and Nutritional Requirements of Biocontrol AgentsInvestigators' Names
(Principal listed first)H. A. J. HoitinkL. V. MaddenY. ChenI. Hadar

Investigators' Institutions

The Ohio State UniversityThe Ohio State UniversityUniversity of JerusalemUniversity of JerusalemProject's starting date: June 7, 1993Type of Report: 1st Annual ☐ 2nd Annual ☐ Final ☒
Signature

Principal Investigator

Signature

Institution's Authorizing Official

**SUSTAINED SUPPRESSION OF PYTHIUM DISEASES:
INTERACTIONS BETWEEN COMPOST MATURITY
AND NUTRITIONAL REQUIREMENTS OF
BIOCONTROL AGENTS.**

(#US-2196-92)

Table of Contents

I.	ABSTRACT	1
II.	OBJECTIVES OF ORIGINAL PROPOSAL	2
III.	RELEVANT DATA	2
	A) Compost Quality:	2
	B) Beneficial Utilization of Compost	3
IV.	DESCRIPTION OF COOPERATION	6
V.	RESEARCH ACHIEVEMENTS/BENEFITS TO AGRICULTURE	6
VI.	LIST OF PUBLICATIONS FROM PROJECT	7
VII.	PATENTS	8
VIII.	APPENDIX I. - RESEARCH PERFORMED IN OHIO	9
IX.	APPENDIX II - RESEARCH PERFORMED IN ISRAEL	56

SUSTAINED SUPPRESSION OF PYTHIUM DISEASES: INTERACTIONS BETWEEN COMPOST MATURITY AND NUTRITIONAL REQUIREMENTS OF BIOCONTROL AGENTS.

(#US-2196-92)

I. ABSTRACT

Several procedures were developed that predict maturity (stability) of composts prepared from municipal solid wastes (MSW). A respirometry procedure, based O_2 uptake by compost, predicted ($R^2=0.90$) the growth response of ryegrass in composts and an acceptable level of maturity. Spectroscopic methods (CPMAS 13 -NMR and DRIFT spectroscopy) showed that the stabilizing compost contained increasing levels of aromatic structures. All procedures predicted acceptable plant growth after approximately 110 days of composting. MSW compost suppressed diseases caused by a broad spectrum of plant pathogens including *Rhizoctonia solani*, *Pythium aphanidermatum* and *Fusarium oxysporum*. A strain of *Pantoea agglomerans* was identified that caused lysis of hyphae of *R. solani*. Evidence was obtained, suggesting that thermophilic biocontrol agents also might play a role in suppression.

13 C-NMR spectra revealed that the longevity of the suppressive effect against *Pythium* root rot was determined by the concentration of readily biodegradable carbohydrate in the substrate, mostly present as cellulose. Bacterial species capable of inducing biocontrol were replaced by those not effective as suppression was lost. The rate of uptake of 14 C-acetate into microbial biomass in the conducive substrate was not significantly different from that in the suppressive substrate although specific activity was higher. The suppressive composts induced systemic acquired resistance in cucumber roots to *Pythium* root rot and to anthracnose in the foliage. Composts also increased peroxidase activity in plants but the conducive substrate did not have these effects. In summary, the composition of the organic fraction determined bacterial species composition and activity in the substrate, which in turn regulated plant gene expression relative to biological control.

II. OBJECTIVES OF ORIGINAL PROPOSAL

The objectives of the original proposal were to enhance the potential for utilization of composted municipal and agricultural waste in agriculture. This was to be achieved by improving methodology for process and quality assessment, and by elaborating suppressive properties to diseases caused by plant pathogens.

The specific objects were:

- 1) To determine changes in microbial activity and total biomass with time, in compost amended soils as related to *Pythium* damping-off severity and to changes in composition of bacterial taxa in the rhizosphere of plants as suppressiveness declined.
- 2) To determine changes in *Pythium* populations and in its reproductive cycle (i.e. mycelial growth, oospore and sporangial formation and/or germination) with time in soil amended with compost to delineate effects on inoculum potential.
- 3) To characterize organic matter transformations in composts and in soil amended with compost: i) by studying transformations of lignin, cellulose and hemicellulose and humic substances using ^{13}C -NMR, NIR and FTIR spectroscopy, and crude fiber analyses; ii) by following changes in nutrient and other ion concentrations; and iii) by relating the formation of toxic and/or plant growth promoting substances with compost maturity.
- 4) To relate compost maturity and nutrient availability to microbial activity, biomass and changes in nutritional status of the biomass, composition of bacterial taxa in the rhizosphere with time of plant growth and suppression of *Pythium* damping-off (analysis of data collected in relation to objectives 1 to 4).
- 5) To define common parameters for maturity of composts produced from municipal solid waste, cattle manure and yard waste (based on results collected according to objectives 1 to 4). These parameters would eventually be used for quality control and standardization of the end-product.

III. RELEVANT DATA

A) Compost Quality:

In both Ohio and Israel, major progress was made in the field of compost

quality. In Ohio, Iannotti et al, 1994, developed a procedure based on oxygen respirometry, that accurately assessed stability of composted municipal solid waste (MSW). Furthermore, an equation predicting growth of rye grass in a compost-amended substrate, based on specific oxygen uptake rate of the compost (SOUR), was developed. This procedure greatly enhances opportunities for successful composting and for utilization of composts in agriculture. Details of the procedures and specific results are presented in the attached publication by Iannotti et al. (1994) in J. Environ. Qual. 23:1177-1183 (Appendix I).

In Israel, Chefetz et al, 1996, examined transformations of organic matter during composting of MSW to define the stability of the product, based on direct spectroscopy. Utilizing both CPMAS ^{13}C -NMR and DRIFT spectroscopy, they showed that the humic acid (HA) content of the compost increased to a maximum after 112 days, corresponding to the highest potential for plant growth (cucumber) and highest 1650/1560 ($\text{cm}^{-1}/\text{cm}^{-1}$) peak ratios calculated from DRIFT spectra. Both NMR and DRIFT spectra revealed that the residual OM contained increasing levels of aromatic structures. Their data showed that MSW compost stabilized and supported excellent plant growth after 110 days of composting. Details of the procedures and specific results of this work are presented in Chefetz et al. (1996) in J. Environ. Qual. 25:776-685 in Appendix II.

These findings correspond closely with those reported by Iannotti et al in Ohio utilizing SOUR respirometry and other analytical procedures developed by that team defining maturity/stability of MSW compost.

B) Beneficial Utilization of Compost

MSW compost prepared in Israel suppressed plant diseases caused by *Rhizoctonia solani*, *Pythium aphanidermatum* and *Fusarium oxysporum*. The effect against *R. solani* was biological in nature.

Several potential biocontrol agents were isolated from the compost, including *Pantoea agglomerans*. Lysis of hyphae of *R. solani* by bacteria was observed. In

summary, the MSW compost induced beneficial effects (Cohen et al, 1996, *Phytoparasitica* 24:145).

In Ohio, the role of soil organic matter decomposition level in sustained suppression of *Pythium* diseases was explored, utilizing both CPMAS ^{13}C -NMR and DRIFT spectroscopy. As the rate of hydrolysis of fluorescein diacetate in a peat mix decreased and *Pythium* populations increased, root rot developed, all according to procedures developed in the mid eighties in both Ohio and Israel under a previous BARD Grant (US662-83). As suppression was lost in the substrate, the concentration of ^{13}C -NMR-detectable polysaccharides (cellulosic substances) declined. At the same time, bacterial species composition in the rhizosphere as well as the soil mix shifted from those able to induce biological control (i. e. *Pseudomonas* spp.) to those that could not (i. e. oligotrophs). The concentration of microbial biomass also declined. However, microbial activity (incorporation of ^{14}C acetate into microbial biomass) did not. In fact, specific microbial activity increased. Detailed procedures and results for this work are presented in Appendix I, in Boehm et al, 1997. *AEM* 62:162-168.

This research established that lack of competition did not explain collapse in suppressiveness and that the polysaccharide fraction in OM is critical in the sustenance of biocontrol agents active in suppression of *Pythium*.

The role of OM decomposition level in the biological suppression of *Pythium* root rot was examined further in a composted separated cow manure-amended substrate by A. G. Stone (Stone et al, 1996, *Phytopathology* 86:S53, attached in Appendix I as a poster). This work presently is being prepared in Ph.D. thesis form and will be submitted to BARD as peer reviewed papers in the near future.

Stone et al investigated temporal relationships between particulate organic matter decomposition level and biological control of *Pythium* root rot caused by *Pythium ultimum* in a composted cow manure-amended sand. Suppressiveness to root rot was sustained over a 9-mon period in a 15% compost/sand mix although *P. ultimum* populations began to increase at 7-mon. As root rot developed and

progressed, the population of *P. ultimum* increased further, while microbial biomass (as determined by phospholipid phosphate) and microbial activity (as determined by FDA hydrolysis) declined. Proportions of actinomycetes declined, while proportions of Gram negative bacteria and fungal and protozoan populations increased (as determined by PLFA analysis). The quantity of particulate organic matter declined at a constant rate over time. Peaks (from DR-FTIR spectroscopy) representing plant-derived polysaccharides (hemicellulose, cellulose) and lignin declined in intensity, while peaks representing microbe-derived polysaccharides and non-lignin aromatics increased in intensity.

Since competition was not the key mechanism underlying biocontrol of Pythium root rot, and parasitism was excluded earlier as a principal factor in control of this disease in compost-amended mixes in both Ohio and Israel (BARD Grant #US 662-63), the role of compost-supported microbiota in the induction of systemic acquired resistance (SAR) against Pythium root rot was explored. It represents the fourth principal mechanism of biological control.

Zhang et al, 1996, revealed that cucumber plants produced in various compost-amended mixes suppressed Pythium root rot as well as anthracnose of cucumber foliage caused by *Colletotrichum orbiculare* through induction of SAR. It was established that plants in the suppressive compost-amended mixes supported higher levels of peroxidase activity immediately upon infection of the host. Zhang et al, 1996 concluded that interactions of the microflora in the compost with the plant and the pathogen during and early after infection were critical for rapid activation of SAR-associated gene expression in cucumber. In summary, soil OM chemistry appeared to indirectly affect plant gene expression relative to Pythium disease control. Details of procedures and results for this work are presented in Zhang et al, 1996 in *Phytopathology* 86:1066-1070 (Appendix I).

IV. DESCRIPTION OF COOPERATION

The principal investigators met for one week in 1994 during an International Symposium held at Rehovot, Israel, and again in 1995 during another International Composting Symposium in Bologna, Italy. Four review papers (Chen et al, 1996; Chen et al, 1996; Chefetz et al, 1996 and Hoitink et al, 1996), were published in which details of this research program were discussed relative to other literature on compost quality. Copies of all publications are enclosed in the report.

Graduate students and post-doctoral scientists active in this project in Ohio and Israel exchanged procedures routinely via E-mail as illustrated by the work on NMR and DRIFT spectroscopy by Boehm et al, 1997; Chefetz et al, 1996 and Stone et al, 1997 in preparation.

V. RESEARCH ACHIEVEMENTS/BENEFITS TO AGRICULTURE

Research supported through this proposal is having a major impact on 1) waste recycling through composting and 2) how diseases caused by soilborne plant pathogens are controlled in agriculture today. In August, 1995, the IPM Practitioner published a major article entitled "Alternatives to Methyl Bromide in Nurseries - Disease Suppressive Media (W. Quarles and J. Grossman, IPM Practitioner, Vol. 17(8) 1-13). This paper reviews the status of the field of composting and present practices in biological control of plant diseases with composts. It concludes that compost-amended mixes not only can serve as a substitute for methyl bromide fumigation but that fungicide applications can be reduced significantly as well and in some cases eliminated. This paper cites 85 references. Sixty percent of the citations dealing with composting (37 of 62 papers) were published by the Ohio and/or Israeli teams. This article illustrates how disease suppressive composts have become a generally accepted phenomenon in the US and elsewhere and that much of the science on which it is based was developed by this team.

The research by Chefetz (Israel) and Stone and Boehm (Ohio) on DRIFT and NMR spectroscopy have given credibility to the earlier work on FDA activity by Mandelbaum and Hadar, 1990 (Israel) and Chen et al, 1988 (Ohio; W. Chen not to be confused with Y. Chen from Israel). FDA activity seems to mimic the CPMAS ^{13}C NMR - detectable concentration of cellulose in substrates which is critical to biocontrol efficacy (Boehm et al, 1997). Even though these laboratory assays (FDA activity, NMR and DRIFT spectroscopy) predict the potential for biological control of Pythium root rot, practical tests useful in field agriculture still remain to be developed.

In our original objectives we planned to evaluate NIR spectroscopy as a low cost procedure for prediction of this beneficial effect in composts. Efforts by A. G. Stone in Ohio to utilize NIR for this application were hampered by slow progress in her work on DRIFT spectroscopy on OM recovered from substrates as suppressiveness declined (Ph.D. thesis in preparation). All other objectives were completed successfully.

VI. LIST OF PUBLICATIONS FROM PROJECT

1. Boehm, M. J., T. Wu, A. G. Stone, B. Kraakman, D. A. Iannotti, G. E. Wilson, L. V. Madden, and H. A. J. Hoitink. 1997. Cross-Polarized Magic-Angle Spinning ^{13}C Nuclear Magnetic Resonance Spectroscopic Characterization of Soil Organic Matter Relative to Culturable Bacterial Species Composition and Sustained Biological Control of Pythium Root Rot. *Appl. and Environ. Microbiol.* 1997. 162-168 pg.
2. Chen, Y., Y. Inbar, B. Chefetz, and Y. Hadar. 1996. Composting and Recycling of Organic Wastes. Pg. 339-360. In: *Modern Agriculture and Environment*. D. Rosen, Y. Chen, E. Tel-Or and Y. Hadar (eds.). Kluwer Academic Publishers, London U. K. (in press).
3. Chen, Y., B. Chefetz, and Y. Hadar. 1996. Formation and properties of humic substances originating from composts. Pg. 382-393. In: *The Science of Composting*, M. de Bertoldi, P. Sequi, B. Lemmes, and T. Papi (eds.). Blackie Academic & Professional, Bologna Italy.
4. Chefetz, B., Y. Chen, and Y. Hadar. 1996. Municipal solid waste composting:

Chemical and biological analysis of the process. Pg. 1105-1108. In: The Science of Composting. M. de Bertoldi, P. Sequi, B. Lemmes, and T. Papi (eds.). Blackie Academic & Professional, Bologna Italy.

5. Chefetz, B., P. G. Hatcher, Y. Hadar, and Y. Chen. 1996. Chemical and biological characterization of organic matter during composting of municipal solid waste. *Journal of Environmental Quality* 25:776-785.
6. Cohen, R. 1996. Suppression of soil-borne pathogens by composted municipal solid waste. M. Sc. Thesis-in Hebrew.
7. Bar-Lev, N. 1996. Use of municipal solid waste compost to suppress soil borne diseases. M. Sc. Thesis-in Hebrew.
8. Cohen, R., N. Bar-Lev, B. Chefetz, Y. Chen, and Y. Hadar. 1996. Suppression of soil borne pathogens by composted municipal solid waste. *Phytoparasitica* 24:145-145.
9. Cohen, R., and Y. Hadar. Utilization composted municipal solid waste for biocontrol of soil borne pathogen diseases. (in preparation)
10. Hoitink, H. A. J., M. E. Grebus, and A. G. Stone. 1997. Impacts of compost quality on plant disease severity. Pg. 361-369. In: *Modern Agriculture and the Environment*. D. Rosen (ed.). Kluwer Academic Publishers, Great Britain.
11. Iannotti, D. A., M. E. Grebus, B. L. Toth, L. V. Madden, and H. A. J. Hoitink. 1994. Oxygen Respirometry to Assess Stability and Maturity of composted Municipal Solid Waste. *J. Environ. Qual.* 23:1177-1183.
12. Zhang, W., W. A. Dick, and H. A. J. Hoitink. 1996. Compost-Induced Systemic Acquired Resistance in Cucumber to Pythium Root Rot and Anthracnose. *Phytopath.* 86:1066-1070.

VII. PATENTS

None

VIII. APPENDIX I. - RESEARCH PERFORMED IN OHIO

Principal Investigators

H. A. J. Hoitink

L. V. Madden

Research Associates

D. A. Iannotti, Ph.D.

A. G. Stone, Ph.D. candidate

W. Zhang, Ph.D. candidate

Publications

Oxygen Respirometry to Assess Stability and Maturity of Composted Municipal Solid Waste. D.A. Iannotti, M.E. Grebus, B.L. Toth, L.V. Madden and H.A.J. Hoitink. 1994. Journal of Environmental Quality 23:1177-1183.

Cross-Polarized Magic-Angle Spinning ^{13}C Nuclear Magnetic Resonance Spectroscopic Characterization of Soil Organic Matter Relative to Culturable Bacterial Species Composition and Sustained Biological Control of Pythium Root Rot. M.J. Boehm, T. Wu, A.G. Stone, B. Kraakman, D.A. Iannotti, G.E. Wilson, L.V. Madden, and H.A.J. Hoitink. 1997. Applied and Environmental Microbiology 63:162-168.

Compost-Induced Systemic Acquired Resistance in Cucumber to Pythium Root Rot and Anthracnose. W. Zhang, W.A. Dick, and H.A.J. Hoitink. 1996. Phytopathology 86:1066-1070.

Impacts of Compost Quality on Plant Disease Severity. H.A.J. Hoitink, M.E. Grebus and A.G. Stone. 1996. Modern Agriculture and the Environment, pp. 361-369.

Dynamics of Pythium Populations, Microbial Biomass, Microbial Activity, and Organic Matter Characterization During the Collapse of Suppression of Pythium Root Rot. A.G. Stone, D.Y. Han, M.M. Wander, S.J. Traina, and H.A.J. Hoitink. 1996. Phytopathology 86:S53.

Oxygen Respirometry to Assess Stability and Maturity of Composted Municipal Solid Waste

D. A. Iannotti, M. E. Grebus, B. L. Toth, L. V. Madden, and H. A. J. Hoitink*

ABSTRACT

The stability and maturity of compost prepared from municipal solid waste (MSW) at a full-scale composting plant was assessed through chemical, physical, and biological assays. Respiration bioassays used to determine stability (O_2 and CO_2 respirometry) were sensitive to process control problems at the composting plant and indicated increasing stability with time. Radish (*Raphanus sativus* L.) and ryegrass (*Lolium perenne* L.) growth bioassays revealed that immature compost samples inhibited growth. Growth of ryegrass in potting mix prepared with cured compost not amended with fertilizer was enhanced as compared to a peat control. Garden cress (*Lepidium sativum* L.) seed germination, used as an indicator of phytotoxicity, revealed inhibition of germination at all compost maturity levels. The phytotoxicity was thought to be salt-related. Spearman rank-order correlations demonstrated that O_2 respirometry, water-soluble organic C, and the water extract organic C to organic N ratio, significantly correlated with compost age and best indicated an acceptable level of stability. Oxygen respirometry also best predicted the potential for ryegrass growth, and an acceptable level of compost maturity.

SUCCESSFUL MARKETING of composts requires that the potential for plant growth is predictable after incorporation of these organic amendments into soil. Many tests have been proposed to assess this aspect of compost quality. Changes in compost stability, or the degree to which composts have been decomposed, can be predicted with respirometry (reviewed in Iannotti et al., 1993), heat output (Jourdan, 1988), by following concentrations of soluble organic C in the water extract of compost, particularly that of readily available organic nutrients (Garcia et al., 1991a), and finally by NMR and IR spectroscopy (Chen and Inbar, 1993). Changes in the percent volatile solids or ash content with time within specific batches of composts also can provide useful information if the necessary precautions are taken. Stability by itself, however, does not necessarily predict the potential for plant growth (Chen and Inbar, 1993).

An acceptable level of compost maturity, which generally implies the potential for the development of beneficial effects upon utilization, is best determined with plant growth bioassays (Chen and Inbar, 1993). Effects of composts on plant health (disease suppression) and soil physical properties (i.e., tilth) are not considered here, although the microflora and organic matter in mature composts critically affects these compost quality parameters (Hoitink et al., 1993; Dick and McCoy, 1993).

The objectives of this work were to apply a battery of procedures previously described for the assessment of compost stability and maturity to municipal solid waste compost (MSW) and to determine which methods best indicate an acceptable level of maturity required for successful utilization of this compost.

METHODS AND MATERIALS

Composting and Curing Process

Raw municipal solid waste, largely devoid of yard trimmings and food wastes, delivered to the full-scale composting facility in St. Cloud, MN, was sorted on the tip floor for recyclable items. The sorted waste was loaded into a drum homogenizer rotating at 0.5 rpm with a residence time of 66 h. The homogenized MSW was conveyed into a trench composting system (Royer Industries, Kingston, PA) equipped with temperature feedback control and forced aeration. The material was turned 13 times, with a total residence time of 52 d. Water was added during the procedure to maintain a moisture content of 50% (w/w) immediately after the compost was turned. After removal from the trench, the compost was screened (25 mm or 1 inch) and transported to The Ohio State University (OSU) laboratory at Wooster, where it was cured in a pile under a pole barn for 112 d. The pile was turned monthly and water was added after the first and second turnings to maintain a moisture content of 50% (w/w). It was allowed to dry thereafter. Temperatures of the compost while in the trench remained above 43°C and below 70°C. Temperatures during curing measured in the center of the pile ranged from 65°C during early curing (Day 72) to 41°C by Day 164.

Sampling Procedures

The compost was sampled immediately after its removal from the drum homogenizer, at five different times during processing in the trench system and, finally, three times during curing. Samples (three replicates/sampling date) were collected within 6 h after the compost was turned. They were taken from just below the surface layer in the center of the trench, and at 25-cm distances from each of the side walls. This ensured maximum homogeneity among samples (Kuter et al., 1985). Samples (4 L) were placed in partially closed polyethylene bags, packed on ice and then shipped overnight for analysis. The samples arrived at OSU with a mean temperature of 4°C. During curing, compost was sampled immediately after turning at monthly intervals (three 4 L replicates per harvest date) and analyzed within 1 h after being collected.

Chemical and Physical Analyses

Samples were screened through a 9.5-mm (0.38 inch) screen to remove large particles. The number of subsamples ranged from three to five depending upon the sensitivity of a particular

Department of Plant Pathology, The Ohio State University/Ohio Agricultural Research and Development Center, Wooster, OH 44691. Present address of D. A. Iannotti, Division of Natural Sciences and Mathematics, St. Mary's College of Maryland, St. Mary's City, MD. Received 12 Aug. 1993. *Corresponding author (hoitink.1@osu.edu).

assay. The number was adequate to obtain means with coefficients of variation less than 30%. Assays on solids were performed on five subsamples for each of the three samples retrieved at each harvest date. Dry solids content (% w/w) was determined after microwave oven drying to a constant weight. The pH of each sample was determined on a saturated paste extract. Ash content, total N (%) and total C (%) were determined at the Research Extension Analytical Laboratory (REAL) at OSU, Wooster. Ash content was determined by heating (550°C for 4 h) in a Thermolyne Furnace (model 30400, Thermolyne Corp., Dubuque, IA). Volatile solids were determined by subtracting ash content from dry solids. Total organic matter was the fraction destroyed by ashing. Total N analysis was performed by oxidative digestion using the Dumas method (Macro N analyzer Foss/Heraeus/UIC, Joliet, IL). The detection limit for this instrument is 200 mg/kg. Total C was determined using CO₂ coulometry (model 5020, Coulometrics, UIC, Joliet, IL). This instrument converts C in the sample to CO₂ by oxidation at 1100°C. The detection limit is <1 mg kg⁻¹.

Compost water extracts (three subsamples for each of the three samples per harvest date) were prepared by the saturated paste method and acidified immediately to avoid NH₃ loss (Brown and Warnke, 1988). Extracts were then frozen and analyzed at REAL for total C, Kjeldahl N as an estimate of organic N, inorganic C, NO₃⁻-N, NH₄⁺-N and electrical conductivity.

Total Kjeldahl N in the water extract was determined by a macro Kjeldahl method. Nitrate-N and NH₄⁺-N were determined by a nitrate electrode and distillation-titration methods, respectively (APHA, 1992). Total water soluble C and water soluble inorganic C determinations were made using C oxidation by coulometry and a coulometry instrumentation modification utilizing a model 5130 acidification module (UIC, Joliet, IL) for inorganic C. Water soluble organic C represents the difference between total water soluble C and water soluble inorganic C. Electrical conductivity was determined with a Solu Bridge Conductivity Indicator (Beckman Instruments, Cedar Grove, NJ).

Compost Stability

The rate of oxygen uptake was determined with dissolved oxygen respirometry (Iannotti et al., 1993). Screened compost samples with a moisture content of approximately 50% (w/w) were placed in 500-mL Erlenmeyer flasks and aerated for 16 h at 37°C. In this procedure, water is added to samples low in moisture content (<45%, w/w) to ensure that matric potential does not limit microbial activity. Excessively wet samples are air-dried down to 50% moisture (w/w). A dissolved oxygen meter and probe assembly were used to determine the change in percent oxygen in air over a 1-h period. The change in % oxygen in air was converted to a rate of oxygen uptake based on volatile solids (VS) by using the equation:

$$\text{Mean O}_2 \text{ uptake} = \frac{-CVS 60 D}{K W_t VS}$$

where O₂ uptake is oxygen uptake expressed as g O₂ per kg of volatile solids per hour (SOUR; g kg⁻¹ VS h). The other variables were:

- C = oxygen content by volume in the air; usually 0.21
- V = volume of air in flask (mL)
- S = slope of relative O₂ uptake rate (% saturation per minute)
- 60 = factor change from minutes to hours
- K = constant factor based on the calibration value corrected

for elevation above sea level, (e.g., 100 at sea level, 98 at 170 m [558 ft], 90 at 880 m [2887 ft], obtained from D.O. meter manufacturer's recommended calibration procedure)

W_t = compost dry matter weight in flask (gram)

D = the density of the oxygen (g L⁻¹) at experimental conditions was converted from STP (Standard Temperature and Pressure, 1 atm pressure, 273.15 K, O₂ density = 1.429 g L⁻¹). The experimental conditions were 310.15 K (37°C) at 100% water vapor saturation in air, O₂ density = 1.1816

VS = the fraction of volatile solids (from 0 to 1.0)

The rate of respiration in compost samples, based on CO₂ evolution, was measured using an adaptation of the procedure of Bartha and Pramer (1965). Screened compost samples were sealed in a 3.8-L vessel with a 1.0 M NaOH solution that absorbed CO₂. After 24, 48, and 72 h of incubation at 25°C, the absorbed CO₂ was precipitated with barium chloride as barium carbonate (Anderson, 1982). The remaining base was titrated with 0.5 M HCl. The quantity of acid required for neutralization was used to calculate CO₂-C evolved per g dry weight of compost per 24 h. The total amount of CO₂ evolved during a 3-d period was then determined.

Plant Growth Bioassays

The aboveground dry matter of cucumber (*Cucumis sativus* L.), radish, and ryegrass produced in compost-amended mixes was compared with that of plants produced in a Canadian Sphagnum peat potting mix. Mixes were prepared by blending compost samples collected at various dates (degrees of maturity) with coarse horticultural grade perlite and Canadian sphagnum peat moss (15:30:55, v/v) in a concrete mixer. Dolomitic lime, gypsum, treble super phosphate and potassium nitrate were added in quantities based on potting mix analyses prepared at REAL to adjust the pH to 5.5 and essential plant nutrients to within the optimum range for potting mixes (Inbar et al., 1993). The mixes were blended 3 min, and water was added to adjust the final moisture content to 50% (w/w). The peat mix used as a control contained peat and perlite (70:30, v/v) amended as described above for the compost mixes.

Cucumber growth bioassays were performed in compost-amended potting mixes amended with 12.5 g L⁻¹ slow release fertilizer (Osmocote 14-14-14, Grace-Sierra Chemical Co., Milpitas, CA). Eight cucumber (cv. Straight Eight, 90% germination) seeds were planted 1 cm deep in each of five pots. After 10 d, the seedlings were thinned to four per pot. The pots were irrigated daily and incubated 21 d in a greenhouse at 20 to 30°C supplemented with 14-h illumination per day (225 μE m⁻² s⁻¹). The aerial portion of the plants was harvested and dried to a constant weight. A mean dry weight per pot was calculated.

Thirty-two radish seeds were planted per pot using a pneumatic seeder. The pots (five per treatment) were incubated 7 d in a growth chamber at a constant temperature of 25°C with continuous illumination (225 μE m⁻² s⁻¹) and irrigated daily. After 7 d, the aerial portion of the radish plants harvested in each pot was dried to a constant weight. Mean dry weights of emerged plants per pot were calculated.

To determine the effect of compost maturity on growth of ryegrass and fertility, a second replicate batch of potting mix was prepared, with and without added potassium nitrate and slow-release fertilizer (Osmocote 14-14-14 as described above). Each pot was seeded with 1 g ryegrass seed spread evenly over the surface of the mix. The pots were incubated in a greenhouse at 23 to 28°C for 21 d under illumination as described above and irrigated by misting twice each day. The

aerial plant material in each pot was harvested after 21 d and dried to a constant weight. The mean dry weight of plant material per pot was calculated.

Cress Phytotoxicity Bioassay

Cress phytotoxicity bioassays were performed on water extracts prepared from compost according to Zucconi et al. (1981a,b). Extracts were centrifuged 15 min at $8000 \times g$ at 5°C and then filter-sterilized through a $0.45\text{-}\mu\text{m}$ membrane (Gelman Scientific, Ann Arbor, MI). A dilution series was prepared (undiluted, threefold, and 10-fold) from each extract, maintaining aseptic conditions. One milliliter of each test solution was pipetted into a sterilized 50-mm glass petri dish lined with filter paper (Whatman no. 1; 4.25-cm diam.). Five replicates were used per dilution. Sterilized distilled water was used as a control. Eight cress seeds (no. 2824, fine curled, Liberty Seed Co., New Philadelphia, OH) were placed in each dish and incubated 24 h at 27°C in the dark. Germination was then stopped by adding 1 mL 50% (v/v) ethanol to each dish. Treatments were evaluated by noting the number of germinated seeds and the length of the root radicle. Ungerminated seeds were defined as having radicles 0 cm long.

The percent germination was calculated by dividing the mean germination of each treatment by the mean germination in the sterile water control and multiplying this number by 100. The percent root length was calculated by dividing the mean root length of each treatment by the mean root length of the water control and multiplying this number by 100.

The germination index was then calculated by multiplying the percent germination by the percent root length.

Experimental Designs and Statistical Analyses

Completely randomized designs were used in all plant bioassays. One-way analyses of variance (ANOVA) were performed with Minitab statistical software (Minitab, State College, PA). Separation of means was based on the least significant difference (LSD, $P \leq 0.05$). Spearman rank-order correlation analysis was used to determine correlation coefficients (r_s) between time and variables characterizing the compost. Regression analysis was also used to develop a model between maturity of compost, based on plant growth and compost characteristics.

RESULTS AND DISCUSSION

Temporal changes in pH, electrical conductivity, volatile solids, and total C/N (Fig. 1) were typical of those reported by others for MSW (Hirai et al., 1983; Saviozzi et al., 1992; Garcia et al., 1992a). The electrical conductivity was high throughout ($>10\text{ dS m}^{-1}$), which also is not atypical for MSW composts (Jourdan, 1988; Iglesias Jimenez and Perez Garcia, 1989).

The volatile solids content of the mature compost after 164 d (Fig. 1C) was 18.7%. The organic matter content was low, therefore. Since small pieces of inert plastic

Fig. 1. Changes in chemical characteristics of MSW compost as a function of composting time. Bars represent standard errors ($n = 3$).

Fig. 2. Mean relative oxygen level (%), based on O_2 respirometry, for MSW compost from a full-scale plant as a function of composting time. Bars represent standard errors ($n = 3$).

were readily apparent in these mature compost samples, the residual concentration of biodegradable organic matter actually was less than 18.7% (w/w).

Compost stability, as determined by O_2 and CO_2 respirometry (Fig. 2 and Table 1), revealed a significant decrease (>11-fold) in rate of respiration with compost age. Temporal changes in % volatile solids during composting of the MSW used in this work (Fig. 1C) revealed that most of the easily biodegradable substances were destroyed during the first 20 d in the trench composting system. Rates of respiration, expressed on the basis of % VS (Table 2), were well within the range of values reported for other MSW composts (Haug, 1980; Keener et al., 1993).

Rates of aeration applied early during the process in this trench MSW composting system were too low to prevent significant levels of fermentation. The organic acids produced during this period lowered the pH (Fig. 1A), particularly because the buffer capacity in fresh compost is low (Chen and Inbar, 1993). The sharp fluctuations in electrical conductivity and high standard errors for this parameter early during the process also suggest that sudden changes in activity of microbial populations and lack of process control prevailed at that time. This may explain why compost samples collected on Day 18 did not follow the decline in rate of respiration observed overall (Fig. 2 and Table 1). These samples emitted strong odors, typical of fermentation, upon their arrival at OSU. Composting system fans operated only

part of the time during the first 18 d. Low activity at this time undoubtedly was due to inhibition of the compost microflora caused by low pH (Cappaert et al., 1976; Kubota and Nakasaki, 1991).

The high rate of respiration observed in the Day 45 sample with O_2 relative to CO_2 respirometry (Fig. 2, Tables 1 and 2) may have been due to the low moisture content of the compost (40%, w/w) when it was collected from the trench system (Miller, 1989). Water added to these dry compost samples to raise the moisture content to 50% (w/w) as required for both respirometry assays (Bartha and Pramer, 1965; Iannotti et al., 1993), initiates a sudden increase in bacterial growth and activity. This yields a short-term false instability reading (Miller, 1989; Inbar et al., 1993). This short burst of activity would not be as readily detectable with the CO_2 procedure because of the long (24 h) incubation period used in that test as compared with the short time (1 h) in the dissolved oxygen meter procedure.

The concentration of soluble organic C declined with composting time after Day 10 (Table 3). The greatest decline occurred during the first approximately 80 d of the process. This decline in water soluble organic C often is used as an indicator of compost maturity (Saviozzi et al., 1992; Inbar et al., 1993; Garcia et al., 1991a,b). Organic C in the water extract of immature MSW compost consists of sugars, phenolic substances, organic and amino acids, peptides, and other easily biodegradable substances. In mature compost, most of the soluble organic C is present as humic substances which are resistant to further decomposition and this explains the increased stability observed with time during composting (Garcia et al., 1991a; Chen and Inbar, 1993).

The slight decrease in ammonium N concentrations with composting time after Day 10 observed in this work (Table 3) agrees with previous reports for MSW (Chanyasak and Kubota, 1981; Garcia et al., 1991b). The general increase in nitrate concentration (Table 3) also is typical of previous reports. The low NO_3^- -N value detected in the Day 164 sample probably was due to denitrification rather than N immobilization. An excessive amount of water inadvertently had been added during turning on Day 116 to this low organic matter compost. This induced temporary anaerobic conditions, as evidenced by fermentation odors emitted for several weeks after the water was added, and probable denitrification during curing. Immobilization of N probably was

Table 1. Changes in rates of respiration based on O_2 uptake and CO_2 evolution per kg dry solids (DS) during composting of MSW.

Compost age	Mean rate of CO_2 evolution				Mean O_2 uptake
	0-24 h	24-48 h	48-72 h	Total/72 h	
	g CO_2 -C kg ⁻¹ DS				g O_2 kg ⁻¹ DS h ⁻¹
d					
3	6.74 ± 2.1†	5.72 ± 0.48	3.11 ± 0.20	15.6 ± 2.8	1.24 ± 0.03
10	3.11 ± 0.20	4.35 ± 2.5	4.50 ± 1.1	12.0 ± 3.8	1.61 ± 0.17
18	0.11 ± 0.11	0.18 ± 0.12	0.44 ± 0.28	0.73 ± 0.51	0.44 ± 0.09
31	1.09 ± 0.50	1.89 ± 0.54	2.12 ± 0.76	5.10 ± 1.8	0.69 ± 0.14
45	1.55 ± 1.0	1.99 ± 0.39	2.67 ± 0.25	6.21 ± 1.6	1.14 ± 0.01
52	0.59 ± 0.11	1.37 ± 0.10	1.76 ± 0.24	3.72 ± 0.45	0.57 ± 0.05
81	1.54 ± 0.13	1.87 ± 0.19	1.89 ± 0.12	5.30 ± 0.44	0.48 ± 0.04
116	0.63 ± 0.15	0.91 ± 0.19	0.71 ± 0.17	2.25 ± 0.51	0.44 ± 0.06
164	0.63 ± 0.06	0.38 ± 0.10	0.33 ± 0.05	1.34 ± 0.21	0.08 ± 0.01

† Represents standard error based on three replicates.

Table 2. Changes in rates of respiration based on O₂ uptake and CO₂ evolution per g volatile solids (VS) during composting of MSW.

Compost age	Mean rate of CO ₂ evolution				Mean O ₂ uptake
	0-24 h	24-48 h	48-72 h	Total/72 h	
d	g CO ₂ -C kg ⁻¹ VS				g O ₂ kg ⁻¹ VS h ⁻¹
3	14.4 ± 2.5†	12.3 ± 0.77	6.66 ± 0.35	33.4 ± 3.6	2.67 ± 0.13
10	4.91 ± 1.7	8.10 ± 3.0	8.32 ± 1.4	21.3 ± 6.1	2.90 ± 0.33
18	0.19 ± 0.09	0.60 ± 0.28	1.49 ± 0.35	2.28 ± 0.72	1.63 ± 0.48
31	4.31 ± 0.88	7.78 ± 0.98	8.48 ± 0.88	20.6 ± 2.7	2.74 ± 0.41
45	2.88 ± 0.55	6.01 ± 0.65	8.07 ± 0.34	17.0 ± 1.5	3.46 ± 0.11
52	2.08 ± 0.20	4.89 ± 0.18	6.25 ± 0.53	13.2 ± 0.91	2.04 ± 0.22
81	7.24 ± 2.14	8.87 ± 0.90	8.92 ± 0.72	25.0 ± 3.8	2.28 ± 0.28
116	2.88 ± 0.23	4.03 ± 0.47	3.18 ± 0.43	10.1 ± 1.1	1.94 ± 0.28
164	3.63 ± 0.10	2.19 ± 0.38	1.91 ± 0.14	7.73 ± 0.62	0.48 ± 0.07

† Represents standard error based on three replicates.

not a factor because (i) the C/N ratio of the compost at this time was low (12.5, Fig. 1D) and (ii) added fertilizer did not affect ryegrass growth in the 164-d compost treatment (Table 4).

The organic C to water soluble organic N ratio in MSW compost water extracts has been proposed as a maturity assay that predicts the potential for plant growth in MSW compost-amended substrates (Chanyasak and Kubota, 1981; Chanyasak et al., 1983). This predictive procedure later was extended to include composts prepared from sewage sludges and other wastes (Hirai et al., 1983). Values ranging from 5 to 6 reflect maturity. Garcia et al. (1991b) found similar trends for sewage sludge composts. In our work with MSW compost, values were high (12-15) initially and declined to 6.5 by Day 164, which is near that reported for mature composts.

Seed germination assays often are used to test composts for maturity. Cress, because of its small seed size and high rate of germination (25 h at 25°C) and sensitivity to environmental stresses, is used most frequently (Zucconi et al., 1981a,b). The 10-fold diluted compost water extract used in this work with MSW yielded a watercress germination index of only 60%, even after 164 d (Fig. 3). The undiluted extract was inhibitory at all maturity levels. The low organic matter content and high salinity in this compost, which results from a high degree of waste separation (Richard et al., 1993), undoubtedly contributed to the inhibition of germination observed in this work.

Several reports reveal that fermentation by-products, the principal phytotoxic substances in immature MSW composts, are destroyed as composts mature (De-

Vleeschauwer et al., 1981; Chanyasak et al., 1983; Garcia et al., 1992b). Other inhibitors present in plant residues affecting plant growth are allelopathy toxins. The latter also are readily destroyed during composting (Solbraa et al., 1983; Still et al., 1976). Thus, lack of maturity likely was not the factor limiting cress seed germination in the 164 d-old compost. Organic acids produced as a result of fermentation as well as N deficiency in immature compost and salinity at all maturity levels, probably were the key factors affecting germination of cress seed.

Plant growth bioassays (Table 4) revealed that ryegrass, cucumber, and radish differed in their responses to MSW compost treatment. Growth of ryegrass in the mature (Day 164) compost-amended mix was significantly ($P = 0.05$) better than that in the peat mix, independent of fertilization (Table 4). The yield in the Day 164 mix not treated with slow-release fertilizer was more than double that in the unfertilized peat control. On the other hand, growth of radish was significantly ($P = 0.05$) less than that in the fertilized peat control. Growth of cucumber was not affected by maturity and was not significantly different from that in the peat mix

Table 4. Effect of MSW compost maturity on growth of ryegrass, cucumber and radish.

	Without fertilizer	Fertilizer added†		
	Ryegrass‡	Ryegrass‡	Cucumber§	Radish¶
Day 3	119 ± 7#	197 ± 4#	97 ± 9#	26 ± 2#
Day 10	101 ± 6	190 ± 7	100 ± 7	15 ± 1
Day 18	112 ± 8	193 ± 5	85 ± 4	19 ± 1
Day 31	105 ± 11	194 ± 2	97 ± 17	22 ± 3
Day 45	112 ± 3	197 ± 9	124 ± 13	19 ± 2
Day 52	128 ± 6	219 ± 12	140 ± 12	24 ± 1
Day 81	114 ± 7	199 ± 15	117 ± 7	33 ± 1
Day 116	154 ± 31	247 ± 3	98 ± 27	24 ± 3
Day 164	210 ± 20	241 ± 7	116 ± 29	28 ± 2
LSD(0.05)	26	25	48	6
Peat mix††	92 ± 4	212 ± 13	120 ± 21	45 ± 1

† Osmocote 14-14-14, slow-release fertilizer (Grace-Sierra Chemical Co., Milpitas, CA) added at a rate of 12.5 g L⁻¹ potting mix consisting of Canadian sphagnum peat moss, perlite and compost (55:30:15;v/v).

‡ Based on five replicates seeded with 1 g ryegrass seed per pot.

§ Based on five replicates of eight cucumber seeds per pot determined 16 d after planting. Potting mix same as for ryegrass.

¶ Based on five replicates of 32 radish seeds per pot determined 10 d after planting. Potting mix same as for ryegrass.

Values are dry weights (mg) of plant tissue and standard errors ($n = 5$).
†† Peat control mix consisted of Canadian sphagnum peat moss and perlite (70:30;v/v).

Table 3. Chemical analyses of compost water extracts prepared at various times during composting of MSW.

Compost age	NO ₃ ⁻ -N	NH ₄ ⁺ -N	TKN†	Total C
d				
3	83.4 ± 12.7‡	267 ± 50	1267 ± 134	8588 ± 1664
10	88.3 ± 13.9	500 ± 58	1600 ± 204	14656 ± 676
18	84.4 ± 18.6	389 ± 77	1178 ± 1929	11528 ± 1642
31	76.3 ± 9.6	233 ± 33	1000 ± 139	7474 ± 1348
45	117 ± 8.4	422 ± 48	1044 ± 116	8002 ± 1195
52	267 ± 52.2	356 ± 73	1089 ± 128	6482 ± 1049
81	109 ± 9.1	<200‡	367 ± 33	2105 ± 152
116	259 ± 43.8	<200‡	567 ± 33	3104 ± 315
164	30.1 ± 6.6	<200‡	200‡	1361 ± 151

† Total Kjeldahl N; values in columns 2-5 represent mg L⁻¹.

‡ Represents standard error based on 15 replicates for each value.

§ Results at or below detection limit of 200 mg L⁻¹ for method.

Fig. 3. Cross germination index as a function of composting time. Seeds were incubated 24 h at 25°C in two dilutions of MSW compost water extract.

(Table 4). The foregoing bioassays reveal that ryegrass and radish were sensitive to changes in compost maturity, but cucumber was not.

Spearman's rank-order correlation coefficients (Table 5) revealed that O_2 and CO_2 respirometry, electrical conductivity, water soluble organic C and organic C/organic N in water extracts and, finally, ryegrass and radish growth all were significantly ($P < 0.01$) correlated with composting time.

Growth of ryegrass (without fertilizer) was chosen as the preferred measure of compost maturity because of its sensitivity to changes in maturity of the MSW compost used in this work. Growth measurements (ryegrass without fertilizer) were divided by the mean growth value for ryegrass in the peat mix (plus fertilizer), so that a relative scale for maturity (M) could be used. Mean rate of O_2 uptake per kg volatile solids per hour (SOUR) had the highest correlation with M . The developed regression equation was $M = 0.80 - 0.60 \log(\text{SOUR})$, which had a coefficient of determination (R^2) of 0.90 (Fig. 4).

CONCLUSIONS

Oxygen and CO_2 respirometry and soluble organic C as well as the organic C to organic N ratio in the compost

Table 5. Spearman's rank-order correlation coefficients (r_s) between composting time and various physical, chemical, and biological properties of MSW compost.

Technique	r_s
O_2 respirometry	-0.80**
CO_2 absorption respirometry	-0.90**
Ryegrass growth (dry wt.) + fertilizer	0.71**
Ryegrass growth (dry wt.) - fertilizer	0.67**
Radish growth (dry wt.)	0.69**
Cucumber growth (dry wt.)	0.37*
Ash	0.76**
Total C (solid sample)	-0.67**
Total C/N (solid sample)	-0.80**
Electrical conductivity (water extract)	-0.88**
Water extract NO_3^- -N	0.67**
Water extract organic N	-0.74**
Water extract organic C	-0.84**
Water extract organic C/N	-0.85**

*, ** Significance at the 0.05 and 0.01 levels, respectively.

Fig. 4. Relationship between the rate of respiration of MSW compost and growth of ryegrass. SOUR = mean rate of O_2 uptake (g O_2 per kg dry weight volatile solids per hour). M = mean dry weight (mg) of ryegrass per pot produced in MSW compost without fertilizer divided by that produced in a fertilized peat mix.

water extract, all were good indicators of MSW compost stability. Oxygen respirometry also was highly correlated with both stability (based on volatile solids content) and composting time. Regression analyses revealed that the highest coefficient of determination was associated with O_2 respirometry and ryegrass growth. Based on the potential for ryegrass growth, O_2 respirometry could be used to predict acceptable levels of both stability and maturity of the MSW compost used in this work.

Municipal solid waste composts often are prepared from highly heterogeneous, biodegradable raw materials that may have a variety of effects on compost quality. We recommend, therefore, that the results obtained in this work, using MSW compost prepared from raw feed stock in which paper was the principal biodegradable product, be tested for additional MSW composts.

ACKNOWLEDGMENTS

We thank Recomp Inc. of St. Cloud, MN, for supplying the MSW compost. We acknowledge Dr. M.E. Watson, head, Research Extension Analytical Laboratory; B.L. Bishop, senior statistician; and C.A. Musselman, technical assistant, Department of Plant Pathology, Ohio Agricultural Research and Development Center, The Ohio State University, Wooster, OH, for their suggestions and technical assistance. We thank Dr. M.A. Cole, University of Illinois; Dr. B.N. Johnson and Dr. G.W. Hyatt, The Procter & Gamble Co., Cincinnati, OH; and Dr. W.H. Thompson, Univ. of Minnesota, St. Paul, for reviewing the manuscript. Salaries and research support provided by state and federal funds appropriated to the Ohio Agricultural Research and Development Center, The Ohio State University, and by a grant from the Composting Council, Washington, DC, and Grant no. US-2196-22 from BARD, the United States-Israel Binational Research and Development Fund. Manuscript no. 129-93.

REFERENCES

- American Public Health Association, American Water Works Association, and Water Environment Federation. 1992. Standard methods for the examination of water and wastewater. 18th ed. APHA, Washington, DC.
- Anderson, J.P.E. 1982. Soil respiration. p. 831-871. In A.L. Page

- et al. (ed.) Methods of soil analysis. Part 2. 2nd ed. ASA and SSSA, Madison, WI.
- Bartha, R., and D. Pramer. 1965. Features of a flask and method for measuring the persistence and biological effects of pesticides in the soil. *Soil Sci.* 100:68-70.
- Brown, J.R., and D. Warnke. 1988. Recommended cation tests and measures of cation exchange capacity. Recommended chemical soil test procedures for the North Central Region. NCR Publ. 221 (revised). Bull. 499 (revised). p. 15-16. North Dakota Agric. Exp. Stn., North Dakota State Univ., Fargo.
- Cappaert I., O. Verdonck, and M. De Boodt. 1976. Composting of bark from pulp mills and the use of bark compost as a substrate for plant growth. Part 2. The effect of physical parameters on the composting rate of bark. Growth experiments with bark compost. *Compost Sci.* 17:18-20.
- Chanyasak, V., A. Katayama, M.F. Hirai, S. Mori, and H. Kubota. 1983. Effects of compost maturity on growth of komatsuna (*Brassica rapa*, var. *pervidis*) in Neubauer's pot: II. Growth inhibitory factors and assessment of degree of maturity by org-C/org-N ratio of water extracts. *Soil Sci. Plant Nutr.* 29:251-259.
- Chanyasak, V., and H. Kubota. 1981. Carbon/organic nitrogen ratio in water extracts as a measure of compost degradation. *J. Ferment. Technol.* 59:215-221.
- Chen, Y., and Y. Inbar. 1993. Chemical and spectroscopical analyses of organic matter transformations during composting in relation to compost maturity. p. 551-600. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- DeVleeschauwer, D., O. Verdonck, and P. Van Assche. 1981. Phytotoxicity of refuse compost. *BioCycle* 22:44-46.
- Dick, W.A., and E.L. McCoy. 1993. Enhancing soil fertility by addition of compost. p. 622-644. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- Garcia, C., T. Hernandez, and F. Costa. 1991a. Changes in carbon fractions during composting and maturation of organic wastes. *Environ. Manag.* 15:433-439.
- Garcia, C., T. Hernandez, and F. Costa. 1991b. Agronomic value of urban waste and the growth of ryegrass (*Lolium perenne* L.) in a Calciorthid soil amended with this waste. *J. Sci. Food Agric.* 56:457-467.
- Garcia, C., T. Hernandez, F. Costa, and M. Ayuso. 1992a. Evaluation of the maturity of municipal waste compost using simple chemical parameters. *Commun. Soil Sci. Plant Anal.* 23:1501-1512.
- Garcia, C., T. Hernandez, F. Costa, and J.A. Pascual. 1992b. Phytotoxicity due to the agricultural use of urban wastes. Germination experiments. *J. Sci. Food Agric.* 59:313-319.
- Haug, R.T. 1980. Composting engineering: Principles and practice. Ann Arbor Science Publ., Ann Arbor, MI.
- Hirai, M.F., V. Chanyasak, and H. Kubota. 1983. A standard measurement for compost maturity. *BioCycle* 24:54-56.
- Hoitink, H.A.J., M.J. Boehm, and Y. Hadar. 1993. Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. p. 601-621. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- Iannotti, D.A., T. Pang, B.L. Toth, D.L. Elwell, H.M. Keener, and H.A.J. Hoitink. 1993. A quantitative respirometric method for monitoring compost stability. *Compost Sci. Util.* 1:52-56.
- Iglesias Jimenez, E., and V. Perez Garcia. 1989. Evaluation of city refuse compost maturity: A review. *Biol. Wastes* 27:115-143.
- Inbar, Y., Y. Chen, and H.A.J. Hoitink. 1993. Properties for establishing standards for utilization of composts in container media. p. 668-695. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- Jourdan, B. 1988. Determination of the degree of decomposition for waste and waste/sludge derived compost. *Stuttgarter Berichte zur Abfallwirtschaft*. Vol. 30. E. Schmidt, Verlag, Berlin.
- Keener, H.M., C. Marugg, R.C. Hansen, and H.A.J. Hoitink. 1993. Optimizing the efficiency of the Composting Process. p. 59-94. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- Kubota, H., and K. Nakasaki. 1991. Accelerated thermophilic composting of garbage. *BioCycle* 32:66-68.
- Kuter, G.A., H.A.J. Hoitink, and L.A. Rossman. 1985. Effects of aeration and temperature on composting of municipal sludge in a full-scale vessel system. *J. Water Pollut. Control Fed.* 57:309-315.
- Miller, F.C. 1989. Water matric potential as an ecological determinant in compost, a substrate dense system. *Microbiol. Ecol.* 18:59-71.
- Richard, T., P. Woodbury, V. Breslin, and S. Crawford. 1993. MSW composts: Impacts of separation on trace metal contamination. p. 401-421. In H.A.J. Hoitink and H.M. Keener (ed.) Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publ., Worthington, OH.
- Saviozzi, A., R. Levi-Minzi, R. Riffaldi, and A. Benetti. 1992. Evaluating garbage compost. *BioCycle* 33:72-75.
- Solbraa, K., M.D. Sant, A.R. Selmer-Olson, and H.R. Gislerod. 1983. Composting soft and hardwood barks. *BioCycle* 24(4):44-48,59.
- Still, S.M., M.A. Dirr, and J.B. Gartner. 1976. Phytotoxic effects of several bark extracts on mung bean and cucumber growth. *J. Am. Soc. Hortic. Sci.* 101:34-37.
- Zucconi, F., M. Forte, A. Monaco, and M. de Bertoldi. 1981a. Biological evaluation of compost maturity. *BioCycle* 22:27-29.
- Zucconi, F., A. Pera, M. Forte, and M. de Bertoldi. 1981b. Evaluating toxicity of immature compost. *BioCycle* 22:54-57.

Cross-Polarized Magic-Angle Spinning ^{13}C Nuclear Magnetic Resonance Spectroscopic Characterization of Soil Organic Matter Relative to Culturable Bacterial Species Composition and Sustained Biological Control of Pythium Root Rot

MICHAEL J. BOEHM,¹ TIANYUN WU,¹ ALEX G. STONE,¹ BART KRAAKMAN,¹
DONNA A. IANNOTTI,² G. EDWIN WILSON,³ LAURENCE V. MADDEN,¹
AND HARRY A. J. HOITINK^{1*}

The Ohio State University and Ohio Agricultural Research and Development Center, Wooster,¹ and University of Akron, Department of Chemistry, Akron,³ Ohio, and Division of Natural Sciences and Mathematics, St. Mary's College, St. Mary's City, Maryland²

Received 4 June 1996/Accepted 4 November 1996

We report the use of a model system that examines the dynamics of biological energy availability in organic matter in a sphagnum peat potting mix critical to sustenance of microorganism-mediated biological control of pythium root rot, a soilborne plant disease caused by *Pythium ultimum*. The concentration of readily degradable carbohydrate in the peat, mostly present as cellulose, was characterized by cross-polarized magic-angle spinning ^{13}C nuclear magnetic resonance spectroscopy. A decrease in the carbohydrate concentration in the mix was observed during the initial 10 weeks after potting as the rate of hydrolysis of fluorescein diacetate declined below a critical threshold level required for biological control of pythium root rot. Throughout this period, total microbial biomass and activity, based on rates of [^{14}C]acetate incorporation into phospholipids, did not change but shifts in culturable bacterial species composition occurred. Species capable of inducing biocontrol were succeeded by pleomorphic gram-positive genera and putative oligotrophs not or less effective in control. We conclude that sustained efficacy of naturally occurring biocontrol agents was limited by energy availability to this microflora within the organic matter contained in the potting mix. We propose that this critical role of organic matter may be a key factor explaining the variability in efficacy typically encountered in the control of pythium root rot with biocontrol agents.

Biological control of diseases caused by soilborne plant pathogens through the introduction of microorganisms has been explored intensively for decades (42, 45). Despite numerous contributions aimed at improving our understanding of the four principal mechanisms involved in biological control, competition, antibiosis, hyperparasitism, and systemic acquired resistance in the host plant (1, 14, 26, 29, 30), variability in efficacy of treatments still limits application and commercialization (45). Recent literature on the use of biocontrol agents emphasizes specific interactions between this microflora and pathogens in the rhizosphere (1, 45). It is generally assumed that rhizodeposition products hold sufficient energy reserves to sustain activity of biocontrol agents and thus disease suppression. The role of edaphic soil organic matter (OM) in disease suppression has largely been overlooked.

Knowledge of the role of OM in disease control was derived decades ago from studies utilizing specific amendments, composts, and manures and, more recently, from comparisons of organic and conventional farming systems (7, 21, 49, 50). Before disease control can be achieved, OM must be fully colonized, thus avoiding direct stimulation of plant pathogens (6, 17). Concentrations of readily available nutrients must be low enough that competition prevails and the production of lytic enzymes and antimicrobial compounds involved in hyperparasitism of the pathogen is not repressed (4, 6, 9, 12, 15). Microbiostasis prevails in these suppressive soils (4, 29). Many microorganisms contribute to this mechanism of biological control, particularly for pathogens such as *Pythium* and *Phyto-*

phthora spp. causing root rots (3, 51). It is often referred to as general suppression sensu Gerlach (18). The basis for the length of time that an amendment remains effective (i.e., sustainability) is unknown. However, the duration of suppressiveness presumably depends in part upon the concentration of biologically available energy reserves in OM (7).

Procedures allowing nondestructive quantitative analysis of such energy reserves in OM were not available until recently (23). In the absence of this technology, microbiologists developed indirect approaches to the analysis of soil properties important to biological control. For example, dehydrogenase activity and rates of respiration are higher in suppressive than in conducive highly decomposed substrates (32, 34). The rate of hydrolysis of fluorescein diacetate (FDA activity) predicts the potential for the development of diseases caused by several soilborne plant-pathogenic fungi in both field soil and soilless potting mixes (4, 5, 22, 34, 50, 51). FDA is hydrolyzed by both free and membrane-bound esterases, lipases, and proteases in soil (38, 48). Thus, FDA activity appears to mimic OM decomposition level. FDA activity has provided meaningful insights concerning the microbial carrying capacity of soil OM relative to biological control of diseases caused by several soilborne plant pathogens sensitive to microbiostasis (2-4, 34, 50, 51).

Another approach used to monitor the impact of OM on biological soil properties has been to analyze the microflora itself. Bacterial species diversity among suppressive substrates high in biologically available carbon versus more decomposed conducive substrates does not differ (3). On the other hand, the composition of bacterial species and other microorganisms in the rhizosphere and edaphic soil is affected dramatically by

* Corresponding author.

OM decomposition level in both field soil and soilless potting mixes (3, 21, 49). This also has been shown through analysis of phospholipid fatty acids extracted directly from the rhizosphere and edaphic substrates (43). Pseudomonads, *Pantoea* spp. (formally *Enterobacter* spp.), and other copiotrophic microorganisms such as *Trichoderma* spp. predominate in suppressive potting mixes and soils high in lignocellulosic substances (3, 20, 27). Isolates and strains of these genera rank among the most efficacious biocontrol agents (1, 45). In conducive highly stabilized OM, pleomorphic gram-positive bacteria and putative oligotrophic microorganisms are abundant (3). The latter microflora is not effective in biological control of *Pythium* spp., for example, when applied as seed treatments (40, 41) and has been associated with highly mineralized niches in soils (24). What remains uncharacterized is a direct relationship between OM decomposition level and the dynamics of shifts in microbial populations and functional activities relative to biological control. Answers to these questions should begin to more critically illustrate the role of OM in sustenance of biological control.

In this paper we report on the role of OM in biological control in a light peat potting mix that changes from suppressive to conducive to pythium root rot over a relatively short time during the production of poinsettia. We apply cross-polarized magic-angle spinning ^{13}C nuclear magnetic resonance spectroscopy (CPMAS ^{13}C -NMR) to the nondestructive analysis of the fate of the carbohydrate fraction (mostly cellulosic substances) in OM in a slightly decomposed light sphagnum peat potting mix. We reveal the temporal changes in FDA activity, microbial activity, and biomass, based on the rate of incorporation of [^{14}C]acetate into phospholipids, as well as culturable bacterial species composition over time. It has been shown previously that (i) trends in disease suppressiveness and FDA activity in this potting mix are negatively correlated with pathogen (*Pythium ultimum*) population and disease (poinsettia root rot) development during the production of a crop (2), (ii) the composition of bacterial species varied significantly within peats of various decomposition levels (3), and (iii) peat decomposition level significantly affected the efficacy of biological control agents (3). While others have shown the correlation of OM to biological control, this is the first analysis of the dynamics of the changes in biological energy availability in OM relative to these other factors affecting biological control of root rot.

MATERIALS AND METHODS

Peat potting mixes. Three different sources of sphagnum peat, H_2 , H_3 , and H_4 on the von Post decomposition scale (37), stored in bales at -15°C until use to prevent further decomposition, were from the same batches as those used by Boehm and Hoitink (2). Peats were prepared into potting mixes, treated with fertilizer, planted with poinsettia, and incubated in a greenhouse as described previously (2). The suppressiveness of these peat mixes to pythium root rot has been characterized previously (2).

CPMAS ^{13}C -NMR characterization of peat decomposition level. The decomposition level of H_2 , H_3 , and H_4 sphagnum peat samples was characterized by CPMAS ^{13}C -NMR spectroscopy (11, 19, 36). Changes in the decomposition level of the H_3 peat mix as suppressiveness declined were determined with the NMR spectral subtraction procedure described previously by Hammond et al. (19). At various times after potting, H_3 peat mix samples incubated as described previously (2) were fractionated by washing through a series of mesh sieves (4.00, 2.00, 1.00, 0.500, 0.250, 0.105, and 0.053 mm). Fractionated samples (three replicates per treatment) were lyophilized and stored under vacuum. Two fractions, one containing coarse particles (1- to 2-mm diameter) and the other containing fines (105- to 205- μm diameter), were analyzed by CPMAS ^{13}C -NMR spectroscopy. Spectra were acquired on a home-built 200-MHz spectrometer equipped with a Chemagnetics multinuclear probe utilizing a Tecmag pulse generator (Technology for Magnetic Resonance, Houston, Tex.). Samples were spun at the magic angle at -3 kHz in a zirconium rotor. A contact time of 2 ms for all samples was used to assume full polarization of all carbon atoms and accurate integrations. A 1.0-s pulse delay was used. Chemical shifts were referenced to the methyl of

hexamethylbenzene at 17.0 ppm. The number of scans acquired for each sample was 10,000. The number of datum points was 512 per scan. All peaks in the samples exhibited a single relaxation time, as determined by variation of contact time. Postprocessing of data was performed with the software programs NMRI (New Methods Research, Inc., Syracuse, N.Y.) and Tecmag MACNMR (Technology for Magnetic Resonance).

Microbial activity and biomass. Microbial activity was measured as the rate of hydrolysis of FDA as described previously (2) and by a slightly modified version of the procedure described by van Veldhoven and Mannaerts (44). Microbial activity and total microbial biomass, based on the rate of incorporation of [^{14}C]acetate into phospholipids, were determined as follows. At various times after potting, 5-g peat samples (three replicates per treatment) were collected, suspended in distilled water (95 ml), and shaken. Next, 2.85-ml aliquots of the resulting suspensions were transferred in duplicate to 20-ml glass scintillation vials and amended with 150 μl of [^{14}C]acetate. Controls consisted of vials containing peat suspensions without [^{14}C]acetate and vials containing [^{14}C]acetate alone. Metabolic activity was stopped after a 2-h incubation in the dark on a rotary shaker (175 rpm) by the addition of 7.5 ml of methanol and 3.75 ml of CHCl_3 to establish a methanol- CHCl_3 -water ratio of 2:1:0.8 (vol/vol/vol) (46). The mixture was shaken vigorously for 5 h on a rotary shaker (175 rpm, 24°C). After 18 h, equal volumes of CHCl_3 and water were added to yield a final methanol- CHCl_3 -water ratio of 1:1:0.9 which partitioned the lipid-containing CHCl_3 fraction from the aqueous phase. The CHCl_3 phase (7.0 ml) was then transferred to a new vial and dried under a stream of N_2 (55°C). Finally, the dried lipid was redissolved in 4 ml of CHCl_3 and fractionated for microbial activity and biomass determinations. Specific activity was calculated on the basis of microbial activity (^{14}C incorporation) per unit of biomass. This entire experiment was performed twice.

Bacterial species composition. Shifts in culturable rhizosphere bacterial populations in the H_3 peat mix were assessed by baiting the H_3 peat mix with cucumber seedlings (*Cucumis sativus* L. 'Straight eight', 90% germination) at 8, 38, 57, and 172 days after potting. Rhizosphere bacteria were isolated as described by Boehm et al. (3). Shifts in culturable bacterial populations in the H_3 peat mix were assessed directly by dilution plating of the mix at 6, 100, and 172 days after potting. Bacteria were enumerated, purified, and identified by gas chromatographic-fatty acid methyl ester (GC-FAME) analysis as described previously (3). Strains with a similarity index of ≥ 0.5 were assigned to a taxon, strains with values < 0.5 and ≥ 0.1 were assigned to a genus, and strains with values < 0.1 were assigned to a group (GC similarity group) based on the similarity of their fatty acid profiles. Pure cultures were stored at -70°C in a 15% (vol/vol) sterile glycerol-water solution (39). The relative population density of bacterial taxa for each sample was determined as $100n_i/N$, in which n_i is the number of strains assigned to the i th taxon and N is the total number of strains for a given root tip segment. This experiment was repeated three times.

Efficacy of rhizosphere bacteria as biocontrol agents. The ability of bacteria isolated from cucumber root tips 172 days after potting in the H_3 peat mix to induce biocontrol of damping-off and root rot caused by *P. ultimum* was determined in cucumber bioassays, as described previously (4). The frequency of rhizosphere bacteria with biocontrol activity against *Pythium* recovered from cucumber roots in the H_3 and H_4 peat mixes 10 days after potting was described previously by Boehm et al. (3).

Statistical analyses. Bacterial species (or taxon) diversity was determined for each root tip or peat mix sample (i.e., sampling unit [SU]) at each harvest date as described previously (3). Similarity in bacterial species composition among SUs was measured by using Dice (3, 31) similarity coefficients (resemblance functions) and polar ordination (3, 31). Dice similarity coefficients were calculated as $2a/(2a + b + c)$, in which a is the number of common taxa between two SUs (X and Y), b is the number of unique taxa which occur in SU X but not in SU Y , and c is the number of unique taxa which occur in SU Y but not in SU X . This resemblance function is equal to 0 at "no similarity" and approaches 1 when the SUs are identical in bacterial species composition (31). The polar ordination method of Bray and Curtis (31) was used to position SUs within a coordinate system such that the distances between SUs reflected their similarity in bacterial species composition as well as their relation to underlying environmental gradients such as OM decomposition level. In this study, SUs were arranged on the basis of percent dissimilarity. Percent dissimilarity between SUs was calculated as $100(1 - \text{Dice similarity index})$. The computer programs used for these analyses were SPASSOC.BAS, SUDIST.BAS, and PO.BAS (31).

Chi-square tests were used to determine significant differences in the relative population densities over time (first to last assessment) of (i) gram-positive and gram-negative bacteria (among identified taxa), (ii) GC groups (unknown taxa), and (iii) strains lost upon subculturing during purification and identification (10).

RESULTS

CPMAS ^{13}C -NMR characterization of peat decomposition level. Subtraction spectra of potting mix samples prepared from fine (105- to 250- μm -diameter)-particle fractions of the H_2 and H_4 peats were distinguished from each other by CPMAS ^{13}C -NMR spectroscopy (Fig. 1). Distinct resonances cen-

FIG. 1. Solid-state CPMAS ^{13}C -NMR spectra revealing differences in carbohydrate content within the fine (105- to 250- μm -diameter)-particle fraction of a slightly decomposed light (H_2) and a decomposed dark (H_4) Canadian sphagnum peat. The subtraction spectrum is the H_2 spectrum minus the H_4 spectrum.

tered at ~ 73 and ~ 107 ppm were due to carbohydrates (11, 13, 16, 47). Such resonances in the subtraction spectra (H_2 minus H_4) revealed that the carbohydrate concentration decreased as the degree of peat decomposition increases. Similar results were obtained for the coarse (1- to 2-mm-diameter) particles for these peat mixes (data not shown).

The subtraction spectra of the fine particles (105- to 205- μm diameter) recovered by wet sieving from the H_3 peat mix at 14 (suppressive to root rot) and 77 (conductive) days after planting (2) reveal a decrease in carbohydrate content with time as evidenced by the distinct resonances centered at ~ 73 and ~ 107 ppm (Fig. 2). The equivalent subtraction spectra of the coarse particles (1- to 2-mm diameter) recovered from the same potting mix samples did not reveal this difference (data not shown). NMR spectra of the coarse and fine particles recovered on day 14 were indistinguishable from one another (data not shown). A slight although not significant difference in carbohydrate content was detected in the subtraction spectra of the coarse minus the fine fractions on day 77. In summary, the magnitude of the decrease in carbohydrate content appears directly related to the ratio of surface area to volume for these particles, being most pronounced for the fine particles.

Microbial activity and biomass. FDA activity at the time of planting in the light H_3 peat mix was $4.2 \mu\text{g}$ of FDA $\text{min}^{-1} \text{g}$ (dry weight) of mix $^{-1}$ (Fig. 3). A steady and significant ($P = 0.05$) decline was observed thereafter until after 50 days, where it was below the critical threshold level of $3.2 \mu\text{g}$ of FDA $\text{min}^{-1} \text{g}$ (dry weight) of mix $^{-1}$ required for sustained suppression of pythium root rot (2). FDA activity in the conducive H_4 peat mix was low at all times, thus verifying our previous report for these peat mixes (2).

Microbial biomass increased significantly ($P = 0.05$) after planting (Fig. 4A). It stabilized at $\sim 0.25 \mu\text{mol}$ of $\text{P}_i \cdot \text{g}$ (dry

FIG. 2. CPMAS ^{13}C -NMR spectra revealing the carbohydrate content within the fine (105- to 250- μm -diameter)-particle fractions of H_3 peat samples collected 14 and 77 days after planting. The subtraction spectrum is the 14-day spectrum minus the 77-day spectrum.

weight) of substrate $^{-1}$ between days 10 and 50. Microbial biomass in the conducive H_4 peat mix (Fig. 4A) followed similar trends. It increased significantly ($P = 0.05$) immediately after planting and then stabilized within the range of 0.11 to 0.19 μmol of $\text{P}_i \cdot \text{g}$ (dry weight) of substrate $^{-1}$. Although similar in trends, microbial biomass in the conducive H_4 peat mix was significantly ($P = 0.01$) below that in the H_3 peat mix at all times (Fig. 4A).

Uptake of [^{14}C]acetate revealed a burst in microbial activity within days after potting in the H_3 and H_4 peat mixes (Fig. 4B).

FIG. 3. Microbial activity, based on the rate of hydrolysis of FDA over time in a light (H_3 ; $\square$) and dark (H_4 ; $\blacksquare$) peat mix during the production of poinsettia. Vertical bars represent standard errors ($n = 3$).

FIG. 4. Comparison of total microbial biomass, microbial activity, and specific microbial activity in a slightly decomposed suppressive (H_3 ; □) and a more decomposed conducive (H_4 ; ■) peat. (A) Total microbial biomass, based on the concentration of total phospholipid phosphate; (B) microbial activity, based on the rate of [^{14}C]acetate uptake; (C) specific microbial activity, calculated as the level of microbial activity per unit of biomass. Vertical bars represent standard errors ($n = 6$).

After 24 days, microbial activity remained at a relatively high level even though FDA activity declined (2). Interestingly, rates of [^{14}C]acetate incorporation in the conducive H_4 peat mix did not differ significantly ($P = 0.05$) from those in the H_3 peat mix (Fig. 4B). Specific microbial activity (uptake of [^{14}C]acetate per micromole of P_i per minute) in the conducive H_4 peat mix was consistently higher than that in the H_3 peat mix (Fig. 4C). In a second experiment, trends in microbial biomass, uptake of [^{14}C]acetate, and specific microbial activity were similar to those presented in Fig. 4.

Bacterial species composition. The composition of culturable bacterial species was affected dramatically as the substrate decomposed (Tables 1 and 2). The first of two distinct shifts occurred sometime between 8 and 57 days after potting. *Pseudomonads* and other facultative oligotrophic gram-nega-

TABLE 1. Relative population density of rhizosphere bacteria isolated on 0.1% tryptic soy broth agar from cucumber seedling root tip segments after potting in a slightly decomposed light (H_3) peat potting mix

Taxon ^a	Relative population density at ^b :								
	8 Days in:			57 Days in:			172 Days in:		
	SU 1	SU 2	SU 3	SU 1	SU 2	SU 3	SU 1	SU 2	SU 3
<i>Aeromonas</i> spp.				1.7					
<i>Agrobacterium</i> spp.				3.4	5.6	8.3			
<i>Alcaligenes</i> spp.	3.0			3.4					
<i>Arthrobacter</i> spp.				1.7	1.9	2.1			
<i>Aquaspirillum</i> spp.		2.4							
<i>Aureobacterium</i> spp.				1.7		2.1			2.2
<i>Bacillus</i> spp.		2.4				2.1	2.3		
<i>Bordetella</i> spp.				1.7					4.4
<i>Clavibacter</i> spp.				1.7					2.2
<i>Comamonas</i> spp.							2.3		2.2
<i>Curtobacterium</i> spp.							2.3		
<i>Cytophaga</i> spp.							2.3		2.2
<i>Flavobacterium</i> spp.		4.6					4.5	10.0	2.2
<i>Hydrogenophaga</i> spp.		46.6	20.4	20.7	24.1				
<i>Klebsiella</i> spp.			1.9	5.2					
<i>Microbacterium</i> spp.		2.4					2.3		
<i>Micrococcus</i> spp.	3.0							4.0	26.4
<i>Morganella</i> spp.			3.7						
<i>Ochrobacterium</i> spp.									4.4
<i>Pantoea</i> spp.			7.4	3.4					
<i>Phyllobacterium</i> spp.				3.4	9.3	8.3			
<i>Pseudomonas</i> spp.	60.6	23.2	40.7	36.2	11.2	4.2	4.5	6.0	6.6
<i>Salmonella</i> spp.			5.6						
<i>Sphingobacterium</i> spp.	21.2							2.0	
<i>Variovorax</i> spp.									4.4
<i>Xanthomonas</i> spp.							2.0		6.6
GC similarity groups									
1		2.4	3.7		9.3	12.5			
2			1.9	1.7		10.4			
4			1.9			14.6		7.0	
5		2.4	7.4		1.9	2.1	27.3	28.0	4.4
6						9.3		2.0	
7				1.9	3.4	3.7	10.4		
9		9.1		1.9	1.7	7.4		2.3	
10					1.7	3.7	6.3		
12							4.5		
13									13.2
14							9.1	16.0	
15									2.2
16							6.8	4.0	2.2
17									4.4
Lost ^c	3.0	13.8	1.9	7.3	12.6	16.7	29.5	24.0	8.8

^a Strains were identified by GC-FAME analysis based on the similarity of their phospholipid fatty acid profiles. Strains with a similarity index ≥ 0.5 were assigned to a taxon, values < 0.5 and ≥ 0.1 , were assigned to a genus, and strains with values < 0.1 were assigned to a GC similarity group. Taxa often were identified to the species level, but only genus results are shown.

^b The relative population density of bacterial taxa on each root tip segment (SU) was calculated as follows: $100n_i/N$, where n_i is the number of strains assigned to the i th taxon and N is the total number of strains for a given root tip segment.

^c Strains lost upon subculturing.

tive genera such as *Hydrogenophaga* predominated during the initial 57 days of the growth period. A second shift was observed much later (100 to 172 days). After 172 days, gram-positive pleomorphic bacterial taxa such as *Micrococcus* spp., nonidentifiable strains, and strains lost upon subculturing (possibly oligotrophs) predominated in the rhizosphere and edaphic niches (Tables 1 and 2). Less than 1% of the bacterial

TABLE 2. Temporal dynamics in bacterial species composition as OM decomposed and suppressiveness collapsed during production of poinsettia in an H₃ decomposition level peat substrate

Group	Relative population density ^a at:			
	8 Days ^b		172 Days ^c	
	Mean	Range	Mean	Range
Rhizosphere				
Gram negative	72	49–84	19	14–22****
Gram positive	7	0–17	16	4–36**
<i>Pseudomonas</i> spp.	38	23–61	6	5–6***
GC similarity groups ^e	13	5–23	47	34–54***
Lost	8	3–11	21	8–30**
Peat mix				
Gram negative	27	26–30	23	11–32
Gram positive	13	10–16	7	4–10 ⁺
<i>Pseudomonas</i> spp.	8	4–13	11	7–13
GC similarity groups	53	49–60	65	54–79**
Lost	7	2–10	8	4–13*

^a The relative population densities of bacterial taxa on root tip segments and peat samples were calculated as follows: $100n_i/N$, where n_i is the number of strains assigned to the i th taxon and N is the total strains isolated from a given root tip segment or peat sample.

^b H₃ peat 8 days after potting prior to the decline in disease suppression.

^c H₃ peat 172 days after potting after the decline in disease suppression had taken place.

^d Significant change in the frequency of strains with a given character between 8 and 172 days: *, $P = 0.05$; **, $P = 0.01$; ***, $P = 0.001$.

^e Strains with similarity index values <0.1 were placed in GC similarity groups.

^f Strains lost during subculturing.

strains recovered from the cucumber rhizosphere after 172 days, after the collapse in FDA activity and suppressiveness, were effective as biocontrol agents when applied as cucumber seed treatments against *P. ultimum*.

The Bray-Curtis polar ordination plot (31) based on Dice resemblances (Fig. 5) clearly revealed this shift in composition. SUs representing rhizosphere and niches on day 8 (suppressive) clustered in the upper left portion of the plot. SUs representing bacterial populations recovered from the nonrhizosphere peat mix as suppressiveness collapsed (days 38, 54, and 100) clustered in the upper right portion. Those SUs representing bacterial populations harvested after 172 days, long after the collapse in suppressiveness had occurred, were clearly distinct and clustered in the lower right corner.

DISCUSSION

Differences in carbohydrate content between the slightly decomposed H₂ and the more decomposed H₄ peat samples (Fig. 1) demonstrated here by CPMAS ¹³C-NMR spectroscopy verify previous findings by others interested in plant material transformations to peat and coal (11, 19, 36). The slightly decomposed H₂ peat samples clearly contained higher carbohydrate concentrations compared to the more decomposed H₄ peat. The increases in *P. ultimum* population and root rot and the decrease in FDA activity in the H₃ peat mix, as described previously by Boehm and Hoitink (2 [Fig. 1]), were accompanied by a decrease in NMR-detectable carbohydrate in the fine-particle fraction of this substrate (Fig. 2). The concentration of CPMAS ¹³C-NMR-detectable carbohydrates after 172 days in the now-conductive H₃ peat mix was similar to that in the consistently conductive H₄ peat mix (Fig. 1 and 2). Lack of a detectable decline in the coarse particles suggests that these particles at days 14 and 77 were similar in composition. Carbohydrate content in the fines would most likely be more

FIG. 5. Bray-Curtis polar ordination plot for separation of rhizosphere (closed symbols) or peat (open symbols) SUs based on similarity of bacterial species composition. Data were collected over time as the concentration of biologically available energy reserves decreased and suppressiveness was lost. Sixty to seventy bacterial strains were isolated on 0.1 × tryptic soy broth agar at each sampling time throughout the growth period and identified by GC-FAME analysis. Symbols represent days after potting: Δ , 6; $\blacktriangle$, 8; $\bullet$, 38; $\blacksquare$, 57; $\square$, 100; $\diamond$ and $\blacklozenge$, 172. Dice resemblance functions (31) assessed the relative similarity between bacterial populations. Axes represent percent dissimilarity (31). Differences between points in the two ordination dimensions can be used to access degree of dissimilarity.

accessible to degradation by soil microorganisms because of a greater surface area available for colonization. Although adequate carbohydrate reserves were present in the coarse particles, even after 77 days, the total surface area of these particles relative to that of the fines may have been too low to support a level of microbial activity adequate for effective biological control of pythium root rot of poinsettia. Alternatively, lack of availability of carbohydrates present in the interior components of the coarse particles with decomposed outer surfaces may also have accounted for the decline in microbial activity. A direct spectroscopic procedure, such as near-infrared reflectance spectroscopy, capable of assessing in situ surface chemistry may provide additional information (35).

A significant shift in culturable rhizosphere bacterial species composition accompanied the decline in carbohydrate content in the H₃ peat mix (Tables 1 and 2). At potting in this mix, 10% of the rhizosphere microflora (mostly pseudomonads and *Pantoea* spp.) was capable of inducing biological control of pythium damping-off and root rot (3), compared to <1% on day 172. This is similar to the percentage of bacteria recovered from the conductive H₄ peat mix capable of causing biocontrol at 10 days after planting (3). Pleomorphic gram-positive bacteria and oligotrophs typical of those recovered from highly mineralized soil fractions (24) and the conductive H₄ peat mix (3) were abundant in the conductive H₃ peat mix at 172 days (Tables 1 and 2). In the H₃ mix, gram-negative bacteria had significantly declined in populations after 172 days, whereas gram positives and oligotrophs increased (Table 2). As the carbohydrate content of the H₃ mix decreased (Fig. 2) and suppressiveness declined (2), a microflora capable of inducing biological control was replaced by another that could not. This finding, in conjunction with that presented previously regarding the effect of OM decomposition level on the efficacy of introduced biocontrol agents (3), suggests strongly that OM is critical to maintenance and activity (sustained activity) of a microflora capable of inducing biological control.

In light of the previous work done with this H₃ peat mix (2, 3), it is interesting that suppressiveness was lost before bacteria capable of inducing biocontrol of pythium root rot had declined to low relative population densities. The H₃ peat mix was conducive after approximately 50 days, when FDA activity was low (2). At this time after potting, the population of pseudomonads, and other facultative oligotrophic gram-negative genera often described as biocontrol agents, was still relatively high (Table 2) even though a significant shift in species composition had already occurred (Fig. 5). This indicates, quite possibly, that the nutritional status of the biocontrol agents at that time was inadequate to sustain biological control activity.

The lack of a difference in microbial activity (based on the rates of incorporation of [¹⁴C]acetate into phospholipids) in the H₃ peat mix (Fig. 4B) indicates that lack of competition for nutrients could not solely explain the loss of suppressiveness observed previously (2). This observation supports an earlier report by Liebman and Epstein (28), which showed that competition for nutrients is not the principal factor limiting germination of fungal spores when microbiostasis prevails but rather that other factors played a significant role. The significantly ($P = 0.05$) higher level of specific microbial activity observed in the conducive H₄ peat mix (Fig. 4C), relative to that in the H₃ peat mix, suggests that the microflora in the conducive environment utilized the nutrients more efficiently.

The overall conclusions of this work are that energy availability in soil OM critically determined survival and efficacy of bacteria in biological control of pythium damping-off and root rot. Rhizodeposition products such as root exudates, mucigel, and sloughed root cells, although important as energy sources for rhizosphere bacteria (8, 25, 33), appeared inadequate to sustain prolonged suppressiveness in the conducive mixes. It appears that an adequate supply of biologically available energy must be available within the soil system prior to the introduction of a plant to maintain or carry a critical threshold level of biocontrol agent activity through the cropping period as roots grow through the soil. The foregoing offers an explanation for the lack in efficacy often associated with introduced biocontrol agents in field soils where this role of OM is usually not taken into consideration.

ACKNOWLEDGMENTS

We gratefully acknowledge C. A. Musselman and M. S. Krause for technical assistance and W. D. Bauer and L. H. Rhodes for critical review of the manuscript.

Salaries and research support were provided in part by state and federal funds appropriated to the Ohio Agricultural Research and Development Center, The Ohio State University, and from grant US-2196-22 BARD, United States-Israel Binational Agricultural Research and Development Fund.

REFERENCES

- Adams, P. B. 1990. The potential of mycoparasites for biological control of plant diseases. *Annu. Rev. Phytopathol.* 28:59-72.
- Boehm, M. J., and H. A. J. Hoitink. 1992. Sustainment of microbial activity in potting mixes and its impact on severity of *Pythium* root rot of poinsettia. *Phytopathology* 82:259-264.
- Boehm, M. J., L. V. Madden, and H. A. J. Hoitink. 1993. Effect of organic matter decomposition level on bacterial species diversity and composition in relationship to pythium damping-off severity. *Appl. Environ. Microbiol.* 59:4171-4179.
- Chen, W., H. A. J. Hoitink, and L. V. Madden. 1988. Microbial activity and biomass in container media for predicting suppressiveness to damping-off caused by *Pythium ultimum*. *Phytopathology* 78:1447-1450.
- Chen, W., H. A. J. Hoitink, A. F. Schmitthenner, and O. H. Tuovinen. 1988. The role of microbial activity in suppression of damping-off caused by *Pythium ultimum*. *Phytopathology* 78:314-322.
- Chung, Y., H. A. J. Hoitink, W. A. Dick, and L. J. Herr. 1988. Effects of organic matter decomposition level and cellulose amendment on the inoculum potential of *Rhizoctonia solani* in hardwood bark media. *Phytopathology* 78:836-840.
- Cook, R. J., and K. F. Baker. 1983. The nature and practice of biological control of plant pathogens. American Phytopathological Society. St. Paul, Minn.
- Curl, E. A., and B. Truelove. 1986. The rhizosphere. Springer-Verlag, New York, N.Y.
- de la Cruz, J., M. Rey, J. M. Lora, A. Hidalgo-Galleso, F. Dominguez, J. A. Pintor-Toro, A. Llobell, and J. Benitez. 1993. Carbon source control on β -glucanases, chitinase and chitinase from *Trichoderma harzianum*. *Microbiology* 159:316-322.
- Devore, J., and R. Peck. 1986. Statistics: the exploration and analysis of data. West Publishing Co., New York, N.Y.
- Dudley, R. L., C. A. Fyfe, and C. M. Preston. 1990. A ¹³C-CPMAS NMR spectroscopic study of the transformation of plant material to peat and coal. *Can. J. Spectrosc.* 35(2):31-38.
- Elad, Y., and I. Chet. 1987. Possible role of competition for nutrients in biocontrol of *Pythium* damping-off. *Phytopathology* 77:190-195.
- Filip, Z., R. H. Newman, and J. J. Alberts. 1991. Carbon-13 nuclear magnetic resonance characterization of humic substances associated with salt marsh environments. *Sci. Total Environ.* 101:191-199.
- Fravel, D. R. 1988. Role of antibiosis in the biological control of plant diseases. *Annu. Rev. Phytopathol.* 26:75-91.
- Friedlander, M., Y. Inbar, and I. Chet. 1993. Biological control of soilborne plant pathogens by a β -1,3 glucanase-producing *Pseudomonas cepacia*. *Soil Biol. Biochem.* 25:1211-1221.
- Fründ, R., and H.-D. Lüderrmann. 1991. Quantitative characterization of soil organic matter and its fraction products by solid state high resolution C-13 (CPMAS) spectroscopy. *Z. Naturforsch.* 46:982-988.
- Garrett, S. D. 1970. Pathogenic root-infecting fungi. University Press, Cambridge, United Kingdom.
- Gerlagh, M. 1968. Introduction of *Ophiobolus graminis* into new polders and its decline. *Neth. J. Plant Pathol.* 749(Suppl.2):1-97.
- Hammond, T. E., D. G. Cory, W. M. Ritchey, and H. Morita. 1985. High resolution solid state ¹³C n.m.r. of Canadian peats. *Fuel* 64:1687-1695.
- Hardy, G. E., and K. Sivasithamparan. 1991. How container media and matrix potential affect the production of sporangia, oospores and chlamydospores by three *Phytophthora* species. *Soil Biol. Biochem.* 23:31-39.
- Hassink, J., J. H. Oude Voshaar, E. H. Nijhuis, and J. A. van Veen. 1991. Dynamics of the microbial populations of a reclaimed-polder soil under a conventional and a reduced-input farming system. *Soil Biol. Biochem.* 23:515-524.
- Inbar, Y., M. J. Boehm, and H. A. J. Hoitink. 1991. Hydrolysis of fluorescein diacetate in sphagnum peat container media for predicting suppressiveness to damping-off caused by *Pythium ultimum*. *Soil Biol. Biochem.* 23:479-483.
- Inbar, Y., Y. Chen, and Y. Hadar. 1989. Solid state carbon-13 nuclear magnetic resonance and infrared spectroscopy of composted organic matter. *Soil Sci. Soc. Am. J.* 53:1695-1701.
- Kanazawa, S., and Z. Filip. 1986. Distribution of microorganisms, total biomass, and enzyme activities in different particles of brown soils. *Ecology* 67:205-215.
- Klein, D. A., J. L. Salzwedel, and F. B. Dazzo. 1990. Microbial colonization of plant roots, p. 189-225. In J. P. Nakas, and C. Hagedorn (ed.), *Biotechnology of plant-microbe interactions*. McGraw-Hill, Inc., New York, N.Y.
- Kuc, J., and N. E. Strobel. 1992. Induced resistance using pathogens and non-pathogens, p. 295-300. In E. C. Tjamos (ed.), *Biological control of plant diseases*. Plenum Press, New York, N.Y.
- Kuter, G. A., E. B. Nelson, H. A. J. Hoitink, and L. V. Madden. 1983. Fungal populations in container media amended with composted hardwood bark suppressive and conducive to *Rhizoctonia* damping-off. *Phytopathology* 73:1450-1456.
- Liebman, J. A., and L. Epstein. 1992. Activity of fungistatic compounds from soil. *Phytopathology* 82:147-153.
- Lockwood, J. L. 1988. Evolution of concepts associated with soilborne plant pathogens. *Annu. Rev. Phytopathol.* 26:93-121.
- Loper, J. E., and J. S. Buyer. 1991. Siderophores in microbial interactions on plant surfaces. *Mol. Plant-Microbe Interact.* 4:5-13.
- Ludwig, J. A., and J. F. Reynolds. 1988. Statistical ecology: a primer on methods and computing. John Wiley and Sons, New York, N.Y.
- Lumsden, R. D., P. D. Millner, and J. A. Lewis. 1986. Suppression of lettuce drop caused by *Sclerotinia minor* with composted sewage sludge. *Plant Dis.* 70:197-201.
- Lynch, J. M., and J. M. Whipps. 1990. Substrate flow in the rhizosphere. *Plant Soil* 129:1-10.
- Mandelbaum, R., and Y. Hadar. 1990. Effects of available carbon source on microbial activity and suppression of *Pythium aphanidermatum* in compost and peat container media. *Phytopathology* 80:794-804.
- McLellen, T. M., J. D. Aber, M. E. Martin, J. M. Melillo, and K. J. Nadelhoffer. 1991. Determination of nitrogen, lignin, and cellulose content of decomposing leaf material by near infrared reflectance spectroscopy. *Can. J. For. Res.* 21:1684-1688.

36. Preston, C. M., S.-E. Shipitalo, R. L. Dudley, C. A. Fyfe, S. P. Mathur, and M. Levesque. 1987. Comparison of ^{13}C CPMAS NMR and chemical techniques for measuring the degree of decomposition in virgin and cultivated peat profiles. *Can. J. Soil Sci.* 67:187-198.
37. Puustjärvi, V., and R. A. Robertson. 1975. Physical and chemical properties, p. 23-28. *In* D. W. Robinson and J. G. D. Lamb (ed.), *Peat in horticulture*. Academic Press, Inc., New York, N.Y.
38. Schnürer, J., and T. Rosswall. 1982. Fluorescein diacetate hydrolysis as a measure of total activity in soil and litter. *Appl. Environ. Microbiol.* 43:1256-1261.
39. Sleesman, J. P., and C. Leben. 1978. Preserving phytopathogenic bacteria at -70°C or with silica gel. *Plant Dis. Rept.* 62:910-913.
40. Sugimoto, E. E., H. A. J. Hoitink, and O. H. Tuovinen. 1990. Enumeration of oligotrophic rhizosphere pseudomonads with diluted and selective media formulations. *Biol. Fertil. Soils* 9:226-230.
41. Suslow, T. V., and M. N. Schroth. 1982. Rhizobacteria of sugar beets: effects of seed application and root colonization on yield. *Phytopathology* 72:199-206.
42. Taylor, A. G., and G. E. Harman. 1990. Concepts and technologies of selected seed treatments. *Annu. Rev. Phytopathol.* 28:321-339.
43. Tunlid, A., H. A. J. Hoitink, C. Low, and D. C. White. 1989. Characterization of bacteria that suppress *Rhizoctonia* damping-off in bark compost media by analysis of fatty acid biomarkers. *Appl. Environ. Microbiol.* 55:1368-1374.
44. van Veldhoven, P. P., and G. P. Mannaerts. 1987. Inorganic and organic phosphate measurements in the nanomolar range. *Anal. Biochem.* 161:45-48.
45. Weller, D. M. 1988. Biological control of soilborne plant pathogens in the rhizosphere with bacteria. *Annu. Rev. Phytopathol.* 26:379-407.
46. White, D. C., W. M. Davies, J. S. Nickels, J. D. King, and R. J. Bobbies. 1979. Determination of the sedimentary microbial biomass by extractable lipid phosphate. *Oecologia* 40:51-62.
47. Wilson, M. A. 1987. NMR techniques and applications in geochemistry and soil chemistry, p. 237-247. Pergamon Press, Oxford, United Kingdom.
48. Wirth, J. S. 1992. Detection of soil polysaccharide *endo*-hydrolase activity profiles after gel permeation chromatography. *Soil Biol. Biochem.* 24:1185-1188.
49. Workneh, F., and A. H. C. van Bruggen. 1994. Microbial density, composition, and diversity in organically and conventionally managed rhizosphere soil in relation to suppression of corky root of tomatoes. *Appl. Soil Ecol.* 1:219-230.
50. Workneh, F., A. H. C. van Bruggen, L. E. Drinkwater, and C. Shennan. 1993. Variables associated with corky root and *Phytophthora* root rot of tomatoes in organic and conventional farms. *Phytopathology* 83:581-589.
51. You, M. P., and K. Sivasithamparam. 1994. Hydrolysis of fluorescein diacetate in an avocado plantation mulch suppressive to *Phytophthora cinnamomi* and its relationship with certain biotic and abiotic factors. *Soil Biol. Biochem.* 26:1355-1361.

Compost-Induced Systemic Acquired Resistance in Cucumber to Pythium Root Rot and Anthracnose

W. Zhang, W. A. Dick, and H. A. J. Hoitink

The Ohio Agricultural Research and Development Center, The Ohio State University, Wooster 44691-4096.

This research was supported by AgriEnergy Resources, Princeton, IL, by grant US-2196-22 from BARD—The United States-Israel Binational Research and Development Fund—and by state and federal funds appropriated to the Ohio State University and the Ohio Agricultural Research and Development Center, Wooster. W. Zhang also obtained support from the National Laboratory of Grassland Ecological Engineering, Northeast Normal University, Changchun, P. R. China.

We thank J. Kuć, Department of Plant Pathology, University of Kentucky, Lexington, for supplying a culture of *Colletotrichum orbiculare*.

J. Durkalski and C. A. Musselman provided valuable technical assistance.

Accepted for publication 8 July 1996.

ABSTRACT

Zhang, W., Dick, W. A., and Hoitink, H. A. J. 1996. Compost-induced systemic acquired resistance in cucumber to *Pythium* root rot and anthracnose. *Phytopathology* 86:1066-1070.

Pythium root and stem rots cause problems in production of greenhouse and nursery crops. Composts, however, can provide biological control of these diseases. We planted and germinated cucumber (*Cucumis sativus* 'Straight Eight') seeds in compost-amended (spruce or pine bark composts) or in highly decomposed sphagnum peat (H₄ on the von Post decomposition scale) mixes suppressive and conducive to *Pythium* root rot, respectively. Two-week-old seedlings were transplanted, using the split-root technique, into the compost-amended mixes and the peat mix. Split-root pairings were peat/compost, peat/peat, and compost/compost. Only one side of the split roots was grown in potting mix infested with *Pythium ultimum* and *P. aphanidermatum*. Root rot in the infested mix, averaged across all split-root pairings, was significantly ($P \leq 0.05$) less severe and mean root dry weights were significantly ($P \leq 0.05$) higher in plants germinated in the compost-amended mixes than in the peat mix.

Also, root rot in the infested peat mix was significantly less severe on roots paired with the compost-amended mixes than those paired with the peat mix. Plants grown from seed in the compost-amended mixes or the peat mix also were tested for resistance to anthracnose. Three weeks after planting, anthracnose on the second leaf of cucumber 'Straight Eight' inoculated with *Colletotrichum orbiculare* was significantly less severe ($P \leq 0.05$) on plants grown in the compost-amended mixes than in the peat mix. Peroxidase activity, a putative marker of systemic acquired resistance (SAR) in cucumber, was significantly ($P \leq 0.05$) enhanced in plants grown in the compost-amended mixes compared to the peat mix. However, the activity of a peroxidase isozyme in the second leaf of cucumber plants was greater when plants were grown in compost as well as after prior inoculation with *C. orbiculare* than if grown in peat and with or without prior inoculation. The interaction of compost and the pathogen appeared critical for rapid activation of SAR-associated gene expression in cucumber plants.

Additional keyword: cucumber anthracnose.

Since the early 1970s, composts have been introduced as peat substitutes and used effectively for control of plant diseases caused by soilborne plant pathogens (12). In the nursery industry, disease-suppressive compost-amended container media have been effective enough to replace methyl bromide (25). Numerous reports reveal that microbiostasis and mycoparasitism can explain the disease control activity of composts (5,10,17,18,20). Foliar diseases also may be affected by composts incorporated into soil (31,36). Systemic acquired resistance (SAR) has been proposed as a mechanism underlying this activity (36), but conclusive evidence has not been presented.

Many reports show that composts suppress *Pythium* root rots (2,3,5-7,10,12,20,21). Microbiostasis is involved in the suppression induced by composts against *Pythium* spp. (3,5,20,21), but competition for carbon can only partially explain the suppressive effect (37). *Pseudomonas* spp. provide biological control and predominate in both rhizosphere and edaphic niches of compost-amended substrates suppressive to *Pythium* root rot (2,3). In contrast, gram-positive pleomorphic genera and putative oligotrophs that are incapable of inducing biological control are most abundant in consistently conducive, highly decomposed sphagnum peat substrates

(3). Populations of pseudomonads and other biocontrol agents decline as organic matter decomposes, and suppressiveness to *Pythium* root rot is lost (37). Zhou and Paulitz (41), utilizing a split-root technique with cucumber, recently showed that plant growth-promoting pseudomonads provide *Pythium* root rot control of cucumber by inducing SAR. It is plausible, therefore, that SAR plays a role in suppression of *Pythium* root rot in compost-amended substrates.

Plant growth-promoting rhizobacteria and fungi also can induce SAR against foliar diseases of plants (19,22,23,34,35). A procedure has been developed to induce SAR against anthracnose of cucumber, whereby the first cucumber leaf is treated with the necrogenic pathogen *Colletotrichum orbiculare* (Berk. & Mont.) Arx (formerly *C. lagenarium* (Pass.) Ellis & Halst.) (16,34,38). Resistance to a second challenge-inoculation by *C. orbiculare* on the second leaf of the cucumber plant is then used as a test for the SAR response. An acidic peroxidase isozyme has been identified as a molecular marker of SAR in plants (1,11,24,27,28).

The objectives of this work were (i) to determine whether compost-amended potting mixes induce SAR in cucumber against *Pythium* root rot, utilizing a split-root procedure (41), and (ii) against anthracnose in cucumber foliage caused by *C. orbiculare* and (iii) to determine whether peroxidase activity is increased in plants grown in composts suppressive to root rot compared to plants grown in a Sphagnum peat mix conducive to root rot. A preliminary report has been published (40).

Corresponding author: W. A. Dick; E-mail address: dick.5@osu.edu

Publication no. P-1996-0812-01R
© 1996 The American Phytopathological Society

MATERIALS AND METHODS

Potting mixes and inocula. A composted pine bark-amended potting mix was obtained from Strong-Lite, Inc., Pine Bluff, AR (experiment 1). A similar composted pine bark-amended mix and a composted spruce bark-amended mix were obtained from Earthgro, Inc., Lebanon, CT (experiments 2 through 4). These compost-amended mixes were naturally suppressive to *Pythium* root rot (2, 12). A dark, decomposed sphagnum peat (H_4 on the von Post decomposition scale) mix that was consistently conducive to *Pythium* root rot (12) was prepared as described previously (2). Soil inocula of *Pythium ultimum* Trow 211 and *P. aphanidermatum* (Edson) Fitzp. 20 were prepared as described previously (6), using Ko and Hora's chopped-potato soil medium (15). Potting mixes were infested by incorporating 0.5 g of *P. ultimum* and 0.2 g of *P. aphanidermatum* soil inocula per liter. The mixes were amended with 18 g of slow-release fertilizer (Osmocote, 14-14-14) per liter.

Pythium root rot bioassay on split-root cucumber plants. Cucumber seeds (*Cucumis sativus* L. 'Straight Eight') were germinated either in the compost amended mixes or the peat mix as described previously (2). Roots of 2-week-old seedlings were removed from the germination mixes, washed in running tap water, divided into two approximately equal portions, and transplanted as split-root plants in 600-ml polystyrene pots (one plant per pot). Roots were split and placed in separate compartments, using a Plexiglas shield sealed with silicone to prevent direct contact or exchange of materials, including water or solution flux. The history of whether plants were germinated in the compost-amended mixes or the peat mix was recorded, and split-root pairings were prepared; they consisted of peat/compost, peat/peat, and compost/compost. One side of each pot was infested with *Pythium* soil inocula prepared as described above. Controls were not infested on either side of the pot containing the split roots. Root dry weight (60°C for 24 h) and root rot severity were measured 35 days after planting in the infested side of the pot. Root rot severity was rated by a scale in which 1 = symptomless, 2 = mild root rot (less than one-third of root rotted), 3 = moderate root rot (one- to two-thirds of root rotted), and 4 = severe root rot (more than two-thirds of root rotted or plant dead). Infested roots were plated on selective medium to verify the identity of the pathogen, as described previously (6).

Cucumber anthracnose bioassay. An isolate of *C. orbiculare* was obtained from J. Kuć, Department of Plant Pathology, University of Kentucky, Lexington. Classic SAR was induced 14 days after seeding by prior inoculation on the abaxial side of the first leaf with a conidial suspension of *C. orbiculare* (30 10- μ l drops, 8.6×10^4 CFU ml⁻¹) (16). Inoculated plants were placed in a growth chamber (25°C, 75% relative humidity, 14 h of light per 10 h of dark). Resistance was detected via challenge-inoculation 7 days later by placing 30 10- μ l drops of a conidial suspension (10^4 to 10^5 CFU ml⁻¹) on the second leaf of the same plants and incubating in a moisture chamber (25°C, 24 h of dark). Anthracnose severity on the second leaf was rated 7 days later by a rating system that involved counting the number and diameter of lesions (experiment 3) and overlaying the leaves with a transparent sheet containing a lined grid (5 $\times$ 5 mm square) followed by photographing of the leaves (experiment 4). The necrotic lesion area and total leaf area were measured by counting the number of squares (5 $\times$ 5 mm) represented by each. Disease severity was expressed as the percentage of total leaf area affected by necrosis.

Peroxidase activity. For the split-root experiments, the second leaf of the cucumber plant was removed 21 days after planting and was frozen in liquid nitrogen until further analyses. Peroxidase activity, assayed using guaiacol as the hydrogen donor, as described previously (11), was expressed as change in absorbance at 470 nm min⁻¹ mg⁻¹ of protein.

Electrophoretic analysis for peroxidase. The second leaf of the cucumber plant was removed 7 days after challenge-inocula-

tion on the same leaf and was frozen in liquid nitrogen. Previously frozen leaves were placed in a porcelain mortar with additional liquid nitrogen and ground using a pestle cooled in the refrigerator (4°C). Equal amounts of protein were loaded for each treatment, and peroxidase isozymes were separated and detected as previously described (4). Gels were stained at room temperature with a solution containing 5 ml of 3% hydrogen peroxide (H_2O_2) and 10 ml of *O*-dianisidine solution until bands were detected (about 1 h) (11). Destaining was accomplished with a 50% glycerol solution, and the stained gel was photographed. Analysis was performed for plants grown in the peat, composted pine bark-amended (experiment 2), and composted spruce bark-amended mixes (experiment 3).

Experimental designs and statistical analysis. All treatments were blocked (six replications) with treatments randomized within each block. Each experiment was performed at least twice. Data were subjected to analysis of variance. When a significant ($P \leq 0.05$) *F* test was obtained for treatments, separation of means was accomplished by the least significant different test (LSD_{0.05}).

RESULTS

Pythium root rot. *Pythium* root rot on cucumber seedlings germinated in compost-amended mixes was significantly ($P \leq 0.05$) less severe than on seedlings germinated in peat mix, regardless of whether both sides of the split roots were transplanted into compost or peat mixes (Table 1, top). The overall treatment mean for root rot severity of plants germinated in the composted pine bark-amended mix was 1.5 versus 1.8 in the peat mix. The overall treatment mean was 1.6 for plants germinated in the composted spruce bark-amended mix versus 2.0 in the peat mix. Dry weight of cucumber roots also was significantly ($P \leq 0.05$) greater for plants germinated in the two compost-amended mixes than in the peat mix (Table 1, top). Root dry weights of plants germinated in the compost-amended mixes was 0.12 g per plant compared to 0.09 g per plant germinated in the peat mix.

Mean root rot severity was significantly ($P \leq 0.05$) less for split roots in the infested peat mix paired with the compost-amended

TABLE 1. *Pythium* root rot severity (mean of six replicates) in cucumber plants (*Cucumis sativus* 'Straight Eight') grown in infested compost-amended mixes versus a peat mix

Treatment	Exps. 1 and 2		Exps. 3 and 4	
	Root rot severity ^a	Root dry wt (g)	Root rot severity ^a	Root dry wt (g)
Germination mix ^b				
Composted pine bark	1.5	0.12		
Composted spruce bark			1.6	0.12
Peat	1.8	0.09	2.0	0.09
LSD _{0.05}	0.1	0.02	0.3	0.01
Paired split root ^c				
P/C (pine bark)	1.5	0.12		
P/P	2.2	0.07		
LSD _{0.05}	0.3	0.03		
P/C (spruce bark)			1.6	0.11
P/P			2.4	0.07
LSD _{0.05}			0.4	0.02

^a Rating value: 1 = symptomless; 2 = mild root rot (less than one-third of root rotted); 3 = moderate root rot (one- to two-thirds of root rotted); and 4 = severe root rot (more than two-thirds of root rotted or plant dead).

^b Data were combined from plants for which both sides of the split-root system were grown in the same mixture (i.e., composts or peat), but prior to transplanting, seeds were germinated in either the compost-amended mixes or the peat mix.

^c Treatment means presented are for root rot severity ratings of plants grown in infested peat (P_i) paired with either compost-amended mixes (not infested) or the peat mix (not infested).

mixes than in the infested peat mix paired with the noninfested peat mix (Table 1, bottom). Roots grown in the infested peat mix paired with the composted pine bark-amended mix (not infested) had a root rot severity rating of 1.5 versus a rating of 2.2 when the infested peat mix was paired with the noninfested peat mix. A similar comparison of roots in the infested peat mix paired with the composted spruce bark-amended mix (not infested) yielded root rot severity values of 1.6 versus 2.4. However, root rot severity of plants grown in infested compost-amended mixes paired with compost (not infested) and plants grown in infested peat mix paired with the same compost (not infested) were not statistically different (data not shown). *P. ultimum* and *P. aphanidermatum* were recovered from infested plants on selective medium. Control plants were free of root rot.

After split-root transplanting, dry weight of cucumber roots grown in the infested peat mix paired with either compost-amended mix (not infested) was significantly greater ($P \leq 0.05$) than that of roots grown in the infested peat mix paired with the noninfested

peat mix (Table 1, bottom). Root dry weight was significantly correlated with root rot severity ($r = 0.64$, $n = 36$, $P \leq 0.05$).

Anthraco severity. Anthracnose on the second leaf of cucumber plants was significantly ($P \leq 0.05$) less severe on plants grown in the composted spruce bark-amended mix versus the peat mix (Table 2). Anthracnose severity on plants grown in the composted pine bark-amended mix also was less severe than on plants grown in the peat mix, but the difference was significant at $P \leq 0.10$. Disease severity in the peat and composted pine bark-amended mixes was significantly ($P \leq 0.05$) less on plants after prior inoculation of the first leaf with *C. orbiculare* (induced) compared to plants without prior inoculation. The necrotic lesion areas on the second leaves of noninduced plants ranged from 85.7% for plants grown in the peat mix to 42.7% for plants grown in the composted spruce bark-amended mix and 63.3% for plants grown in the composted pine bark-amended mix. There was no significant difference in disease severity between plants grown in the composted spruce bark-amended mix versus those in which SAR was induced (regardless of mix treatment) through prior inoculation of first leaves with *C. orbiculare* (percentage of lesion area [35.7%], Table 2). Disease severity on plants grown in the composted pine bark-amended mix (percentage of lesion area [63.3%]) was significantly ($P \leq 0.05$) greater than on plants with prior inoculation of first leaves with *C. orbiculare*, but it was significantly less ($P \leq 0.10$) on plants grown in the peat mix.

Peroxidase activity. Peroxidase activity in leaves of cucumber plants was significantly ($P \leq 0.05$) increased in plants germinated in the compost-amended mixes versus those germinated in the peat mix (Table 3, top). After split-root transplanting, peroxidase activities also were greater when roots were paired with the composted spruce bark or pine bark mixes than when roots were paired with the peat mix (Table 3, bottom). The highest peroxidase activity was observed when the peat mix was paired with the composted

TABLE 2. Severity of anthracnose caused by *Colletotrichum orbiculare* on the second leaf of cucumber plants (*Cucumis sativus* 'Straight Eight') in compost-amended mixes versus a peat mix

Potting mix	First leaf inoculation ^y	Anthraco severity (%) ^z
Composted pine bark	—	63.3 ab
Composted spruce bark	—	42.7 bc
Peat	—	85.7 a
Overall mean		63.9
Composted pine bark	+	28.0 c
Composted spruce bark	+	35.7 c
Peat	+	35.7 c
Overall mean		33.5

^y + = systemic resistance induced by prior inoculation of the first leaf with *C. orbiculare*; — = mimic inoculation with distilled water only on the first leaf.

^z Anthracnose severity was measured on the second leaf for all treatments and is expressed as the percentage of total leaf area exhibiting necrosis. Measurements were made by overlaying a transparent sheet containing a lined grid (5 × 5 mm square) on leaves and counting the necrotic lesion area and total leaf area. Experiments were performed four times with six replicates for each experiment. Means with different letters are significantly different at $P \leq 0.05$. LSD_{0.05} = 24.0.

TABLE 3. Peroxidase activity in leaves (mean of six replicates) of cucumber plants (*Cucumis sativus* 'Straight Eight') grown in compost-amended mixes versus a peat mix

Treatment	Peroxidase activity ^a	
	Exps. 1 and 2	Exps. 3 and 4
Germination mix^y		
Composted pine bark	4.3	4.6
Composted spruce bark		
Peat	3.5	3.4
LSD _{0.05}	0.6	0.9
Paired split root^z		
P ₁ /C (pine bark)	4.7	
P ₁ /P	3.7	
LSD _{0.05}	0.8	
P ₁ /C (spruce bark)		5.0
P ₁ /P		3.9
LSD _{0.05}		1.2

^a Peroxidase activity is expressed as the change in absorbance at 470 nm min⁻¹ mg⁻¹ of protein.

^y Data were combined from plants in which both sides of the split-root system were grown in the same mixture (i.e., composts or peat), but prior to transplanting, seeds were germinated in either the compost-amended mixes or the peat mix.

^z Treatment means presented are for root rot severity ratings of plants grown in infested peat (P₁) paired with either the compost-amended mixes (not infested) or the peat mix (not infested).

Fig. 1. Peroxidase isozyme expression in the second leaf of cucumber plants (*Cucumis sativus* 'Straight Eight') grown in compost-amended mixes or a peat mix. Some plants were induced to express systemic resistance by prior inoculation of the first leaf with *Colletotrichum orbiculare*. Challenge-inoculation on the second leaf was with *C. orbiculare* 7 days after prior inoculation of the first leaf. Lanes 1 and 10, plants grown in peat that were induced and challenged with *C. orbiculare*. Lanes 2 and 11, plants grown in peat that were not induced but were challenged with *C. orbiculare*. Lanes 3 and 12, plants grown in peat and not induced or challenged. Lane 4, plants grown in composted spruce bark-amended mix that were induced and challenged with *C. orbiculare*. Lane 5, plants grown in composted spruce bark-amended mix that were not induced or challenged. Lane 6, plants grown in composted spruce bark-amended mix that were induced and challenged with *C. orbiculare*. Lane 7, plants grown in composted pine bark-amended mix that were not induced or challenged. Lane 8, plants grown in composted pine bark-amended mix that were induced but were challenged with *C. orbiculare*. Lane 9, plants grown in composted pine bark-amended mix that were not induced or challenged.

spruce bark-amended mix, and the activity was significantly ($P \leq 0.05$) higher than in plants grown in the peat mix paired with itself. Similar results were obtained when the peat mix was paired with the composted pine bark mix.

Relative activity of a peroxidase isozyme in the second leaves of cucumber plants grown in the peat mix was low (Fig. 1, lanes 3 and 12). Activity in the leaves from plants produced in the composted spruce bark- and pine bark-amended mixes also was very low (Fig. 1, lanes 6 and 9, respectively), although some very faint bands, which do not show in the figure, were evident. Inoculation with *C. orbiculare* on the second leaf dramatically increased the activity of this peroxidase isozyme in plants grown in the peat-amended mixes (Fig. 1, lanes 2 and 11), but this increase was even greater in plants grown in the compost-amended mixes (Fig. 1, lanes 4 and 8). The highest activity occurred when plants were both grown in compost-amended mixes that were prior inoculated as well as challenge-inoculated with *C. orbiculare* (Fig. 1, lanes 5 and 7). Even here, peroxidase levels were higher in the two compost-amended mixes than in the peat mix.

DISCUSSION

The significantly ($P \leq 0.05$) reduced root rot severity observed in split roots of plants produced in the *Pythium*-infested conducive peat mix paired with the suppressive compost-amended mixes (Table 1) suggests that compost-induced SAR played a role in control of *Pythium* root rot. The significantly ($P \leq 0.05$) lower anthracnose severity (Table 2) observed on the second leaves of cucumber plants in compost-amended mixes strengthens this argument. Finally, increased peroxidase activity (Table 3) and enhanced peroxidase isozyme level (Fig. 1) in plants produced in the compost-amended mixes over that in the conducive peat mix are further evidence of the role of compost-induced SAR in control of these diseases.

The mechanisms by which these composts induced SAR remain unknown. Chemical as well as biological factors can induce this effect in plants (14,19,22,23,26,33–35,41). The high populations of biocontrol agents in the compost-amended mixes versus low populations in the conducive peat mix (3) suggest they play a role in SAR, because many of the bacterial strains present in the compost-amended mixes that induce suppression to *Pythium* root rot also induce SAR in plants (19,22,23,33–35,39,41). Furthermore, autoclaving destroys the suppressive effect of compost-amended mixes against *Pythium* root rot (6). Finally, the conducive peat mix, as mentioned above, is colonized predominantly by obligate oligotrophs and pleomorphic gram-positive bacteria that are unable to induce biological control (3,30) and, thus apparently, also are unable to induce SAR.

Many phytotoxic chemicals, including salicylic acid, can induce SAR (14,26,32). Alcohols and organic acids can occur in phytotoxic concentrations in immature composts (9) and, possibly, also in compost steepages (8,31,36). The composts used in this work, however, were cured well beyond the stability level that produces phytotoxic concentrations of such compounds (12). Allelopathic substances also may be present in composts prepared from tree barks (12,29). These products, however, are readily destroyed during composting (29) and, therefore, are thought to not have had a role in SAR induced by composts in cucumber in this work.

Nutritional factors affect the severity of many plant diseases (13). The concentration of plant nutrients used in the peat and compost mixes used in this work were adjusted to within optimum ranges for cucumber (6). Plants in both peat and compost-amended mixes developed at similar rates, and availability of essential plant nutrients was discounted as having a role in the disease control observed in this work.

The systemic protection of split-roots from *Pythium* root rot (Table 1) induced by compost was accompanied by increased peroxidase activity (Table 3) in leaf tissue. Acidic peroxidase is a putative

molecular marker of SAR in cucumber (11,24,27,28). Based on peroxidase isozyme activity in electrophoretic analysis (Fig. 1), however, the composts by themselves were weak inducers of SAR relative to SAR induced by prior inoculation with *C. orbiculare*. The obviously higher peroxidase isozyme activity in leaves of cucumber plants that were prior inoculated, as well as challenge-inoculated, and grown in compost compared to peat (Fig. 1) suggests that as yet unidentified factors in composts complemented the activity of *C. orbiculare* in induction of SAR. The interaction of compost and the pathogen appeared critical for rapid activation of SAR-associated gene expression in cucumber plants.

LITERATURE CITED

- Albert, F., and Anderson, A. J. 1987. The effect of *Pseudomonas putida* colonization on root surface peroxidase. *Plant Physiol.* 85:537-541.
- Boehm, M. J., and Hoitink, H. A. J. 1992. Sustainance of microbial activity in the potting mixes and its impact on severity of *Pythium* root rot of poinsettia. *Phytopathology* 82:259-264.
- Boehm, M. J., Madden, L. V., and Hoitink, H. A. J. 1993. Effect of organic matter decomposition level on bacterial species diversity and composition in relationship to *Pythium* damping-off severity. *Appl. Environ. Microbiol.* 59:4147-4179.
- Cheliak, W. M., and Pitel, J. A. 1984. Techniques for starch gel electrophoresis of enzymes from forest tree species. Pages 35-36 in: Inform. Rep. PI-X-42. Petawawa National Forestry Institute, Agriculture Canada, Chalk River, Ontario.
- Chen, W., Hoitink, H. A. J., and Madden, L. V. 1988. Microbial activity and biomass in container media for predicting suppressiveness to damping-off caused by *Pythium ultimum*. *Phytopathology* 78:1447-1450.
- Chen, W., Hoitink, H. A. J., and Schmitthenner, A. F. 1987. Factors affecting suppression of *Pythium* damping-off in container media amended with composts. *Phytopathology* 77:755-760.
- Craft, C. M., and Nelson, E. B. 1996. Microbial properties of composts that suppress damping-off and root rot of creeping bentgrass caused by *Pythium graminicola*. *Appl. Environ. Microbiol.* 62:1550-1557.
- Elad, Y., and Shtienberg, D. 1994. Effect of compost water extracts on grey mold (*Botrytis cinerea*). *Crop Prot.* 13:109-114.
- Garcia, C., Hernandez, T., Costa, F., and Pascual, J. A. 1992. Phytotoxicity due to the agricultural use of urban wastes. Germination experiments. *J. Sci. Food Agric.* 59:313-319.
- Hadar, Y., Mandelbaum, R., and Gorodecki, B. 1992. Biological control of soilborne plant pathogens by suppressive compost. Pages 79-83 in: *Biological Control of Plant Diseases*. E. S. Tjamos, G. C. Papavizas, and R. J. Cook, eds. Plenum Press, New York.
- Hammerschmidt, R., Nuckles, E. M., and Kuć, J. 1982. Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiol. Plant Pathol.* 20:73-82.
- Hoitink, H. A. J., Inbar, Y., and Boehm, M. J. 1991. Status of compost-amended potting mixes naturally suppressive to soilborne diseases of floricultural crops. *Plant Dis.* 75:869-873.
- Huber, D. M. 1989. Soilborne pathogens—Management of diseases with macro- and microelements: Introduction. Pages 1-8 in: *Soilborne Plant Pathogens: Management of Diseases with Macro- and Microelements*. A. W. Engelhard, ed. The American Phytopathological Society, St. Paul, MN.
- Kessmann, H., Staub, T., Hofmann, C., Maetzke, T., Herzog, J., Ward, E., Uknes, S., and Ryals, J. 1994. Induction of systemic acquired disease resistance in plants by chemicals. *Annu. Rev. Phytopathol.* 32:439-459.
- Ko, W., and Hora, F. K. 1971. A selective medium for the quantitative determination of *Rhizoctonia solani* in soil. *Phytopathology* 61:707-710.
- Kuć, J., and Richmond, S. 1977. Aspects of protection of cucumber against *Colletotrichum lagenarium* by *Colletotrichum lagenarium*. *Phytopathology* 67:533-536.
- Kuter, G. A., Nelson, E. B., Hoitink, H. A. J., and Madden, L. V. 1983. Fungal populations in container media amended with composted hardwood bark suppressive and conducive to *Rhizoctonia* damping-off. *Phytopathology* 73:1450-1456.
- Kwok, O. C. H., Fahy, P. C., Hoitink, H. A. J., and Kuter, G. A. 1987. Interactions between bacteria and *Trichoderma hamatum* in suppression of *Rhizoctonia* damping-off in bark compost media. *Phytopathology* 77:1206-1212.
- Liu, L., Kloepper, J. W., and Tuzun, S. 1995. Induction of systemic resistance in cucumber by plant growth-promoting rhizobacteria: Duration of protection and effect of host resistance on protection and root colonization. *Phytopathology* 85:1064-1068.
- Lumsden, R. D., Lewis, J. A., and Millner, P. D. 1983. Effect of composted sewage sludge on several soilborne plant pathogens and diseases.

- Phytopathology 73:1543-1548.
21. Mandelbaum, R., and Hadar, Y. 1990. Effects of available carbon source on microbial activity and suppression of *Pythium aphanidermatum* in compost and peat container media. *Phytopathology* 80:794-804.
22. Maurhofer, M., Hase, C., Meuwly, P., Métraux, J.-P., and Défago, G. 1994. Induction of systemic resistance of tobacco to tobacco necrosis virus by the root-colonizing *Pseudomonas fluorescens* strain CHA0: Influence of the *gacA* gene and pyoverdine production. *Phytopathology* 84:139-146.
23. Meera, M. S., Shivanna, M. B., Kageyama, K., and Hyakumachi, M. 1995. Responses of cucumber cultivars to induction of systemic resistance against anthracnose by plant growth promoting fungi. *Eur. J. Plant Pathol.* 101:421-430.
24. Rasmussen, J. B., Smith, J. A., Williams, S., Burkhart, W., Ward, E., Somerville, S. C., Ryals, J., and Hammerschmidt, R. 1995. cDNA cloning and systemic expression of acidic peroxidases associated with systemic acquired resistance to disease in cucumber. *Physiol. Mol. Plant Pathol.* 46:389-400.
25. Quarles, W., and Grossman, J. 1995. Alternatives to methyl bromide in nurseries—Disease suppressive media. *IPM Practitioner* 17:1-11.
26. Ryals, J., Uknes, S., and Ward, E. 1994. Systemic acquired resistance. *Plant Physiol.* 104:1109-1112.
27. Smith, J. A., and Hammerschmidt, R. 1988. Comparative study of acidic peroxidase associated with induced resistance in cucumber, muskmelon and watermelon. *Physiol. Mol. Plant Pathol.* 33:255-261.
28. Smith, J. A., Hammerschmidt, R., and Fulbright, D. W. 1991. Rapid induction of systemic resistance in cucumber by *Pseudomonas syringae* pv. *syringae*. *Physiol. Mol. Plant Pathol.* 38:223-235.
29. Still, S. M., Dirr, M. A., and Gartner, J. B. 1976. Phytotoxic effects of several bark extracts on mung bean and cucumber growth. *J. Am. Soc. Hortic. Sci.* 101:34-37.
30. Sugimoto, E. E., Hoitink, H. A. J., and Tuovinen, O. H. 1990. Oligotrophic pseudomonads in the rhizosphere: Suppressiveness to *Pythium* damping-off of cucumber seedlings (*Cucumis sativus* L.). *Biol. Fertil. Soils* 9:231-234.
31. Tränkner, A. 1992. Use of agricultural and municipal organic wastes to develop suppressiveness to plant pathogens. Pages 35-42 in: *Biological Control of Plant Diseases*. E. C. Tjamos, G. C. Papavizas, and R. J. Cook, eds. Plenum Press, New York.
32. Tuzun, S., Rao, M. N., Vogeli, U., Schardl, C. L., and Kuć, J. 1989. Induced systemic resistance to blue mold: Early induction and accumulation of β -1,3-glucanases, chitinases, and other pathogenesis-related proteins (b-proteins) in immunized tobacco. *Phytopathology* 79:979-983.
33. van Peer, R., Niemann, G. J., and Schippers, B. 1991. Induced resistance and phytoalexin accumulation in biological control of Fusarium wilt of carnation by *Pseudomonas* sp. strain WCS417r. *Phytopathology* 81:728-734.
34. Wei, G., Kloepper, J. W., and Tuzun, S. 1991. Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology* 81:1508-1512.
35. Wei, G., Kloepper, J. W., and Tuzun, S. 1996. Induced systemic resistance to cucumber diseases and increased plant growth by plant growth-promoting rhizobacteria under field conditions. *Phytopathology* 86:221-224.
36. Weltzien, H. C. 1992. Biocontrol of foliar fungal diseases with compost extracts. Pages 430-450 in: *Microbial Ecology of Leaves*. J. H. Andrews and S. S. Hirano, eds. Springer-Verlag, New York.
37. Wu, T., Boehm, M. J., Madden, L. V., and Hoitink, H. A. J. 1993. Sustained suppression of *Pythium* root rot: A function of the microbial carrying capacity and bacterial species composition. (Abstr.) *Phytopathology* 83:1365.
38. Ye, X. S., Pan, S. Q., and Kuć, J. 1990. Association of pathogenesis-related proteins and activities of peroxidase, β -1,3-glucanase and chitinase with systemic induced resistance to blue mold of tobacco but not to systemic tobacco mosaic virus. *Physiol. Mol. Plant Pathol.* 36:523-531.
39. Zdor, R. E., and Anderson, A. J. 1992. Influence of root colonizing bacteria on the defense responses of bean. *Plant Soil* 140:99-107.
40. Zhang, W., Dick, W. A., and Hoitink, H. A. J. 1994. Compost induced systemic resistance in cucumber plants to *Pythium* root rot and anthracnose. (Abstr.) *Phytopathology* 84:1138.
41. Zhou, T., and Paulitz, T. C. 1994. Induced resistance in the biocontrol of *Pythium aphanidermatum* by *Pseudomonas* spp. on cucumber. *J. Phytopathol.* 142:51-63.

Impacts of compost quality on plant disease severity

H.A.J. HOITINK, M.E. GREBUS AND A.G. STONE

1. Introduction

Soil organic matter decomposition level plays an important role in soilborne disease severity (Cook and Baker, 1983). Even though this has been recognised for centuries, management of diseases with amendments remains an art rather than a science. Lack of progress in this field in part has been due to our inability to characterise adequately biological energy availability in soil or the amendments. Recent breakthroughs in instrumentation, coupled with advances made in the analysis of microbial populations in soil, promise to solve this problem. Procedures are available for monitoring the stability of composts based on respirometry (Iannotti *et al.*, 1993). Furthermore non-destructive direct spectroscopic procedures are available that in principle allow characterisation of stability or biological energy availability in organic matter (Chen and Inbar, 1993; Inbar *et al.*, 1989). This review emphasises this literature on composting and how it relates to plant disease suppression.

During the 1960s, nurserymen across the United States explored the possibility of using tree bark as a peat substitute to reduce potting mix costs. Since that time, procedures for composting of various types of tree bark have been developed which avoid nitrogen immobilisation and allelopathy toxin problems associated with fresh bark (Hoitink and Kuter 1986; Hoitink *et al.*, 1991). During the past decade, similar procedures have been developed for composts produced from food industry and agricultural wastes, as well as from municipal wastes (Hoitink and Keener, 1993).

Early during the utilisation of bark composts, improved plant growth and decreased losses caused by phytophthora root rots were observed as side benefits by the nursery industry. In practice, control of such root rots with composts can be as effective as that obtained with fungicides (Hardy and Sivasithamparam, 1991; Hoitink *et al.*, 1991; Ownley and Benson, 1991). Therefore, the nursery industry relies heavily on composts for control of diseases caused by these soilborne plant pathogens.

Composts must be of consistent quality and maturity to be used successfully in biological control of diseases of horticultural crops, particularly if used in container media (Inbar *et al.*, 1993a). Variability in this quality parameter is one of the principal factors limiting its utilisation. Maturity is less important in ground bed or field agriculture as long as composts are applied well ahead of planting to allow for additional stabilisation. Lack of maturity frequently causes problems here as well, however.

En: D. Rosen *et al.* (eds): *Modern Agriculture and the Environment*, 361-369.
 © 1996 Kluwer Academic Publishers. Printed in Great Britain.

2. Fate of biocontrol agents during composting

The composting process is often divided into three phases. The first phase occurs during the first 24–48 hr as temperatures gradually rise to 40–50°C and sugars and other easily biodegradable substances are destroyed. During the second phase, when temperatures of 40–65°C prevail, cellulosic substances that are less biodegradable are destroyed. Lignins, the darker components in plant tissues, break down even more slowly but the rate varies among plant species. Thermophilic microorganisms predominate during this part of the process. Plant pathogens and seeds are killed by the heat generated during this high temperature phase (Bollen, 1993; Farrell, 1993). Compost piles must be turned frequently to expose all parts to high temperature and produce a homogeneous product free of pathogens and weed seeds.

The third or curing phase of composting begins as the concentration of readily biodegradable components in wastes declines. As a result, rates of decomposition and heat output and also temperatures decline. At this time, mesophilic microorganisms recolonise the compost from the outer low temperature layer into the pile. Humic substances accumulate in increasing quantities by this time. Mature composts consist largely of lignins, humic substances and microbial biomass, and have a dark colour (Chen and Inbar, 1993).

Unfortunately, beneficial as well as detrimental microorganisms are killed during the high temperature phase of composting. Therefore, suppression of pathogens and/or disease is largely induced during curing as biocontrol agents recolonise composts after peak heating. *Bacillus* spp., *Enterobacter* spp., *Flavobacterium balustinum*, *Pseudomonas* spp., other bacterial genera and *Streptomyces* spp., as well as *Penicillium* spp., a host of *Trichoderma* spp., *Gliocladium virens* and other fungi have been identified as biocontrol agents in composts (Chung and Hoitink, 1990; Hadar and Gorodecki, 1991; Hardy and Sivasithamparam, 1991; Hoitink and Fahy, 1986; Nelson *et al.*, 1983; Phae *et al.*, 1990).

The moisture content of compost critically affects the potential for mesophiles to colonise the substrate after peak heating. Dry composts (< 34% moisture, w/w) become colonized by fungi and are conducive to Pythium diseases, for example. The moisture content must be high enough (40–50%, w/w), so that bacteria can colonise the substrate after peak heating. Water must often be added to composts during curing to avoid this problem.

Variability in suppression of Rhizoctonia damping-off encountered in substrates amended with mature composts in part is due to random recolonisation after peak heating of compost by a microflora with efficacy against *R. solani*. Compost produced in the open near a forest, an environment that is high in microbial species diversity, is more consistently suppressive to Rhizoctonia diseases than the same produced in a partially enclosed facility where few microbial species survive heat treatment (Kuter *et al.*, 1983). Composts prepared from municipal sewage sludges and MSW are consistently conducive to *R. solani* because care is taken to kill faecal pathogens and parasites with heat exposure. As mentioned earlier, this process kills most beneficial microorganisms as well. These composts, although

naturally suppressive to *Pythium* diseases, have to be incubated for a month or more before they become naturally colonised due to chance by the right-specific microflora and, thus, become suppressive to *R. solani* (Kuter *et al.*, 1988). In field soil, several months may pass before suppression is induced (Lumsden *et al.*, 1983).

To solve the problem of variability in suppressiveness of compost to *R. solani* and other plant pathogens, specific fungal and bacterial inoculants have been developed, which when introduced into compost after peak heating, but before significant levels of recolonisation have occurred, induce consistent levels of suppression. Patents have been issued to the Ohio State University for this process (Hoitink, 1990). In Japan, Phae *et al.* (1990) isolated a *Bacillus* strain that induces predictable biological control in composts. It survives peak heating.

3. Mechanisms of biological control in composts

Two mechanisms of biological control, based on competition, antibiosis, hyperparasitism, and induced systemic resistance in plants, have been described for compost-amended substrates. Propagules of plant pathogens, including *Pythium* and *Phytophthora* spp., are suppressed through a mechanism known as 'general suppression' (Boehm *et al.*, 1993; Chen *et al.*, 1988a,b; Cook and Baker, 1983; Hardy and Sivasithamparam, 1991; Mandelbaum and Hadar, 1990). Many types of microorganisms present in compost-amended container media function as biocontrol agents for diseases caused by *Phytophthora* and *Pythium* spp. (Boehm *et al.*, 1993; Hardy and Sivasithamparam, 1991). Propagules of these pathogens, if inadvertently introduced into composts, do not germinate in response to nutrients released in the form of seed or root exudates. The high microbial activity and biomass caused by the 'general soil microflora' in compost-amended substrates, prevents germination of spores of the pathogen and infection of the host, presumably through microbiostasis (Chen *et al.*, 1988a; Mandelbaum and Hadar, 1990). Propagules of these pathogens remain dormant and are typically not killed if introduced in compost-amended soil (Chen *et al.*, 1988a; Mandelbaum and Hadar, 1990). An enzyme assay, that determines microbial activity based on the rate of hydrolysis of fluorescein diacetate, predicts suppressiveness of potting mixes to *Pythium* diseases (Boehm and Hoitink, 1992; Chen *et al.*, 1988a; Mandelbaum and Hadar, 1990). Similar information has been developed for soils on 'organic farms' where soilborne diseases are less prevalent (Workneh *et al.*, 1993).

The mechanism of biological control for *Rhizoctonia solani* in compost-amended substrates is different from that of *Pythium* and *Phytophthora* spp. A narrow group of microorganisms is capable of eradicating sclerotia of *R. solani*. This type of suppression we refer to as 'specific suppression' (Hoitink *et al.*, 1991). *Trichoderma* spp., including *T. hamatum* and *T. harzianum*, are the predominant hyperparasites recovered from composts prepared of lignocellulosic wastes (Kuter *et al.*, 1983; Nelson *et al.*, 1983). These fungi interact with various bacterial strains in biological control of *Rhizoctonia* damping-off (Kwok *et al.*, 1987). It is of interest that

Penicillium spp. are the predominant hyperparasites recovered from sclerotia of *Sclerotium rolfsii* in composted grape pomace, a waste high in sugar content (Hadar and Gorodecki, 1991). *Trichoderma* spp. were not recovered from this compost. The composition of the parent product, as expected, thus appears to have an impact on the microflora in composts active in biological control.

4. Role of organic matter decomposition level

The decomposition level of organic matter in compost-amended substrates has a major impact on disease suppression. For example, *R. solani* is highly competitive as a saprophyte (Garrett, 1962). It can utilise cellulose and colonise fresh wastes but not the low in cellulose mature compost (Chung *et al.*, 1988). *Trichoderma* isolates on the other hand, function as biocontrol agents of *R. solani*, and are capable of colonising fresh as well as mature compost (Nelson *et al.*, 1983). In fresh, undecomposed organic matter, biological control does not occur because both fungi grow as saprophytes and *R. solani* (the pathogen) remains capable of causing disease. presumably, synthesis of lytic enzymes involved in hyperparasitism of pathogens by *Trichoderma* (de la Cruz *et al.*, 1993) is repressed here due to high glucose concentrations in fresh wastes. The same may apply to antibiotic production. In mature compost, where concentrations of free nutrients are low (Chen *et al.*, 1988a), sclerotia of *R. solani* are killed by the hyperparasite and biological control prevails (Nelson *et al.*, 1983).

Because organic matter decomposition level is so important, composts must be stabilised adequately to reach that decomposition level where biological control is feasible. In practice, this occurs in composts (tree barks, yard wastes, etc.) that have been stabilised far enough not to induce phytotoxicity (due to lack of stability) nor immobilise nitrogen during plant growth, but are colonised by the appropriate specific microflora. Practical guidelines that define this critical stage of decomposition in terms of biological control are not yet available. Industry presently controls decomposition level by maintaining constant conditions during the entire process and adhering to a given time schedule.

Excessively stabilised organic matter, the opposite end of the decomposition scale, does not support adequate activity of biocontrol agents. As a result suppression is lacking and soilborne diseases are severe in highly mineralised soils, where humic substances are the predominant forms of organic matter (Workneh *et al.*, 1993).

The length of time that composts incorporated in soil support adequate levels of activity for biocontrol has not yet been determined. Presumably, it varies with soil temperature and the type of organic matter from which the compost was prepared. Loading rates and farming practices of course also play a role.

We have studied this 'carrying capacity' of soil organic matter in potting mixes prepared with sphagnum peat to bring a partial solution to this problem (Boehm and Hoitink, 1992, Boehm *et al.*, 1993). Sphagnum peat typically competes with compost as a source of organic matter in horticulture. Both the microflora and the

organic matter in peat itself can affect suppression of soilborne diseases. The literature on that effect is therefore briefly reviewed here.

Because of its stability, organic matter in sphagnum peat (resistance to decomposition) generally does not support high microbial activity. Dark, more decomposed sphagnum peat, harvested from a foot or greater depths in peat bogs, is low in activity and consistently conducive to *Pythium* and *Phytophthora* root rots (Boehm and Hoitink, 1992). On the other hand, light sources of sphagnum peat harvested from the surface of peat bogs, are less decomposed, have a higher microbial activity (FDA activity) and suppress disease. The suppressive effect of light peat to *Pythium* root rots is of short duration, however (Boehm and Hoitink, 1992; Tahvonen, 1982; Wolffhechel, 1988). Light peats, therefore, can be used most effectively for short production cycles, such as in plug and flat mixes used in the ornamentals industry. Composts have longer lasting effects (Boehm and Hoitink, 1992; Boehm *et al.*, 1993).

The rate of hydrolysis of fluorescein diacetate (FDA) predicts suppressiveness of compost-amended substrates and of peat mixes to *Pythium* root rot (Boehm and Hoitink, 1992). As FDA activity in suppressive light sphagnum peat substrates declines to less than $3.2 \mu\text{g FDA hydrolysed min}^{-1} \text{g}^{-1}$ dry weight mix, the population of *Pythium ultimum* increases and root rot develops. During this collapse in suppressiveness, bacterial species diversity in the rhizosphere or the edaphic substrate does not change. However, species composition is affected (Boehm *et al.*, 1993). A microflora typical of suppressive soils, which includes *Pseudomonas* spp. and other rod-shaped Gram negative bacteria as the predominant rhizosphere colonisers, is replaced by pleomorphic Gram-positive bacteria and putative oligotrophs (Wu *et al.*, 1993). Thus, the microflora of the conducive substrate resembles that of highly mineralised niches in soil (Kanazawa and Filip, 1986).

Non-destructive analysis of soil organic matter, utilising Fourier Transform infrared spectroscopy (FT-IR) and cross polarisation magic angle spinning - ^{13}C nuclear magnetic resonance spectroscopy (CPMAS- ^{13}C NMR), allows characterisation of biodegradable components of soil organic fractions (Chen and Inbar, 1993; Inbar *et al.*, 1989). CPMAS- ^{13}C NMR allows quantitative analysis of concentrations of readily biodegradable substances such as 'carbohydrates' (hemicellulose, cellulose, etc.) versus lignins and humic substances in soil organic matter (reviewed in Chen and Inbar, 1993). In a preliminary report, Wu *et al.* (1993) have shown that the concentration of 'carbohydrates' in a suppressive light peat mix declines as suppressiveness is lost. During the same time, bacterial genera capable of causing biological control are replaced by those that cannot. The concentration of 'carbohydrates' in the 'worn out' light peat has reached a final concentration not different from that in the conducive dark more decomposed peat. They also have shown that biocontrol agents inoculated into the more decomposed peat are not able to induce sustained biological control of *Pythium* root rot. Thus, biocontrol of this disease is determined by the 'carrying capacity' of the substrate which regulates species composition and activity and, thus in turn, the potential for sustenance of biological control.

5. Compost steepages for disease control

During the past decade, a series of projects have been published on the control of plant diseases of above ground plant parts with compost water extracts (Weltzien, 1992). The steepages are prepared by soaking mature composts in water (still culture; 1:1, w/w) for 7–10 days. The steepage is filtered and then sprayed on plants. Efficacy varies with composts, crops and the disease under question. Sackenheim (1993), utilising plate counting procedures, has shown that aerobic microorganisms predominate in these steepages. The microflora includes strains of bacteria and isolates of fungi known as biocontrol agents. A number of enrichment strategies, that included nutrients as well as microorganisms, were developed to improve efficacy of the steepages. Disease control induced by compost steepages has in the past been attributed in part to systemic acquired resistance induced in plants by microbes present in the extracts. The recent work by Sackenheim (1993) on grape does not support this assumption, however.

A factor that has been entirely overlooked but could play a role in efficacy of steepages is the condition of organic matter and the associated microflora in the soil in which treated plants are produced. Soils naturally suppressive to soilborne plant pathogens (e.g. compost-amended soils) harbour active populations of biocontrol agents (Boehm *et al.*, 1993) which could induce protection to foliar pathogens in the leaves of plants (Maurhofer *et al.*, 1994; Wei *et al.*, 1991). Thus, further research may show that plants produced in suppressive soils not only have less root rot but could also be less prone to attack by vascular wilt and foliar pathogens and respond better to compost steepages.

6. Disease suppression – future outlook

Success in biological control of diseases caused by *Fusarium*, *Pythium* and *Phytophthora* spp. is possible only if all factors involved in the production of composts are defined and kept constant. Most composts cannot meet these criteria because emphasis on quality control in general is still inadequate. Therefore, composted bark and light sphagnum peat remain the principal organic components used for the preparation of potting mixes or soils naturally suppressive to soilborne plant pathogens. Composted manures, yard and food wastes, are steadily gaining in popularity, and offer the same potential (Gorodecki and Hadar, 1990; Grebus *et al.*, 1994; Inbar *et al.*, 1993b; Marugg *et al.*, 1993; Schüller *et al.*, 1993).

The general suppression phenomenon, based on present concepts, at best covers those diseases caused by pathogens suppressed through microbiostasis. Microbial activity in a mix, based on the rate of hydrolysis of fluorescein diacetate, is one procedure that now can be used effectively to determine suppressiveness to *Pythium* and *Phytophthora* root rots. This procedure by itself, however, does not predict how long the effect will last. The potential for biodegradable carbon in a mix to support an active and effective microbial biomass determines that phenomenon. CPMAS-¹³CNMR spectroscopy predicts this property for peat. The same basic information

will have to be developed for composts and other sources of organic amendments. Predictive utilisation of composts for disease control will remain an art until more quantitative guidelines are developed.

Controlled inoculation of composts with biocontrol agents is a procedure that must be developed on a commercial scale to induce consistent levels of suppression to pathogens such as *R. solani* (Grebus *et al.*, 1993; Hoitink *et al.*, 1991; Phae *et al.*, 1990). This new field of biotechnology is in its infancy, however. Major research and development efforts will need to be directed into this approach to disease control. Recycling through composting is increasingly being chosen as the preferred strategy for waste treatment. This also applies to farm manures (Rynk, 1992). For this reason, composts are becoming available in greater quantities. Peat, on the other hand, is a limited resource that cannot be recycled. Future opportunities for both natural and controlled-induced suppression of soilborne plant pathogens, using composts as the food stuff for biocontrol agents, therefore appear bright.

Acknowledgements

Salaries and research support provided by State and Federal Funds appropriated to the Ohio Agricultural Research and Development Center, The Ohio State University and by Grant No. US-2196-92 from BARD, the United States-Israel Binational Agricultural Research and Development Fund.

References

- Boehm M J and Hoitink H A J 1992 Sustenance of microbial activity and severity of *Pythium* root rot of poinsettia. *Phytopathology* 82, 259-264.
- Boehm M J, Madden L V and Hoitink H A J 1993 Effect of organic matter decomposition level on bacterial species diversity and composition in relationship to *Pythium* damping-off severity. *Appl. Environ. Microbiol.* 59, 4171-4179.
- Bollen G J 1993 Factors involved in inactivation of plant pathogens during composting of crop residues. In *Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects*. Eds H A J Hoitink and H M Keener, pp 301-318. Renaissance Publications, Worthington, OH.
- Chen Y and Inbar Y 1993 Chemical and spectroscopical analyses of organic matter transformations during composting in relation to compost maturity. In *Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspect*. Eds H A J Hoitink and H M Keener. pp 551-600. Renaissance Publications, Worthington, OH.
- Chen W, Hoitink H A J, Schmitthenner A F and Tuovinen O H 1988a The role of microbial activity in suppression of damping-off caused by *Pythium ultimum*. *Phytopathology* 78, 314-322.
- Chen Y, Hoitink H A J and Madden L V 1988b Microbial activity and biomass in container media predicting suppressiveness to damping-off caused by *Pythium ultimum*. *Phytopathology* 78, 1447-1450.
- Chung Y R and Hoitink H A J 1990 Interactions between thermophilic fungi and *Trichoderma hamatum* in suppression of *Rhizoctonia* damping-off in a bark compost-amended container medium. *Phytopathology* 80, 73-77.
- Chung Y R, Hoitink H A J and Lipps P E 1988 Interactions between organic-matter decomposition level and soilborne disease severity. *Agric. Ecosys. Environ.* 24, 183-193.
- Cook R J and Baker K F 1983 *The Nature and Practice of Biological Control of Plant Pathogens*. Am. Phytopathol. Soc., St Paul, MN.
- de la Cruz J, Rey M, Lora J M *et al.* 1993 Carbon source control on β -glucanases, chitinase and chitinase from *Trichoderma harzianum*. *Microbiology* 159, 316-322.

- Farrell J B 1993 Fecal pathogen control during composting. In Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects. Eds H A J Hoitink and H M Keener. pp 282-300. Renaissance Publications, Worthington, OH.
- Garrett, S D 1962 Decomposition of cellulose in soil by *Rhizoctonia solani* Kuhn. Trans. Br. Mycol. Soc. 45, 114-120.
- Gorodecki B and Hadar Y 1990 Suppression of *Rhizoctonia solani* and *Sclerotium rolfsii* in container media containing composted separated cattle manure and composted grape marc. Crop Protection 9, 271-274.
- Grebus M E, Feldman K A, Musselman C A and Hoitink H A J 1993 Production of biocontrol agent fortified compost-amended potting mixes for predicatable disease suppression. Phytopathology 83, 1406 (Abstract).
- Grebus M E, Watson M E and Hoitink H A J 1994 Biological, chemical and physical properties of composted yard trimmings as indicators of maturity and plant disease suppression. Compost Sci. Util. 1, 57-71.
- Hadar Y and Gorodecki B 1991 Suppression of germination of sclerotia of *Sclerotium rolfsii* in compost. Soil. Biol. Biochem. 23, 303-306.
- Hardy G E St J and Sivasithamparan K 1991 Suppression of *Phytophthora* root rot by a composted *Eucalyptus* bark mix. Aust. J. Bot. 39, 153-159.
- Hoitink H A J 1990 Production of disease suppressive compost and container media, and microorganism culture for use therein. US Patent 4960348. Feb. 13, 1990.
- Hoitink H A J and Fahy P C 1986 Basis for the control of soilborne plant pathogens with composts. Ann. Rev. Phytopathol. 24, 93-114.
- Hoitink H A J and Keener H M (Eds) 1993 Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects. Renaissance Publications, Worthington, OH.
- Hoitink H A J and Kuter G A 1986 Effects of composts in growth media on soilborne plant pathogens. In The Role of Organic Matter in Modern Agriculture. Eds Y Chen and Y Avnimelech. pp 289-306. Martinus Nyhoff Publishers, Dordrecht and The Netherlands.
- Hoitink H A J, Inbar Y and Boehm M J 1991 Status of composted-amended potting mixes naturally suppressive to soilborne diseases of floricultural crops. Plant Dis. 75, 869-873.
- Hoitink H A J, Boehm M J and Hadar Y 1993 Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. In Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects. Eds H A J Hoitink and H M Keener. pp 601-621. Renaissance Publications, Worthington, OH.
- Iannotti D A, Pang T, Toth B L, Elwell D L, Keener H M and Hoitink H A J 1993 A quantitative respirometric method for monitoring compost stability. Compost Sci. Util. 1, 52-65.
- Inbar Y, Chen Y and Hadar Y 1989 Solid-state carbon-13 nuclear magnetic resonance and infrared spectroscopy of composted organic matter. Soil Sci. Soc. Am. J. 53, 1695-1701.
- Inbar Y, Chen Y and Hoitink H A J 1993a Properties for establishing standards for utilization of composts in container media. In Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects. Eds H A J Hoitink and H M Keener. pp 668-694. Renaissance Publications, Worthington, OH.
- Inbar Y, Hadar Y and Chen Y 1993b Recycling of cattle manure: The composting process and characterization of maturity. J. Environ. Qual. 22, 857-863.
- Kanazawa S and Filip Z 1986 Distribution of microorganisms, total biomass, and enzyme activities in different particles of brown soils. Microb. Ecol. 12, 205-215.
- Kuter G A, Nelson E B, Hoitink H A J and Madden L V 1983 Fungal populations in container media amended with composted hardwood bark suppressive and conducive to *Rhizoctonia* damping-off. Phytopathology 73, 1450-1456.
- Kuter G A, Hoitink H A J and Chen W 1988 Effects of municipal sludge compost curing time on suppression of *Pythium* and *Rhizoctonia* diseases of ornamental plants. Plant Dis. 72, 751-756.
- Kwok O C H, Fahy P C, Hoitink H A J and Kuter G A 1987 Interactions between bacteria and *Trichoderma hamatum* in suppression of *Rhizoctonia* damping-off in bark compost media. Phytopathology 77, 1206-1212.
- Lumsden R D, Lewis J A and Millner P D 1983 Effect of composted sewage sludge on several soilborne pathogens and diseases. Phytopathology 73, 1543-1548.
- Mandelbaum R and Hadar Y 1990 Effects of available carbon source on microbial activity and suppression of *Pythium aphanidermatum* in compost and peat container media. Phytopathology 80, 794-804.
- Marugg C, Grebus M E, Hansen R C, Keener H M and Hoitink H A J 1993 A kinetic model of the yard waste composting process. Compost Sci. Util. 1, 38-51.

- Maurhofer M, Hase C, Meuwly P, Métraux, J-P, and Défago G 1994 Induction of systemic resistance of tobacco to tobacco necrosis virus by the root-colonizing *Pseudomonas fluorescens* strain CHAO: Influence of the *gac*, a gene and of pyoverdine production. *Phytopathology* 84, 139-146.
- Nelson E B, Kuter G A and Hoitink H A J 1983 Effects of fungal antagonists and compost age on suppression of *Rhizoctonia* damping-off in container media amended with composted hardwood bark. *Phytopathology* 3, 1457-1462.
- Ownley B H and Beason D M 1991 Relationship of matric water potential and air-filled porosity of container media to development of *Phytophthora* root rot of rhododendron. *Phytopathology* 81, 936-941.
- Phae C G, Saski M, Shoda M and Kubota H 1990 Characteristics of *Bacillus subtilis* isolated from composts suppressing phytopathogenic microorganisms. *Soil Sci. Plant Nutr.* 36, 575-586.
- Rynk R (Ed.) 1992 On-Farm Composting Handbook. Coop. Ext., NE Reg. Agric. Eng. Serv., Ithaca, NY 14853-5701. NRAES-54.
- Sackenheim R 1993 Untersuchungen über Wirkungen von wässrigen, mikrobiologisch aktiven Extracten aus kompostierten Substraten auf den Befall der Weinrebe (*Vitis vinifera*) mit *Plasmopora viticola*, *Uncinula necator*, *Botrytis cinerea* und *Pseudopezizicola tracheiphila*. PhD thesis. Rheinische Friedrich-Wilhelms Universität, Bonn, Germany.
- Schiller C, Pikny J, Nasir M and Vogtmann H 1993 Effects of composted organic kitchen and garden waste on *Mycosphaerella pinodes* (Berk, et Blox) Vestergr., causal organism of foot rot on peas (*Pisum sativum* L.) *Biol. Agric. and Hort.* 9, 353-360.
- Tahvonen R 1982 The suppressiveness of Finnish light coloured Sphagnum peat. *J. Sci. Agric. Soc. Finl.* 54, 345-356.
- Wei G, Kloepper J W and Tuzun S 1991 Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology* 81, 1508-1512.
- Weltzien H C 1992 Biocontrol of foliar fungal diseases with compost extracts. In *Microbial Ecology of Leaves*. Eds J H Andrews and S Hirano. pp 430-450. Brock Springer Series in Contemporary Bioscience. BSNB 0387-97579-9.
- Wolffhechel H 1988 The suppressiveness of Sphagnum peat to *Pythium* spp. *Acta Hort.* 221, 217-222.
- Workneh F, Van Bruggen A H C, Drinkwater L E and Sherman C 1993 Variables associated with a reduction in corky root and *Phytophthora* root rot of tomatoes in organic compared to conventional farms. *Phytopathology* 83, 581-589.
- Wu T, Boehm M J, Madden L V and Hoitink H A J 1993 Sustained suppression of *Pythium* root rot: A function of microbial carrying capacity and bacterial species composition. *Phytopathology* 83, 1406 (Abstract).

Dynamics of *Pythium* populations, microbial biomass, microbial activity, and organic matter characterization during the collapse of suppression of *Pythium* root rot .
A.G. Stone, D.Y. Han, M.M. Wander, S.J. Traina and H.A.J. Hoitink.
Phytopath. 86:S53 (abstract).

Abstract:

Temporal relationships between particulate organic matter decomposition level and biological control of *Pythium* root rot caused by *Pythium ultimum* were investigated in a composted cow manure-amended sand. Suppressiveness to root rot was sustained over a 9-mon period in a 15% compost/sand mix although *P. ultimum* populations began to increase at 7-mon. As root rot developed and progressed, the population of *P. ultimum* increased further, while microbial biomass (as determined by phospholipid phosphate) and microbial activity (as determined by FDA hydrolysis) declined. Proportions of actinomycetes declined, while proportions of Gram negative bacteria and fungal and protozoan populations increased (as determined by PLFA analysis). The quantity of particulate organic matter declined at a constant rate over time. Peaks (from DR-FTIR spectroscopy) representing plant-derived polysaccharides (hemicellulose, cellulose) and lignin declined in intensity, while peaks representing microbe-derived polysaccharides and non-lignin aromatics increased in intensity.

Introduction:

Nursery growers have recognized for decades that organic components utilized in container media vary in their abilities to sustain biological control of root rots (Table 1) but the characteristics of organic materials which confer suppressiveness have not been identified.

Organic matter-supported suppression of *Pythium* root rot occurs through the activities of many different biocontrol microorganisms interacting with both the pathogen and the plant host. These activities include antibiotic production, nutrient competition and induction of plant resistance responses (3). Interestingly, recent work suggests that cellulases produced by *Trichoderma* spp. saprophytes may turn on host defense responses (13), a biocontrol activity tied directly to substrate chemistry.

Peat loses suppressiveness to *Pythium* root rot as the peat decomposes and the microbial community shifts in composition from species active in biocontrol (eg. *Pseudomonas* spp., *Pantoea* spp., *Flavobacterium* spp., etc) to those inactive (eg. *Arthrobacter*, oligotrophs, etc.). At the same time, FDA hydrolysis (a measure of microbial extracellular hydrolase activity and therefore substrate availability) and the carbohydrate content of the peat declines (as determined by CP MAS ^{13}C -NMR). This has led to the conclusion that carbohydrate content in peat may play a role in regulating the composition of the microbial community and its activities relative to the biological control of *Pythium* root rot (3).

Some of these trends have been documented in field agriculture. The rate of FDA hydrolysis was negatively related to the severity of *Phytophthora* root rot in a decomposing straw mulch (15), *Pythium* damping-off in creeping bentgrass (5) and the severity of *Phytophthora* root rot and corky root in tomato in organic and conventionally managed tomato fields (14). Suppressiveness to root rots has been positively related to fungal populations (15) and fluorescent pseudomonad and actinomycete populations (14).

Only one study in peat-based soilless mixes (2) has attempted to directly relate organic matter chemistry to suppressiveness to root rots. Suppressiveness to corky root in the field has been indirectly related to organic matter chemistry through FDA hydrolysis and nitrogen mineralization potential (6).

In this study our goal was to identify and quantify shifts in compost chemistry most closely related to the decline in suppressiveness to *Pythium* root rot and relate these events to shifts in *Pythium* populations, microbial community composition, biomass and activity.

Methods:

Preparation of sand mixes: Sawdust-bedded separated cow manure was composted in outdoor windrows and formulated into sand mixes with 15% compost (v/v). A 20% sphagnum peat (H4 on the von Post decomposition scale) sand mix consistently conducive to root rot was used as a control.

Disease suppression bioassay: Suppressiveness to *Pythium* damping-off was monitored with a cucumber seedling bioassay utilizing *Pythium ultimum* as the pathogen(4).

Microbial activity, bioamass and community composition: Hydrolysis of fluorescein diacetate was monitored as before (1). Total bacteria and fluorescent pseudomonads were monitored by dilution plating onto 0.1 TSA and King's B, respectively. The population of *P. ultimum* was monitored by dilution plating on a *Pythium* selective medium (1). Microbial biomass was measured by a phospholipid phosphate method (7) and microbial community composition was determined by Microbial Insights (Nashville, TN) by phospholipid fatty acid (PLFA) analysis (11).

Organic matter characterization: Compost/sand mixes were sampled at potting (suppressive to *Pythium* root rot) and at 14 mon (conductive). Organic matter (OM) was decanted from sand mixes, oven-dried and weighed to determine total particulate OM. Dried OM fractions were ground, mixed with KCl and characterized by diffuse reflectance FT-IR spectroscopy as in (12). Spectral sections from 1835-865 cm^{-1} were baseline-corrected to absorbance 0.00 at 1835 cm^{-1} and 0.99 (the mean of absorbance values) at 865 cm^{-1} and sections from 3015 - 2805 cm^{-1} (aliphatics) were baseline-corrected to absorbances 0.00 and -0.015. Six replicate spectra were averaged for each treatment and the resulting average spectra were subtracted to produce a single subtraction spectrum.

Twelve spectra (6 per treatment) (1750-750 cm^{-1} , 187 wavenumbers; 3015-2805 cm^{-1} , 56 wavenumbers) were analysed by principal component analysis (PCA) to partition total variability into fewer variables and to select peaks contributing highly to treatment differences. Scatter plots of PC scores were constructed to determine which PCs best described the variability between treatment groups. PC scores were analysed by discriminant analysis to verify cluster separation and t-tests to determine significance of differences between treatment groups.

Results:

The compost mix began to decline in suppressiveness at day 360 and at day 426 had declined to the same low level as the peat control. Suppressiveness never declined to a fully conducive level (disease severity rating 3.0-4.0) (Table 2).

Populations of *P. ultimum* began to increase after day 200 in the H4 peat mix and increased dramatically between days 280 and 370. Populations of *P. ultimum* in the 15% compost mix increased after day 280 but at a slower rate, and populations in a 25% compost mix increased even more slowly (Fig. 1). Pathogen population data could not be taken after day 370 because the host plant died.

Total particulate organic matter declined at a constant rate. The rate of FDA hydrolysis, while extremely variable, also declined (Fig. 2).

Microbial biomass declined from 19,257 picomoles PLFA gdw soil mix⁻¹ (3.85×10^8 cells gdw sand mix⁻¹) at potting to 11,337 picomoles (2.27×10^8 cells gdw⁻¹) 8 months after potting, and remained constant thereafter. Prokaryotic biomass declined from 17,282 to 9,740 picomoles

PLFA gdw^{-1} at 8 months and to 8,739 picomoles at 14 months. Eukaryotic biomass declined from 1,971 picomoles PLFA gdw^{-1} to 1,597 picomoles at 8 months and 14 months. The prokaryotic to eukaryotic ratio declined from 9:1 at potting to 6:1 at 8 and 14 months.

The proportion of Gram negative bacteria (monoenoics) increased, while the proportion of Gram positives (terminal branched saturates) remained constant. The proportion of mid-chain branched saturates, which represent sulfate reducers, actinomycetes and certain other Gram positive bacteria, declined. The proportions of the protozoan PLFA 20:4w6 and the fungal PLFA 18:2w6 increased, while proportions of both the actinomycete PLFA 10me18:0 and PLFA i17:0 (certain Gram positive bacteria) declined (data not shown).

Dramatic changes in chemistry occurred during the composting process including reductions in the concentrations of aliphatics, amorphous and crystalline cellulose, hemicellulose, (1-3)-glucans and lignin (Fig. 3). In contrast, changes over the course of the experiment in the chemistry of the particulate organic matter removed from sand mixes were of much smaller magnitude, as shown by smaller differences in peak heights relative to those seen during the composting process (Fig. 4).

Principal component analysis of the spectral region $1750\text{-}775\text{ cm}^{-1}$ clearly separated spectra into treatment groups (Fig. 7). PCs 1-5 explained 52, 18, 14, 7 and 5% of the total variability, respectively. Discriminant analysis with cross validation 100% correctly grouped all replicates into two treatment groups by analysis of PC1 scores only. PC1 and PC2 scores of treatment groups were significantly different ($P=0.003$ and 0.09 , respectively). Aliphatics region spectra were not separated into treatment groups by PCA (data not shown).

The subtraction spectrum and plots of PC coefficients yielded highly similar profiles of peaks differing in intensity among suppressive and conducive spectral groups (Figs. 5 and 6). All peaks in Table 3 were identified both in the subtraction spectrum (Fig 5) and in coefficient plots of PC1 and PC2 (Fig. 6).

Peaks declining in intensity as suppressiveness declined represent hemicellulose, (1-3)-glucans (1252 cm^{-1}), crystalline and amorphous cellulose, lignin, and amides I and II. Peaks increasing in intensity represent non-cellulosic polysaccharides, (1-3)-glucans (1075 cm^{-1}), amides V, aromatics, and peaks (912 cm^{-1} and 985 cm^{-1}) seen in spectra of fungal structural glucans lutean and pustulan (Table 3).

Changes in constituent concentration could not be quantified because peak intensities could not be related to a known constituent concentration. Peak intensities at different wavenumbers cannot be related because each chemical constituent has a unique molar extinction coefficient at each wavenumber which is proportional to peak intensity.

In conclusion, microbial biomass had declined to 59% by day 245 after potting and then remained constant. *Pythium* populations began to increase slowly at day 250 and then more quickly between days 300 and 350. Suppressiveness only began to decline between days 375 and 426. Both total particulate organic matter and rate of FDA hydrolysis declined over time, but the variability in the FDA data makes a comparison of the two trends difficult (Fig. 8). There was a shift in microbial community composition and substrate chemistry from beginning to end of the experiment.

Discussion and Conclusions:

Overall, trends seen as suppressiveness declines in the compost mix are similar to those seen in the peat system (3): reductions in FDA activity and microbial biomass, an increase in *P. ultimum* population, and a shift in microbial community composition and substrate chemistry

The positive relationship between FDA activity and suppressiveness to root rot reported in field studies (5, 6, 14, 15) and in potting mixes (1) is also seen in this study.

A shift in community composition occurred as suppressiveness declined as in previous studies (3, 14, 15). However, specific results of this study cannot be compared directly to previous studies because of differences in methodology (total extractable PLFAs vs. culturable organisms)

Plant-derived polysaccharides declined in concentration as suppressiveness declined, but other polysaccharides increased in concentration. Future work will determine whether total carbohydrates declined. Substrate bioavailability declined as determined by the rate of FDA hydrolysis. Additional spectral data on particulate organic matter at 4, 8 and 12 months after potting should better relate changes in substrate chemistry over time to the decline in suppressiveness and changes in FDA activity

Wood-rot fungi produce extracellular polysaccharides/lignin complexes as lignin is degraded, and spectra of these are similar to the wood-bulked compost spectra in this study (9). The appearance of bands found in fungal wall glucans, the increase in intensity of (1-3)-glucan band (1075 cm^{-1}) and the increase in proportion of marker fungal PLFAs suggest that polysaccharides increasing in intensity as suppression declines may be of microbial origin.

References:

1. Boehm, M.J., and H.A.J. Hoitink, 1992. Sustenance of microbial activity in potting mixes and its impact on severity of *Pythium* root rot of poinsettia. *Phytopathology* 82:259-264.
2. Boehm, M.J., L.V. Madden, and H.A.J. Hoitink, 1993. Effect of organic matter decomposition level on bacterial species diversity and composition in relationship to *Pythium* damping-off severity. *Appl. Environ. Microbiol.* 59:4171-4179.
3. Boehm, M.J., T. Wu, A.G. Stone, B. Kraakman, D.A. Iannotti, G.E. Wilson, L.V. Madden and H.A.J. Hoitink, 1996. CPMAS ^{13}C -NMR spectroscopic characterization of soil organic matter relative to culturable species composition and sustained biological control of *Pythium* root rot. *Appl. Environ. Microbiol.*, in press.
4. Chen, W., H.A.J. Hoitink and L.V. Madden, 1988. Microbial activity and biomass in container media for predicting suppressiveness to damping-off caused by *Pythium ultimum*. *Phytopathology* 78:1447-1450.
5. Craft, C. M. and E.B. Nelson, 1996. Microbial properties of compost that suppress damping-off and root rot of creeping bentgrass caused by *Pythium graminicola*. *Appl. Env. Microbiol.* 62:1550-1557.
6. Drinkwater, L.E., D.K. Letourneau, F. Workneh, A.H.C. van Bruggen and C. Shennan, 1995. Fundamental differences between conventional and organic tomato agroecosystems in California. *Ecol. Applic.* 5:1098-1112.
7. Findlay, R.H., G.M. King, and L. Watling, 1989. Efficacy of phospholipid analysis in determination of microbial biomass in sediment. *Appl. Environ. Microbiol.* 55:2888-2893.
8. Gilbert, C., S. Kokot and U. Meyer, 1993. Application of DRIFT spectroscopy and chemometrics for the comparison of cotton fabrics. *Appl. Spectr.* 47:741-748.
9. Gutierrez, A., P. Bocchini, G.C. Galletti and A.T. Martinez, 1996. Analysis of lignin-polysaccharide complexes formed during grass lignin degradation by cultures of *Pleurotus* species. *Appl. Environ. Microbiol.* 62:1928-1934.
10. Stevenson, F.J., 1982. Humus chemistry. John Wiley and Sons, NY.
11. Tunlid, A. and D.C. White, 1991. Biochemical analysis of biomass, community structure, nutritional status, and metabolic activity of the microbial community in soil. In: Bollag J-M. and Stotzky, G. (eds.) *Soil Biochemistry* 7:229-262.
12. Wander, M.M. and S.J. Traina, 1996. Organic matter fractions from organically and conventionally managed soils: II. Characterization of composition. *Soil Sci. Soc. Am. J.* 60:1087-1095.
13. Wang, H. and R.W. Jones, unpublished data. Analyzing the role of a cellulose binding domain in fungal-plant interactions. Poster, American Phytopathological Society Meetings. August, 1996.

14. Workneh, F. and A.H.C. van Bruggen, 1994. Microbial density, composition, and diversity in organically and conventionally managed rhizosphere soil in relation to suppression of corky root of tomatoes. *Appl. Soil Ecol.* 1:219-230.
15. You, M.P. and K. Sivasithamparan, 1994. Hydrolysis of fluorescein diacetate in an avocado plantation mulch suppressive to *Phytophthora cinnamomi* and its relationship with certain biotic and abiotic factors. *Soil Biol. Biochem.* 26:1355-1361.

DR-FTIR Peak Assignment References

- Akin, D.E., D.S. Himmelsbach, R.T. Carl, W.W. Hanna and F.E. Barton, 1993. Mid-infrared microspectroscopy to assess lignin in plant tissues related to digestability. *Agron. J.* 85:171-175.
- Baes, A.U. and P.R. Bloom, 1989. Diffuse reflectance and transmission Fourier transform infrared (DRIFT) spectroscopy of humic and fulvic acids. *Soil Sci. Soc. Am. J.* 53:695-700.
- Evans, P.A., 1991. Differentiating "hard" from "soft" woods using Fourier transform infrared and Fourier transform Raman spectroscopy. *Spectr. Acta* 47:1441-1447.
- Gerasimowicz, W.V., D.M. Byler and H. Susi, 1986. Resolution-enhanced FT-IR spectra of soil constituents: humic acid. *Appl. Spectr.* 40:504-507.
- Gilbert, C., S. Kokot and U. Meyer, 1993. Application of DRIFT spectroscopy and chemometrics for the comparison of cotton fabrics. *Appl. Spectr.* 47:741-748.
- Guo, Y., J.J. Renton and J.H. Penn, 1996. FTIR microspectroscopy of particular liptinite-(lopinite-) rich, Late Permian coals from Southern China. *Int. J. Coal. Geol.* 29:187-197.
- Gutierrez, A., P. Bocchini, G.C. Galletti and A.T. Martinez, 1996. Analysis of lignin-polysaccharide complexes formed during grass lignin degradation by cultures of *Pleurotus* species. *Appl. Environ. Microbiol.* 62:1928-1934.
- Mathlouthi, M. and J.L. Koenig, 1986. Vibrational spectra of carbohydrates. *Adv. Carboh. Chem. Biochem.* 44:7-89.
- Michell, A.J. and G. Scurfield, 1970. An assessment of infrared spectra as indicators of fungal cell wall composition. *Aust. J. Biol. Sci.* 23:345-360.
- Painter, P.C., R.W. Snyder, M. Starsinic, M.M. Coleman, D. W. Kuehn and A. Davis, 1981. Concerning the application of FT-IR for the study of coal: a critical assessment of band assignments and the application of spectral analysis programs. *Appl. Spectr.* 35:475-489.
- Southerland, I.W., 1990. *Biotechnology of microbial polysaccharides.* Cambridge Univ. Press.
- Stevenson, F.J., 1982. *Humus chemistry.* John Wiley and Sons, NY.
- Stewart, D., H.M. Wilson, P.J. Hendra and I.M. Morrison, 1995. Fourier transform infrared and Raman spectroscopic study of biochemical and chemical treatments of oak wood (*Quercus rubra*) and barley (*Hordeum vulgare*) straw. *J. Agric. Food Chem.* 43:2219-2225.
- Stewart, D. and I.M. Morrison, 1992. FT-IR spectroscopy as a tool for the study of biological and chemical treatments of barley straw. *J. Sci. Food Agric.* 60:431-436.

Table 1.

BIOLOGICAL CARRYING CAPACITY*

Relative to the biocontrol of *Pythium* root rot;

based on grower observations and laboratory measurements

Hardwood bark compost	2-3 yr
Separated cow manure compost	1-2 yr
Pine bark compost	6-12 mo
Light peat (H2-H3 on the von Post decomposition scale)	2-10 mo
Dark peat (H4 on the von Post decomposition scale)	0-1 wk

*sensu Boehm et al, 1993

Table 2: Decline in Suppressiveness to *Pythium Damping-Off* of Cucumber in Compost- and Peat-Amended Sand Mixes.

	Time after potting (days)				
	15	186	274	375	426
15% compost mix¹	1.4²	1.4	1.0	1.5	2.4
20% H4 peat mix	2.4	2.2	3.8	3.1	2.5
LSD_{0.05} = 0.70					

¹ Infested with 0.5g soil inoculum of *Pythium ultimum* 21 l per liter mix (mean disease severity in controls ranged from 1.0-1.3)

² Mean damping-off severity, based on 8 cucumber seeds per pot (n=5), determined 10 days after planting. Disease severity scale; 1=symptomless, 2=diseased but not damped-off, 3=post-emergence damping-off, 4=pre-emergence damping-off.

Table 3. Peaks Changing in Intensity as Suppression Declines*

Peaks decreasing in intensity		Peaks increasing in intensity	
cm ⁻¹	Assignment	cm ⁻¹	Assignment
1735	carbonyls; hemicelluloses and lignins	1600	aromatic ring str. assoc. w/ methoxyl str.
1652	C=O str., amide I	1595	aromatic ring str. assoc. with C-O str.
1540-1520	N-H bend; C-N str., amide II	1270	C-OH stretch, phenolic OH
1460	polar asymm. arom. ring stretch, lignin	1075	(1-3)-glucan linkage
1437	1430 - CH ₂ wagging	1033	C-O-C stretch, non-cellulosic polysacch. and/or aromatic-associated methoxy grp.
1373	C-H def. str., (1-3)-glucan linkage	1009	aliphatic ethers, alcohols
1311	syringyl units	985	found in fungal glucan spectra
1252	(1-3)-glucan linkage	912	found in fungal glucan spectra
1200	O-H in-plane bending	797	arom. C-H (2 or 3 adj. H out-of-pl. bend)
1170	C-(CH ₃) ₂ skeletal stretch	775	aromatic C-H out-of-pl. bend
1161	antisymm. brdg. C-O-C, cellulose	695	amide V (N-H oop bend)
1098	principal crystalline cellulose absorb.		
900	antisym. oop ring str., amorph. cellulose		

*for references see handout

Fig. 1. Changes in Populations of *Pythium ultimum* in Compost- and Peat-Amended Sand Mixes

Fig. 2. Decline in Rate of FDA Hydrolysis and Total Particulate Organic Matter

Fig. 3 DR-FTIR Spectra of Separated Cow Manure through the Composting Process

Fig. 4. DR-FTIR Spectra of Particulate Organic Matter from Suppressive and Conductive Compost-Amended Sand Mixes

Fig. 5. Changes in Chemistry Accompanying the Decline in Suppressiveness to Pythium Root Rot: Subtraction Spectrum

Fig. 6. Changes in Chemistry Accompanying the Decline in Suppressiveness to Root Rot: PCA Coefficient Plots

Fig. 7. Scatter Plot of PC Scores Representing Spectra of Particulate Organic Matter from Suppressive And Conductive Compost-Amended Sands

Fig. 8. Timeline

IX. APPENDIX II - RESEARCH PERFORMED IN ISRAEL

Principal Investigators

Prof. Yitzhak Hadar

Prof. Yona Chen

Investigators

Benny Chefetz

Rony Coehn

Nir Bar-Lev

Table of Contents

1. Composting and recycling of organic wastes
2. Formation and properties of humic substances originating from composts
3. Chemical and biological characterization of organic matter during composting of municipal solid waste
4. Suppression of soil-borne pathogens by composted municipal solid waste

Composting and Recycling of Organic Wastes

Introduction

Both quality and quantity of organic matter (OM) in soils affect plant growth and health. Crop residues and root exudates affect the activity of beneficial microorganisms, as well as pathogens (Chang et al., 1986). The effects of OM decomposition level in the soil on interactions of beneficial microorganisms and pathogens, and on plant growth have been largely overlooked (Hoitink and Fahy, 1986; Chen et al., 1988; Grebus et al., 1994). Both the mineral fraction and the yearly introduced organic residues, as well as their resulting transformations in soils, play a role in this complex. In addition, humic substances (HS), the most stable organic components in soil, may have both direct and indirect effects on plant growth (Stevenson, 1985; Chen and Aviad, 1989; Chen et al., 1994).

The composting of organic wastes has been defined as a controlled microbial aerobic decomposition process. Recently composting has gained interest in western countries as a result of the accumulation of large amounts of agricultural, industrial and municipal wastes (de Bertoldi et al., 1987; de Bertoldi, 1993; Steuteville, 1994a,b; Golsdtein and Steuteville, 1995). These wastes create a severe disposal problem involving a number of environmental aspects related to air, soil, ground water and surface water pollution. During the course of the composting process, transformations of various OM fractions can be followed separately without analytical interference from OM already present in the soil. The composting process therefore provides a unique opportunity to evaluate effects of specific OM transformations on plant growth and health and may serve as a model for field soil systems. The main products of aerobic composting are CO₂, water, minerals and stabilized OM often called humus. Well-composted OM is traditionally used as a source of nutritional elements and/or soil conditioner directly in the field. In the last decade, the demand for peat as a potting medium in horticulture has increased while its availability has decreased (Chen et al., 1992). A number of organic wastes such as bark, leaf mold, municipal solid waste (MSW), sewage sludge, sawdust, and treated animal excreta have been introduced as peat substitutes in growth media after proper composting (Cull, 1981; Bik, 1983; Lohr et al., 1984; Verdonck, 1984; Inbar et al., 1985; Raviv et al., 1986; Chen et al., 1988; Chen et al., 1992). Some of the well-matured composts have been found to be suppressive of soilborne plant pathogens when utilized as container media (Hoitink and

Fahy, 1986; Hoitink and Kuter, 1986; Chen et al., 1988; Mandelbaum et al., 1988; Hoitink et al., 1993).

Principles of composting

Composting is a form of waste stabilization that requires special conditions, particularly of moisture and aeration, to yield temperatures conducive to thermophilic microorganisms (Hoitink and Fahy, 1986). Zucconi and de Bertoldi (1987) proposed the following definitions: Composting is a controlled biooxidative process that: (i) involves a heterogeneous organic substrate in its solid state; (ii) evolves by passing through a thermophilic phase and a temporary release of phytotoxins; and (iii) leads to production of CO₂, water, minerals and stabilized OM (compost). Compost is the stabilized and sanitized product of composting which is beneficial to plant growth. The OM has undergone an initial, rapid stage of decomposition and is in the process of humification.

Controlled environmental conditions distinguish composting from natural rotting or putrefaction, which often occurs in open dumps, manure heaps, or field soil. A simplified flowchart of the overall composting process is shown in Fig. 1. This flowchart illustrates the interactions between the organic waste, microorganisms, moisture and oxygen. The organic wastes, comprised of several components, normally have an indigenous population of microorganisms derived from the atmosphere, water or soil. Once the moisture content of the waste is brought to an appropriate level and the mass is aerated, microbial metabolism speeds up. Energy is obtained by biological oxidation of part of the carbon; some of this energy is used in metabolism, the rest is given off as heat. The end product (compost) is made up of the more resistant residues of the OM, breakdown products, dead and living microorganisms, together with products from further chemical reactions between these materials (Gray and Biddlestone, 1981).

The temperature trend in most composting processes is presented in Fig. 2. Basically, the process can be divided into four stages: (i) mesophilic stage - an initial phase of 1-2 days, during which the mesophilic strains of microorganisms start to decompose the readily degradable compounds, heat is given off and the temperature rises. The pH falls slightly as organic acids are produced; (ii) thermophilic stage - the

second stage, lasting 2-4 weeks, when above 40°C the thermophilic strains take over. If the temperature rises above 60°C, the fungi become deactivated, and the reaction is continued by thermophilic bacteria. In this phase the more readily degradable substances such as sugars, fats, starch and proteins are rapidly consumed and most of the human and plant pathogens are destroyed; the pH becomes alkaline as ammonia is liberated from the proteins. The reaction rate decreases as the more resistant materials are attacked; the compost enters (iii) the cooling-down stage - as the temperature falls, the thermophilic fungi reinvade the composted material from the cooler extremities and start to attack the cellulose, hemicellulose and other polymers. These first three stages of composting last from a few weeks to 2 months, depending on the composted material; the last stage (iv) stability and maturity - requires several months, where little heat is generated and the final pH is normally slightly alkaline. During this phase, mesophilic microorganisms as well as macrofauna colonize the compost, Intense competition for food take place between the microorganisms involving: antagonism and the formation of antibiotics. Humification occurs in the residual OM to produce the stable composted product (Gray and Biddlestone, 1981; Diaz et al., 1982; Hoitink and Fahy, 1986; Zucconi and de Bertoldi, 1987; He et al., 1992; Jimenez and Garcia, 1992; Chen and Inbar, 1993; Ciavatta et al., 1993; Inbar et al., 1993; Adani et al., 1995).

The composting of organic wastes is a dynamic and extremely complicated ecological process in which temperature, pH and nutrient availability are constantly changing. As a consequence, the numbers and species of organisms responsible also change markedly. Thermophilic bacteria, particularly *Bacillus* spp. and *B. cereus* (Finstein and Morris, 1975; Kwok et al., 1987), appear to dominate the early phase of high activity, but thermophilic actinomycetes predominate thereafter (Nakasaki et al., 1985). The rate of decomposition depends on the rates of microbial activity, the latter being greater when composts are maintained at low (38-55°C) rather than high temperatures (60-70°C) (Kuter et al., 1985). In addition to temperature, optimum starting material values for composting are: C/N ratio 30-35; C/P ratio 75-150; moisture content 50-60%; particle size 15-35 mm; pile size 1.5 m high and 2.5 m wide

Compost maturity

Compost maturity is defined as the degree of OM decomposition during composting. Any definition of maturity must be based on the potential use of the compost. For example, compost used in potting mixes (container media) must be mature enough so that no plant growth inhibition occurs within days after formulating of the mix. On the other hand, composts utilized in field agriculture may be applied weeks or months before planting, giving them time to "stabilize" in the soil. Thus, the following definitions are proposed for mature composts: (i) Greenhouse utilization: OM composted to the degree of decomposition which has no adverse effects on container grown plants. (ii) Field application: OM composted to the degree of decomposition that has no adverse effects on the growth of various crops when applied at annual rates of up to 50 tons/ha, at least 6 weeks before sowing or planting (during the warm season, in which decomposition can take place).

Chemical parameters for compost maturity

The chemical nature of the decomposition process and the evaluation of compost maturity are a difficult problem, since several parameters are required to provide a definite evaluation for the readiness of the material for use in horticulture (Morel et al., 1985; Zucconi and de Bertoldi, 1987). Some of the main changes taking place in the solid phase of the OM during decomposition in soils and/or in partially controlled environments such as a compost pile are: an increase in ash content, a decrease in cellulose and hemicellulose, and generally, as a result of concentration, a relative increase in lignin (Harada and Inoko, 1980b; Inoko, 1982).

Several procedures for testing stability or maturity of composts have been proposed. Some examples are: starch content, cellulose content, C/N ratio, C/N ratio in water extracts of composts, concentration of HS, presence of indicator microorganisms, respiration rate, and others (Harada and Inoko, 1980a,b; Chanyasak and Kubota, 1981; Harada et al., 1981; Chanyasak et al., 1982; Lax et al., 1987; Inbar et al., 1989; Inbar et al., 1990b; Inbar and Chen, 1993).

The C/N ratio represents one of the major guidelines for the composting process. However often the C/N ratio of bulk composts does not provide useful information; in wood wastes for example, much of the carbon is not readily biodegradable. For such wastes, availability must be considered. High "available" C/N ratios of immature or

inadequately composted materials cause immobilization of nitrogen and nitrogen deficiency in plants. High microbial activity existing in immature compost requires O_2 and consequently may cause oxygen deficiency to roots. During the decomposition of fresh or partially decomposed OM, adverse effects on plant growth result from toxic fermentation byproducts produced in the early stages of composting. This phenomenon is well-known in relation to the incorporation of crop residues into soils. Lynch et al. (1980) observed a reduction in root growth of young barely plants cultured in different soils amended with straw. They attributed the reduction to an accumulation of acetic acid produced by decomposition of the straw. Toussan et al. (1968) identified benzoic acid and phenylacetic acid as phytotoxins present in the decomposition products of barley sampled in the field. Acetic acid has also been identified as a phytotoxin in immature composts by De Vleeschauwer et al. (1981). Anid (1986) identified high levels of acetic, propionic and butyric acids in immature compost, whereas those volatile acids disappeared when stabilization was achieved. Wong and Chu (1985) noted severe suppression of germination when seeds of *Brassica chinesis* were treated with extracts of fresh refuse compost. Less adverse effects were found with 6 week-old compost. Differences from the control were not found with the 12 month-old compost. These investigators speculated that many factors could be influencing germination and root elongation including pH, salt and ammonia contents, ethylene oxide concentration, and heavy metals. Zucconi et al. (1981a,b) tested the response of a wide range of plant species in various bioassays to fresh, immature and mature composts. In general, all plants were inhibited by immature compost. Growth usually improved upon contact with mature compost. These authors concluded that phytotoxicity during composting appears to be strictly associated with the initial stage of decomposition and that it is a transient condition, possibly connected to the presence of readily metabolizable material. Low molecular weight organic acids (acetic, propionic, and butyric) were the most toxic compounds found in the early stages of the composting process (De Vleeschauwer et al., 1981; Zucconi et al., 1981a,b; Chanyasak et al., 1983; Hirai et al., 1986). Finally, allelopathic toxins in wastes can cause problems, but they are generally destroyed within a few weeks of composting. Hoitink and Kuter (1986) concluded that the negative responses of plants grown in bark composts were due to high cellulose

content resulting in nitrogen immobilization, allelopathic chemicals, toxic levels of metabolites of anaerobic metabolism, high salt content and, possibly, other factors. They concluded that appropriate curing is essential to stabilize compost and eliminate or reduce negative plant responses (Hoitink and Fahy, 1986). Inbar et al. (1988) measured O₂ consumption by fresh and composted separated manure (CSM) and grape marc (CGM) at various moisture levels, with and without the addition of nitrogen, as a function of composting time in a simulation system. For both materials, cumulative O₂ consumption increased rapidly at first, then leveled off. Maximum O₂ uptake was obtained at moisture levels of 60 to 70% (on a net weight basis). The addition of nitrogen increased the O₂ uptake rate in some instances, more markedly in the CGM which is low in nitrogen content. The O₂ uptake rate vs. time approached leveling off after a few days of composting in the pilot-scale system, exhibiting a slow approach to a steady state in microbial activity. As opposed to the fresh OM for both materials, the composted products had low O₂ uptake rates, indicating that O₂ consumption provides a good measure of the rate of decomposition and the degree of stability. A similar relationship between O₂ uptake and MSW composting age was observed by Iannotti et al. (1994). This measure, as well as CO₂ release, has been used by microbiologists for many years for this purpose. However, leveling off or a lower rate of O₂ uptake is often achieved before compost can be considered mature for greenhouse utilization (Inbar et al., 1988). This may not necessarily be the case for compost applied to soil well ahead of planting, where only some stability is required.

To avoid problems caused by carbon availability, the ratio of water-extractable organic C/organic N was proposed as an indicator of compost maturity by researchers in Japan (Chanyasak and Kobuta, 1981; Chanyasak et al., 1982, 1983). These researchers found values of 5 to 6 for the soluble organic C/organic N ratio of mature compost of MSW. A highly significant negative correlation was established between the cation exchange capacity (CEC) and the soluble C/N ratio. However, this was not confirmed for CSM by Inbar et al. (1989), nor for CGM by Inbar et al. (1990a). It also seems, as proposed by Hoitink and Kuter (1986), that this procedure is not effective for low C/N ratio wastes such as sewage sludge which, after the mixing of raw sludge and bulking agents, already has low ratios in the extract.

Biological parameters of compost maturity

The ultimate, and probably most attractive evaluation of compost maturity is based on bioassays. Most are based on plant growth over a few days or weeks, and this limits their usefulness. Data given in Table 1 (Hadar et al., 1985) show that extended composting was needed to obtain maximum growth of plants in separated manure. Growth of unfertilized plants was retarded but fertilization alone could not sufficiently enhance growth unless the compost was mature. This suggests that nitrogen deficiency was not the only growth-limiting factor in this immature compost. An experiment conducted using water extracts of CSM (Inbar et al., 1989) showed that tomato seedlings were inhibited in water extracts obtained from composts less than 40 days old. An increase in inhibition resulting from phytotoxic substances was observed in compost extracts prepared during the thermophilic stages of composting (20-40 d). From day 40 onward, growth in compost extract was significantly better than that in a standard nutrient solution. It was concluded that phytotoxic compounds and competition for O₂ in the immature compost were responsible for the inhibition in plant growth. Grebus et al. (1994) reported that 60-70 days of composting were required to destroy the inhibitory growth effect of a yard-trimmings mixture on radish. Plant growth bioassays revealed that ryegrass, cucumber and radish differ in their responses to MSW compost age (Iannotti et al., 1994). These bioassay revealed that ryegrass and radish treated with slow-release fertilizer are sensitive to compost maturity, but cucumber is not. Different results were observed by Chefetz et al. (1995) who reported that a bioassay based on cucumber growth is a sensitive tool for determining MSW compost maturity.

Humic substances in composts

HS have attracted the interest of soil scientists due to their importance to soil ecology, fertility and structure and their effects on plant growth (Vaughan and Malcolm, 1985; Chen and Aviad, 1989; Chen et al., 1994). As more agricultural lands throughout the world are treated with composted organic materials such as composts from sewage sludge, municipal refuse, crop residues, and animal excreta, elaboration of the nature of the organic components (especially the HS) present in such composts, as well as of their formation during the composting process, has become essential. HS

are known to be involved in a number of interrelated, chemical processes in soil and to influence plant growth directly as well as, indirectly. Indirectly, humic acid (HA) may improve soil fertility by increasing the availability of plant nutrients or by sequestering heavy metals such as iron, zinc, and manganese in the soil environment, thereby solubilizing these elements for the benefit of plants (Chen and Aviad, 1989). Some reports also suggest that plant roots can take up HS, thereby allowing them to have direct affect on plant growth and metabolism (Vaughan and Malcolm, 1985).

A variety of techniques have been employed to extract HS from soils. The methods vary among researchers, depending upon the nature of the material to be examined, the discipline of the investigators and other arbitrary factors. The common extraction procedures have been reviewed by Hayes et al. (1975), Hayes (1985), Stevenson (1982), Watanabe et al. (1994) and Adani et al. (1995). They stated that aqueous solutions of NaOH are capable of extracting appreciable fractions of OM, including the more complex, high-molecular-weight and less soluble components of humus. Aqueous solutions of NaOH are known to induce slight oxidation of the extracted OM. The alkali solution also dissolves protoplasmic and structural components from fresh organic tissues and these become mixed with the humified OM (Stevenson, 1982). When fresh plant residues and young peat material were extracted with alkali, substances such as hemicelluloses, proteins, alkali-soluble lignin, and tannins entered into the humic fraction. However, as the decomposition proceeded, the fraction of these substances decreased while that of the HS increased (Inbar et al., 1993; Adani et al., 1995). In the case of strongly humified materials, the danger of extracting the admixtures with the HA is practically negligible. To minimize the chemical changes caused by autooxidation during the alkali extraction, all steps should be conducted in the presence of an inert gas (e.g. N₂) (Stevenson, 1982).

Among the chemical criteria required for the characterization of an stabilized organic waste product (e.g. compost), humus content and quality are highly ranked (Sugahara and Inoko, 1981; Roletto et al; 1982; Roletto and Ottino, 1984; Roletto et al., 1985; Sequi et al., 1986; Saviozzi et al., 1988; Jimenez and Garcia, 1992; Chen and Inbar, 1993). The main trends of several humification parameters (expressed as percentage of variation) over a period of 140 days during the stabilization of wastewater sludge and straw mixtures are shown in Fig. 3 (Saviozzi et al., 1988). The

following humification indexes were suggested: (i) C/N ratio, (ii) CHA - humic acid carbon (percentage of organic C), (iii) CFA - fulvic acid carbon (percentage of organic C), (iv) CHA/CFA ratio, (v) CHA+CFA (percentage of organic C), (vi) HI (index of humification) - the ratio between the organic C of the non-humified fraction (NHF) and that of the HA + FA fraction (after Sequi et al., 1986), (vii) organic C (% DM), (viii) stability index (SI) - the ratio between HA after apolar, polar and acid hydrolysis treatments and NaOH-Na₄P₂O₇ HA extraction (after Adani et al., 1995). During the first 30 to 60 days changes in HI occurred rapidly; thereafter, the rate of change became progressively slower. All the parameters were reported to be linearly and significantly correlated to the C/N ratio. In their work, Saviozzi et al. (1988) suggested minimum values for some humification parameters indicating maturity of farmyard manure (FYM). FYM containing less than 37.1% organic C, and exhibiting an HI value lower than 0.4, a CHA + CFA higher than 16, and a CHA higher than 9.7, could be considered mature. Time-course studies of the composting of various wastes showed an increase in the relative amount of HA (from 18% to 45% of OM in CSM compost and from 5% to 20% in MSW compost), whereas the formation rate of fulvic acid (FA) was inconsistent. Humification ratio (HR=HA/FA) values of 0.9 to 3.4 and 3.0 to 9.2 were typical for immature and mature composts, respectively. Although the values of the HR for various composts differ, they generally increase by more than 80% during composting, indicating a high degree of OM polymerization, since more polymerized components are found in mature compost. HI decrease during composting represents the formation of HS. Elemental and functional-group analysis of the HA originating from various composts indicated only minor differences between composting stages. Moreover, these values fell into a wide range, similar to that of soil HA. In general, even though HA amounts increased during composting, the elemental and functional groups were similar to "young forms" of HA reported in the literature. The E₄/E₆ ratio was higher in compost HA, indicating a lower molecular weight than that of "model" HA (Chen et al., 1977). Elemental and functional-group analysis of compost FA have been reported in only a few works (Roletto and Luda, 1987; Pandeya, 1992). The compost FA had lower carbon and nitrogen contents but higher amounts of oxygen and carboxyl groups than HA, and the overall elemental and

functional groups extracted from compost were similar to "model" FA (Schnitzer, 1978; Roletto and Luda, 1987; Pandeya, 1992).

CPMAS ^{13}C -NMR, FTIR and DRIFT spectroscopies

During the process of compost production, organic materials that are chemically complex and difficult to fractionate are formed. Extraction procedures that are commonly applied to soils and composts remove only a fractions of the C in the system. To profile the organic material of composts during the course of their formation, a method that can be applied to the intact sample is preferable. The only physical method holding some promise for the reliable characterization of such organic materials appears to be nuclear magnetic resonance (NMR) (Wershaw, 1985; Wilson, 1987). Recently, the application of NMR spectroscopy in studies on HA has gained considerable attention. Both ^1H - and ^{13}C -NMR, in the solid and liquid state, have been used. Wershaw (1985) indicated, that although NMR analysis has been applied for a much shorter period of time than most other techniques, it has provided more definitive structural information than other methods. Recent studies have shown that the technique of solid-state CPMAS (cross polarization magic-angle spinning) ^{13}C -NMR can provide carbon "fingerprints" of diverse solid samples such as peats (Wilson et al., 1983; Preston et al., 1987), whole soils (Preston and Ripmeester, 1983; Wilson et al., 1983; Preston et al., 1994), composts (Piotrowski et al., 1984; Gerasimowicz and Byler, 1985; Preston et al., 1986; Inbar et al., 1990a, Chefetz et al., 1995), forages (Eflosen et al., 1984), wood (Kolodzieski et al., 1982; Haw et al., 1984), and fossil fuels (Packer et al., 1983), HS isolated from soils, peat, river water, river sediments, lake water, marine waters, and marine sediments as reviewed by Malcolm (1989). An in-depth review of the state of the art of NMR spectroscopy theory, techniques, and applications to a wide range of organic and inorganic geochemistry and soil chemistry studies are discussed by Wilson (1987). Plant and microbial remains are the substrate for OM transformations in the geosphere. The myriad of chemicals which compose plants and microbes can be grouped into six major types: lignins, polysaccharides, proteins, lipids, resins and pigments (Wilson, 1987). The contribution of CPMAS ^{13}C -NMR assigned chemical shifts for each of these groups is described by Wilson (1987). Wershaw et al. (1986, 1988) compared spectra of HA fractions from several different

environments with published spectra of plant components; they showed that these components, when incorporated into HA retain many of the chemical structural features of the original plant material. They attributed some of the observed bands to lignin-like or melanin-like structures. Sharp and well-defined carbohydrate bands were observed in the spectra of some of the fractions. Lignins are composed of three basic building blocks, namely syringyl, guaiacyl, and p-hydroxyphenolic units. Resonances from these structures can be observed in all lignins. Typical ^{13}C -NMR assignments to these structures are presented in Fig. 4. A number of structural differences between lignins from various sources have been reported, but in general, many investigators agree that the basic units are those shown in Fig. 4. The intensity in the ^{13}C -NMR spectral region from 160 ppm to 110 ppm is due to the aromatic ring carbons of lignin. In addition, the signal around 56 ppm is usually attributed to methoxy carbons, principally of lignin origin (Himmelsbach and Barton, 1980; Haw et al., 1984; Wilson, 1987).

The second most important organic precursors to HS are polysaccharides, of which cellulose is the most important. Cellulose consists of long chains of glucose units, each of which is combined by a β -glucoside link to the C4 hydroxyl or another glucose unit (Fig. 5). Enzymatic hydrolysis of cellulose can vary in nature, as can the number of glucose units that are present. It is not uncommon to find 2-4000 glucose units joined together (Wilson, 1987). In the solid state, cellulose exists in at least four forms- the most common of which are cellulose I and II that may have slightly different chemical shifts. Moreover, these cellulose units may exist in various degrees of crystallinity (Haw et al., 1984; Wilson, 1987). The features of cellulose I (the native or natural form of cellulose) are: a peak at 105 ppm attributed to the C_1 carbon of the anhydroglucose repeating unit (Fig. 5); signals at 89 and 84 ppm, resulting from the C_4 carbon of cellulose; the highest absorption intensities for C_2 , C_3 and C_5 carbons in the 70 to 80 ppm region; peaks at 63 to 66 ppm are assigned to the C_6 carbon of cellulose (Haw et al., 1984; Wilson, 1987). Hemicellulose exhibits spectral features which are similar to cellulose. In addition, it exhibits peak at around 103 ppm due to carbons that are structurally analogous to the C_1 carbon of cellulose, and signals at 22 and 174 ppm which result from acetate groups present in hemicellulose. Some of the absorption

intensity of the 174 ppm peak could be due to other carbonyl species present in hemicellulose (e.g. CO₂H groups of uronic acid) (Kolodziejewski et al., 1982).

The ¹³C-NMR spectra can be divided into five regions (Hatcher et al., 1983). Region I (0-50 ppm) is assigned to paraffinic carbon or carbons that are bonded only to other carbons. Region II (50-112 ppm) is assigned to C-O, C-N bonds as in carbohydrates, alcohols, esters and amines. Region III (112-163 ppm) is assigned to aromatic and phenolic carbons. Region IV (163-190 ppm) represents carboxyl, ester and amide carbons groups. Region V (190-215 ppm) represent the carbonyl carbons. The 0 to 112 ppm region was calculated as aliphatic carbon, the 60 to 112 ppm region as polysaccharide carbon and the 112 to 163 ppm region as aromatic carbon. Total aromaticity was calculated by expressing aromatic C (112-163 ppm) as percentage of aliphatic C (0-112 ppm) plus aromatic C (Hatcher et al., 1981; Kögel-Knabner et al., 1991).

To date, many studies have been conducted on HA compost using CPMAS ¹³C-NMR spectroscopy, but only a few used this method to study bulk compost (Inbar et al., 1989; Chefetz et al., 1995). Inbar et al., (1989) reported only minor differences between the spectra of HA from fresh and composted manure (12% increase in carboxyl content; 8% increase in aromatic carbon and 8% decrease in alkyl carbon). During composting of MSW, HA aromatic carbon increased by 39% while the aliphatic carbon content decreased by 19% (Garcia et al., 1992). During composting of MSW, the following distinct changes in the CPMAS ¹³C-NMR spectra of the bulk compost resulting from the composting process were noted: (i) Total aliphatic carbon decreased by 14.4% from 74.8% in the raw material to 64.0% in the mature compost; the polysaccharides decreased by 17.0%, from 42.3% to 35.1% in the mature compost; and the alkyl groups (0-60 ppm) decreased by 11.0% during the composting process; (ii) the aromatic components, including aromatic and phenolic carbons, increased by 54%, from 15.9% in the raw material up to 24.5% of the total carbon in the mature compost. Total aromaticity increased by 58.2%, from 17.5% in the raw material to 27.7% in the mature compost; (iii) carboxyl carbon increased by only 9.5%; (iv) the C=O carbonyl carbon increased by 73.7%, from 1.9% in the raw material to 3.3% in the mature compost (Chefetz et al., 1995).

Infrared (IR) spectroscopy and a wide variety of chemical methods are being used to characterize HS. Fourier transform infrared (FTIR) spectroscopy serves as a qualitative tool to nondestructively characterize the principal classes of chemical groups that compose the HA. Negative feedback as to the usefulness of IR spectral analysis is probably due, in part, to the complexity of the IR spectra. However, IR analysis has contributed substantially to the knowledge of HA chemistry (Schnitzer, 1978; Stevenson, 1982; MacCarthy and Rice, 1985), and complements the information obtained from solid-state CPMAS ^{13}C -NMR spectroscopy (Gerasimowicz and Byler, 1985; Lobartini and Tan, 1988). The structural complexity of humic materials and the abundance of strongly hydrogen-bonded groups have, until recently, severely limited the usefulness of infrared spectroscopy for structural and compositional studies. With dispersive instruments only broad, ill-defined bands have been observed (Gerasimowicz et al., 1986). FTIR spectroscopy, which provides a much higher signal-to-noise ratio and improved accuracy in the frequency of the instrument, increased the information attainable from the spectra (Griffiths, 1983). Gerasimowicz and Byler (1985) and Gerasimowicz et al. (1986) obtained FTIR spectra on solid HA extracted from materials sampled at four selected stages of a municipal wastewater treatment process. From the FTIR spectra, several important differences between the decomposition stages were observed and discussed. Sugahara and Inoko (1981) investigated HA extracted from city refuse at different stages of decomposition by IR spectroscopy. They observed decreases in aliphatic C-H stretching, increases in COO- and decreases in C-O stretching as composting proceeded. On the whole, the shape of the IR spectra of city refuse HA became featureless during composting. Similar changes were observed by Zech et al. (1987) for IR spectra of fresh and decomposed spruce and pine litter. In many instances, CPMAS ^{13}C -NMR spectroscopy data lend themselves somewhat better to quantitative analysis than do the FTIR results. In particular, the relative aromatic nature of the HA was more evident from the NMR spectra and could not be seen as readily from FTIR data (Gerasimowicz and Byler, 1985). The development of an attachment mounted onto FTIR spectrophotometers has allowed the determination of diffuse reflectance infrared Fourier transform (DRIFT) spectra. Sample preparation for DRIFT spectroscopy is much simpler than for FTIR spectroscopy, interferences due to water adsorption are reduced, and resolution is

improved. The spectra obtained using DRIFT spectroscopy had a higher degree of resolution as compared with FTIR spectroscopy. Moreover, the bands indicative of aliphatic C-H, carboxyl and carboxylate functional groups, aromatic C=C, and the C-O stretch of polysaccharides were prominent and very well resolved. The DRIFT spectra obtained can also be used to fingerprint OM acquired from various sources. Relative concentrations of functional groups were found to be fairly constant regardless of sample concentration. The DRIFT spectra peak intensity ratio can be used to determine the humification process. The application of DRIFT spectroscopy to OM research will prove especially useful for characterizing heterogeneous bulk samples such as peat and composts. A combination of quantitative elemental analysis supplemented by modern, nondestructive NMR, FTIR and DRIFT characterizations could have significant biochemical implications in the study of the nature and properties of HS and other complex heterogeneous matrices (Niemeyer et al., 1992).

Raw material used for composting

Municipal solid waste:

Problems caused by MSW in modern society have become more severe over the last decade due to increasing amounts of waste and decreasing availability of landfill space (Alter, 1991; Finstein, 1992). The amount of MSW generated in the US rose to 323 million tons in 1994, an increase of 5%, whereas during the period between 1988 and 1994 the number of landfills decreased by more than half (from 8000 to 3558 in 1994). The remaining capacity of the landfills has also decreased and in some states such as Ohio, Missouri, and Vermont the landfills can last another 10 years or less. Together with the decreasing landfill capacity, the tipping fee has increased and can reach up to 75\$ per ton (Steuteville, 1994a,b, 1995). The European community produces more than 2500 tons of garbage a year of which the MSW fraction can reach up to 10%. Thus recycling has become an attractive solution for MSW treatment. The organic fraction in the MSW varies between 30 and 50%. Thus recycling this fraction can reduce the cost of transportation by 50%. For the organic fraction, composting seems to be a desirable option, having the capacity of reducing the volume and weight by approximately 50% and resulting a product which can be useful for agriculture (He et al., 1992). An alternative solution for organic waste management is the waste to

energy program, but its high tipping fee and its minor daily capacity make composting the ideal solution for the organic fraction. The main problems facing MSW compost production and utilization stem from the heterogeneous nature of the raw material. Its composition can change with climate, standard of living, type of garbage collection system and season.

Yard waste:

Yard waste can be defined as the amount of leaf fall, grass clippings and wood waste diverted from the MSW stream. In 1988, the EPA estimated that 18% of MSW is yard waste. Yard waste is not homogenized waste and its characteristics vary with weather, area and collection policy. The characteristics of the material affect the collection treatment, the processing equipment and the properties of the end product. Yard waste humidity varies from 30 to 75%, with that of leaves varying between 19 and 28%, of grass from 54 to 64% and of the wood fraction from 17 to 18%. The C/N ratio of yard waste is usually high (90-125) due to the large levels of lignin in woody waste. To reduce the initial C/N ratio to a level suitable for composting (30-35), yard waste is usually mixed with animal manure, sewage sludge, poultry manure or any other waste that can provide additional nitrogen. Recovery of yard waste by composting is an increasingly popular method of recycling solid organic waste (Steuteville, 1995). The common systems used for composting yard waste are static piles and turned windrows. Yard waste compost is useful as a substrate in container media and unlike composts produced from other wastes (MSW and sewage sludge) this compost seems to be "cleaner".

Sewage sludge:

The level of sewage sludge, the by product of normal wastewater treatment operations, is increasing worldwide. Utilization of sewage sludge in agriculture as an organic fertilizer is regarded as a beneficial and economic process. Chemical properties of sewage sludge indicate high rates of nitrogen and phosphorus. Sewage sludge can be applied to soil as a liquid or solid amendment. An increasing concern over human pathogens in the sludge has enhanced the interest in using composting as a process that can destroy plant and human pathogens and produce useful agricultural products. The main problem facing composting of sewage sludge is its moisture content. Thus before composting, sewage sludge humidity must be brought to less than 60% by drying or by

adding a dewatering agent (lime). Usually sewage sludge cannot be used as only raw material for composting, so yard waste or straw litter are used as bulking agents.

Separated cattle manure:

The total amount of organic waste products in the US from livestock and poultry production has been estimated at over 1.5 billion tons per annum (on a dry matter basis). Of this amount, about 1 billion tons is cattle manure (Stevenson, 1985). Because of technological and economic considerations, the monetary value of manures and other wastes has significantly decreased in recent years, with the result that these materials are now regarded as a liability rather than an asset to the total farm operation. Farming has become highly specialized and livestock and poultry production is being concentrated in to large-scale confinement-type enterprises. These changes imply that large volumes of wastes have to be disposed of in a relatively small area (Stevenson, 1985). Stevenson (1985) summarized the limitations regarding the use of farmland for the disposal of animal manure as follows: (i) animal manure as a source of plant nutrients is a bulky, low-grade fertilizer of variable composition; (ii) concentration of nutrients, soluble salts, trace elements, and water vary tremendously and are seldom known. Optimum rates of application are thus difficult to predict; (iii) in many soils, the leaching of nutrients present in high concentrations in manures (especially NO_3^-) in to the ground water may limit application rates; (iv) on-farm management problems are created by the physical properties of manure. Application techniques are inefficient and time consuming; (v) production of animal manures is continuous, while the need for fertilizers is seasonal; (vi) odor and associated nuisances; and (vii) the presence of some human and plant pathogens and weeds restrict the use of manures by farmers. Development of new management systems which will increase the quality of the products, decrease salt concentration, and improve the physical structure and ease of handling are needed. One of the proposed systems now used on farms is based on separating of the manure into solids and liquid (Huijsmans and Lindley, 1984; Bohnhoff and Converse, 1987). This material can be successfully used, after proper composting, as a peat substitute or complementary product for peat mixtures in container media, while the effluent is used as organic liquid fertilizer (Chen et al., 1984a,b; Levanon et al., 1984; Raviv et al., 1984; Hadar et al., 1985; Chen and Hadar, 1987).

Grape marc:

The primary by-product of grape juice processing and wine production is grape marc (GM; pomace). It consists primarily of processed skins, seeds and stems. GM represents as much as 20% of the wet weight of the original fruit and its moisture content is about 65% (Hang, 1988). Annual world production of dried GM is estimated at 7,500,000 metric tons (Aguilera, 1987). Direct incorporation of GM, without any pretreatment, into vineyard soils is a universal practice. However, this method has become a serious problems since degradation materials that are highly incompatible with plant roots are released. Composting GM has the potential to solve the environmental problems related to this byproduct and to create a useful product for greenhouse agriculture.

Conclusions

Composting seems to be a desirable solution for organic waste management. Readily available wastes such as cattle manure, MSW, sewage sludge, winery waste and yard waste have been found to be suitable materials for composting. The composting process has the capacity to reduce the volume and weight of these wastes by approximately 50% and to produce a product which can be useful in agriculture. Composts produced from various wastes have been found to be suitable materials for container media having the ability to reduce peat and pesticide use thereby solving environmental problems. The positive effects of the composts on plant growth can be explained by their physical, chemical and biological properties. Their physical properties fit well with optimal requirements. In greenhouse setups, where fertigation is continuously applied, the nutrient content is not as important. However, the slow release of nutrients between irrigations, the high contents of HS and the bioactivity of the medium improves plant growth. Mature composts have also been found to be suppressive to several soilborne plant pathogens such as *P. aphanidermatum*, *R. solani* and *S. rolfsi*. As a result, application rates of hazardous pesticides can be reduced or eliminated. Amendment of composts to soils has been shown an increase crop yield, while amendment of compost with mineral fertilizer increases crop yield as compared to the amendment of either compost or mineral fertilizer alone. The positive effects of mature compost on plant growth are connected to its high HS content. The HS formed

during composting seem to have the same characteristics as HS that are formed in soils over long periods. Spectroscopic methods (CPMAS ^{13}C -NMR, DRIFT and FTIR) are useful tools to determine the transformations of the organic fraction during composting, and can help to define compost maturity. During composting, many parameters must be monitored (temperature, humidity, pH, C/N ratio, aeration), although a number of the parameters stabilize at the maturation stage providing no tools for the actual maturity stage. Only a few parameters can be used to define compost maturity. Many authors have reported that HI, C/N ratio, respiration and microbial activity can be used as maturity indexes. We hypothesize that plant growth may serve as an integrative signal for compost maturity.

References

- Adani, F., P.L. Genevini, and F. Tambone. 1995. A new index of OM stability. *Compost Science & Utilization* 3:25-37.
- Aguilera, J.F. 1987. Improvement of olive cake and grape by-products for animal nutrition. p. 45-54. In J.M. van der Meer, B.A. Rijkens and M.P. Ferranti (Eds.) *Degradation of Lignocellulosics in Ruminants and in Industrial Processes*. Elsevier Applied Science, London.
- Alter, H. 1991. The future course of solid waste management in the U.S. *Waste Management & Research* 9:3-9.
- Anid, P.J. 1986. Evaluating maturity and metal transfer of MSW compost. *BioCycle* 46-47.
- Bik, A.R. 1983. Substrates in floriculture. *Proc. XXI Intern. Hortic. Cong. Hamburg* 11:811-822.
- Bohnhoff, D.R., and J.C. Converse. 1987. Engineering properties of separated manure solids. *Biological Wastes* 19:91-106.
- Chang, Y.C., R. Maker, O. Kleifield, and I. Chet. 1986. Increasing growth of plants in the presence of biological control agent Trichoderma harzianum. *Plant Disease* 70:145-148.
- Chanyasak, V., and H. Kubota. 1981. Carbon/organic nitrogen ratio in water extract as a measure of composting degradation. *J. Ferment. Technol.* 59:215-219.

- Chanyasak, V., M. Hirai, and H. Kubota. 1982. Changes of chemical components and nitrogen transformation in water extracts during composting of garbage. *J. Ferment. Technol.* 60(5):439-446.
- Chanyasak, V., A. Katayama, M.F. Hirai, S. Mori, and H. Kubota. 1983. Effects of compost maturity on growth of komatsuna (*Brassica rapa* var. *peridis*) in Neubauer's pot. II. Growth inhibitory factors and assessment of degree of maturity by org.-C/org.-N ratio of water extract. *Soil Sci. Plant Nutr.* 29:251-259.
- Chefetz, B., P.G. Hatcher, Y. Hadar, and Y. Chen. 1995. Chemical and biological characterization of OM during composting of municipal solid waste. *J. Environ. Qual.* [In Press].
- Chen, Y., N. Senesi, and M. Schnitzer. 1977. Information provided on humic substances by E4/E6 ratios. *Soil Sci. Soc. Am. J.* 41:352-358.
- Chen, Y., Y. Inbar, M. Raviv, and A. Dovrat. 1984a. The use of slurry produced by methanogenic fermentation of cow manure as a peat substitute in horticulture - physical and chemical characteristics. *Acta Hort.* 150:553-561.
- Chen, Y., Y. Inbar, and M. Raviv. 1984b. Slurry produced by methanogenic fermentation of cow manure as a peat substitute in horticulture. *Proc. 2nd. Int. Peat in Agriculture and Horticulture*, The Hebrew University, Jerusalem pp.297-317.
- Chen, Y., and Y. Hadar. 1987. Composting and use of agricultural wastes in container media. p. 71-77. In M. de Bertoldi, M.P. Ferranti, P. L'Hermite and F. Zucchini (Eds.) *Compost: Production, Quality and Use*. Elsevier Applied Science, London.
- Chen, Y., Y. Inbar, and Y. Hadar. 1988. Composted agricultural wastes as potting media for ornamental plants. *Soil Sci.* 145:298-303.
- Chen, Y., and T. Aviad. 1989. Effects of humic substances on plant growth. In P. MacCarthy, C.E. Clapp, R.L. Malcolm, and P.R. Bloom (Eds.) *Humic substances in Soil and Crop Science*. Am. Soc. Agron., Soil Sci. Am., Madison, Wisconsin. (in press).
- Chen, Y., Y. Inbar and Y. Hadar. 1992. Composted residues reduce peat and pesticide use. *BioCycle* 6:48-51.
- Chen, Y., and Y. Inbar. 1993. Chemical and spectroscopical analyses of OM transformations during composting in relation to compost maturity. p. 551-600. In

- H.A.J. Hoitink and H.M. Keener (Eds) Science and Engineering of Composting. Renaissance Publications, Worthington, OH.
- Chen, Y., H. Magen, and J. Riov. 1994. Humic substances originating from rapidly decomposing OM: properties and effects on plant growth. p. 427-443. In N. Senesi and T.M. Miano, (Eds.) Humic Substances in the Global Environment and Implication on Human Health.. Elsevier, Amsterdam.
- Ciavatta, C., M. Govi, L. Pasotti and P. Sequi. 1993. Changes in OM during stabilization of compost from municipal solid wastes. Bio. Tech. 43:141-145.
- Cull, D.C. 1981. Alternatives to peat as container media. Organic resources in the U.K.. Acta Hort. 126:69-81.
- de Bertoldi, M., M.P. Ferranti, P. L'Hermite, and F. Zucconi (Eds). 1987. Compost: Production, Quality and Use. Elsevire, London.
- de Bertoldi, M. 1993. MSW composting challenges in Europe. BioCycle 34:75-76.
- De Vleeschauwer, D., O. Verdonck, and P. VanAssche. 1981. Phytotoxicity of refuse compost. BioCycle 22:44-46.
- Diaz, L.F., M.S. Savage, and C.G. Golueke. 1982. Composting. p. 95-120. In Resource Recovery from Municipal Solid Wastes. Volume II - Final processing. CRC Press Inc., Boca Raton, Florida.
- Eflosen, R.M., J.A. Ripmeester, N. Cyr, L.P. Milligan, and G. Mathison. 1984. Nutritional evaluation of forages by high-resolution solid-state ¹³C-NMR. Can. J. Anim. Sci. 64:93-102.
- Finstein, M.S., and M.L. Morris. 1975. Microbiology of municipal solid waste composting. Adv. Appl. Microbiol. 19:113-151.
- Finstein, M.S. 1992. Composting in the context of municipal waste management. p. 355-374. In R. Mitchell, (Ed.) Environmental Microbiology. Wiley-Liss, New York.
- Garcia, C., T. Hernandez, and F. Costa. 1992. Comparison of humic acids derived from city refuse with more developed humic acids. Soil Sci. Plant Nutr. 38:339-346.
- Gerasimowicz, W.V., and D.M. Byler. 1985. Carbon-13 CPMAS NMR and FTIR spectroscopic studies of humic acids. Soil Sci. 139:270-278.
- Gerasimowicz, W.V., D.M. Byler, and H. Susi. 1986. Resolution-enhanced FT-IR spectra of soil constituents: Humic Acid Applied Spectroscopy 4:504-507.

- Goldstein, N., and R. Steuteville. 1995. Solid waste composting plants in a steady state. *BioCycle* 36:48-57.
- Gray, K.R., and A.J. Biddlestone. 1981. The composting of agricultural wastes. p. 99-111. In B. Stonehouse (Ed.) *Biological Husbandry*. Butterworths, London.
- Grebus, M.E., M.E. Watson, and H.A.J. Hoitink. 1994. Biological, chemical and physical properties of composted yard trimmings as indicators of maturity and plant disease suppression. *Compost Science and Utilization* 2:57-71.
- Griffiths, P.R. 1983. Fourier transform infrared spectroscopy. *Science* 222:297-302.
- Hadar, Y., Y. Inbar, and Y. Chen. 1985. Effect of compost maturity on tomato seedling growth. *Scientia Horticulturae* 27:199-208.
- Hang, Y.D. 1988. Recovery of food ingredients from grape pomace. *Process Biochemistry* February 1988:2-4.
- Harada, Y., and A. Inoko. 1980a. The measurement of cation exchange capacity of composts for the estimation of the degree of maturity. *Soil. Sci. Plant Nutr.* 26:127-134.
- Harada, Y., and A. Inoko. 1980b. Relationship between cation-exchange capacity and degree of maturity of city refuse composts. *Soil Sci. Plant Nutr.* 26:353-362.
- Harada, Y., A. Inoko, M. Tadaki, and T. Izawa. 1981. Maturing process of city refuse compost during piling. *Soil Sci. Plant Nutr.* 27:357-364.
- Hatcher, P.G., M. Schnitzer, L.W. Dennis, and G.E. Maciel. 1981. Aromaticity of humic substances in soils. *Soil Sci. Soc. Am. J.* 45:1089-1094.
- Hatcher, P.G., I.A. Breger, L.W. Dennis, and G.E. Maciel. 1983. Solid-state ^{13}C -NMR of sedimentary humic substances: new revelations on their chemical composition. p. 37-81. In R.F. Christman and E.T. Gjessing (Eds.) *Aquatic and Terrestrial Humic Materials*. Ann Arbor Science, Ann Arbor, MI.
- Haw, J.F., G.A. Maciel, and H.A. Schroder. 1984. Carbon-13 nuclear magnetic resonance spectrometric study of wood and wood pulping with cross polarization and magic-angle spinning. *Anal. Chem.* 56:1323-1329.
- Hayes, M.H.B., R.S. Swift, R.E. Wardle, and J.K. Brown. 1975. Humic materials from an organic soil: a comparison of extractants and of properties of extracts. *Geoderma* 13:231-245.

- Hayes, M.H.B. 1985. Extraction of humic substances from soil. p. 329-362. In G.R. Aiken, D.M. McKnight, R.L. Wershaw and P. MacCarthy (Eds.) *Humic Substances in Soil, Sediment, and Water*. John Wiley & Sons, New York.
- He, X.T., S.J. Traina, and T.J. Logan. 1992. Chemical properties of municipal solid waste composts. *J. Environ. Qual.* 21:318-329.
- Himmelsbach, D.S., and F.E. Barton II. 1980. ¹³C-nuclear magnetic resonance of grass lignins. *J. Agric. Food Chem.* 28:1203-1208.
- Hirai, M.F., A. Katayama, and H. Kubota. 1986. Effect of compost maturity on plant growth. *BioCycle* 27(4):58-61.
- Hoitink, H.A.J., M.J. Boehm, and Y. Hadar. 1993. Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. p. 601-622. In H.A.J. Hoitink and H.M. Keener, (Eds.) *Science and Engineering of Composting*. Renaissance Publications, Worthington, OH.
- Hoitink, H.A.J., and G.A. Kuter. 1986. Effects of composts in growth media on soilborne pathogens. p. 289-306. In Y. Chen and Y. Avnimelech (Eds.) *The Role of OM in Modern Agriculture*. Martinus Nijhoff Publishers, Dordrecht.
- Hoitink, H.A.J., and P.C. Fahy. 1986. Basis for the control of soilborne plant pathogens with composts. *Ann. Rev. Phytopathol.* 24:93-114.
- Huijsmans, J., and J.A. Lindley. 1984. Evaluation of a solid-liquid separator. *Trans. ASAE* :1854-1858.
- Iannotti, D.A., M.E. Grebus, B.L. Toth, L.V. Madden, and H.A.J. Hoitink. 1994. Oxygen respirometry to assess stability and maturity of composted municipal solid waste. *J. Environ. Qual.* 23:1177-1183.
- Inbar, Y., Y. Chen, and Y. Hadar. 1985. The use of composted slurry produced by methanogenic fermentation of cow manure as growth media. *Acta Hort.* 172:75-85.
- Inbar, Y., Y. Chen, and Y. Hadar. 1988. Composting of agricultural wastes for their use as container media: simulation of the composting process. *Bio. Wastes* 26:247-259.
- Inbar, Y., Y. Chen, and Y. Hadar. 1989. Solid-state carbon-13 nuclear magnetic resonance and infrared spectroscopy of composted organic matter. *Soil Sci. Soc. Am. J.* 53:1695-1701.

- Inbar, Y., Y. Chen, and Y. Hadar. 1990a. Humic substances formed during the composting of OM. *Soil Sci. Soc. Am. J.* 54:1316-1323.
- Inbar, Y., Y. Chen, Y. Hadar, and H.A.J. Hoitink. 1990b. New approaches to compost maturity. *BioCycle* 31:64-69.
- Inbar, Y., Y. Hadar, and Y. Chen. 1993. Recycling of cattle manure: the composting process and characterization of maturity. *J. Environ. Qual.* 22:257-863.
- Inoko, A. 1982. The composting of organic materials and associated maturity problems. ASPAC, Food & Fertilizer Technology Center, Technical Bulletin No. 71.
- Jimenez, E.I., and V.P. Garcia. 1992. Determination of maturity indices for city refuse composts. *Agriculture, Ecosystems and Environment* 38:331-343.
- Kogel-Knabner, I., P.G. Hatcher and W. Zech. 1991. Chemical structural studies of forest soil humic acids: aromatic carbon fraction. *Soil Sci. Soc. Am. J.* 55:241-247.
- Kolodziecki, W., J.S. Frye, and G.E. Maciel. 1982. Carbon-13 nuclear magnetic resonance spectrometry with cross polarization and magic-angle spinning for analysis of lodgepole pine wood. *Anal. Chem.* 54:1419-1424.
- Kuter, G.A., H.A.J. Hoitink, and L.A. Rossman. 1985. Effect of aeration and temperature on composting of municipal sludge in full-scale vessel system. *Journal WPCF* 57(4):309-315.
- Kwok, O.C.H., P.C. Fahy, H.A.J. Hoitink and G.A. Kuter. 1987. Interactions between bacteria and Trichoderma hamatum in suppression of Rhizoctonia damping-off in bark compost media. *Phytopathology* 77:1206-1212.
- Lax, A., F. Costa, A. Roig, M.T. Hernandez, and J. Cegarra. 1987. Indexes of humification grade in manures. *The Science of the Total Environment* 62:457-458.
- Levanon, D., C. Dosoretz, B. Motro, and I. Kahn. 1984. Recycling agricultural waste for mushroom casing. *Mushroom J.* 133:13-17.
- Lobartini, J.C., and K.H. Tan. 1988. Differences in humic acid characteristics as determined by carbon-13 nuclear magnetic resonance, scanning electron microscopy, and infrared analysis. *Soil Sci. Soc. Am. J.* 52:125-130.
- Lohr, V.I., R.G. O'Brien, and D.L. Coeffey. 1984. Spent mushroom compost in soilless media and its effects on the yield and quality of transplants. *J. Amer. Soc. Hort. Sci.* 109:693-697.

- Lynch, J.M., K.B. Gunn, and L.M. Panting. 1980. On the concentration of acetic acid in straw and soil. *Plant and Soil* 56:93-98.
- MacCarthy, P., and J.A. Rice. 1985. Spectroscopic methods (other than NMR) for determining functionality in humic substances. p. 527-543. In G.R. Aiken, D.M. McKnight, R.L. Wershaw and P. MacCarthy (Eds.) *Humic Substances in Soil, Sediment, and Water*. John Wiley & Sons, New York.
- Malcolm, R.L. 1989. Applications of solid-state ^{13}C -NMR to geochemical studies. In M.H.B. Hayes, P. MacCarthy, R. Swift, and R.L. Malcolm (Eds.) *Humic Substances in Soil, Sediments and Water: II. In Search of Structure*. John Wiley, New-York. (in press).
- Mandelbaum, R., Y. Hadar, and Y. Chen. 1988. Composting of agricultural wastes for their use as container media: effect of heat treatments on suppression of Pythium aphanidermatum and microbial activities in substrates containing compost. *Biol. Wastes* 26:261-274.
- Morel, J.L., F. Colin, J.C. Germon, P. Godin, and C. Juste. 1985. Methods for the evaluation of the maturity of municipal refuse. p. 56-72. In J.K.R. Gasser (Ed.) *Composting of Agricultural and Other Wastes*. Elsevier Applied Science Publishers, London.
- Nakasaki, K., M. Sasaki, M. Shoda, and H. Kubota. 1985. Change in microbial numbers during thermophilic composting of sewage sludge with reference to CO_2 evolution rate. *Appl. Environ. Microbiol.* 49:37-41.
- Niemeyer, J., Y. Chen and J.M. Bollag. 1992. Characterization of humic acids, composts, and peat by diffuse reflectance Fourier transform infrared spectroscopy. *Soil Sci. Soc. Am. J.* 56:135-140.
- Packer, K.J., R.K. Harris, A.M. Kenwright, and C.E. Snape. 1983. Quantitative aspects of solid state ^{13}C -NMR of coals and related materials. *Fuel* 62:999-1002.
- Pandeya, S.B. 1992. Characterization of fulvic acids extracted from some organic manures and wastes by potentiometric titration. *Biol. Tech.* 39:77-83.
- Piotrowski, E.G., K.M. Valentine, and P.E. Pfeffer. 1984. Solid-state, C-13, cross-polarization, "magic-angle" spinning, NMR spectroscopy studies of sewage sludge. *Soil Sci.* 137:194-203.

- Preston, C.M., and J.A. Ripmeester. 1983. ^{13}C -labelling for NMR studies of soils: CPMAS observation of ^{13}C -acetate transformation in a mineral soil. *Can. J. Soil Sci.* 63:495-500.
- Preston, C.M., J.A. Ripmeester, S.P. Mathur, and M. Levesque. 1986. Application of solution and solid-state multinuclear NMR to a peat-based composting system for fish and crab scrap. *Can. J. Soil Sci.* 31:63-69.
- Preston, C.M., S.E. Shipitalo, R.L. Dudley, C.A. Fyfe, S.P. Mathur, and M. Levesque. 1987. Comparison of C-13 CPMAS NMR and chemical techniques for measuring the degree of decomposition in virgin and cultivated peat profiles. *Can. J. Soil Sci.* 67:187-198.
- Preston, C.M., R.H. Hewman, and P. Rother. 1994. Using ^{13}C CPMAS NMR to assess effects of cultivation on the OM of particle size fractions in a grassland soil. *Soil Science* 157:26-35.
- Raviv, M., Y. Chen, Z. Geler, S. Medina, E. Putievski, and Y. Inbar. 1984. Slurry produced by methanogenic fermentation of cow manure as a growth medium for some horticultural crops. *Acta Hort.* 150:563-573.
- Raviv, M., Y. Chen, and Y. Inbar. 1986. Peat and peat substitutes as growth media for container-growth plants. p. 257-287. In Y. Chen and Y. Avnimelech (Eds.) *The Role of OM in Modern Agriculture*. Martinus Nijhof, The Hague, The Netherlands.
- Roletto, E., R. Barberis, and V. Zelano. 1982. Gel filtration and absorption spectroscopic investigations on humic substances from organic fertilizers. *Plant and Soil* 66:383-390.
- Roletto, E., and P. Ottino. 1984. Analytical characterization of humic substances from composted lignocellulosic residues. *Agricultural Wastes* 11:181-195.
- Roletto, E., R. Chiono, and E. Barberis. 1985. Investigation on humic matter from decomposing poplar bark. *Agricultural Wastes* 12:261-272.
- Roletto, E., and M.P. Luda. 1987. Chemical and infra-red spectroscopic study of humic and fulvic acids from composted poplar bark. *Biol. Wastes* 19:205-214.
- Saviozzi, A., R. Levi-Minzi, and R. Riffaldi. 1988. Maturity evaluation of organic wastes. *BioCycle* 29 (3):54-56.
- Schnitzer, M. 1978. Humic substances: chemistry and reactions. p. 1-64. In M. Schnitzer and S.U. Khan (Eds.) *Soil Organic Matter*. Elsevier, Amsterdam.

- Sequi, P., M. De Nobili, L. Leita, and G. Cercignani. 1986. A new index of humification. *Agrochimica* 30:175-179.
- Steuteville, R. 1994a. The state of garbage in America (II). *BioCycle* 35:30-36.
- Steuteville, R. 1994b. The state of garbage in America (I). *BioCycle* 35:46-52.
- Steuteville, R. 1995. The state of garbage in America (I). *BioCycle* 36:54-58.
- Stevenson, F.J. 1982. *Humus Chemistry*. John Wiley & Sons, Inc.
- Stevenson, F.J. 1985. Geochemistry of soil humic substances. p. 13-52. In G.R. Aiken, M.D. McKnight, R.L. Wershaw, and P. MacCarthy (Eds.) *Humic Substances in Soil, Sediments and Water*. John Wiley & Sons.
- Sugahara, K., and A. Inoko. 1981. Composition analysis of humus and characterization of humic acid obtained from city refuse compost. *Soil Sci. Plant Nutr.* 27:213-224.
- Toussan, T.A., A.R. Weinhold, R.G. Linderman, and Z.A. Patrick. 1968. Nature of phytotoxic substances produced during plant residue decomposition in soil. *Phytopathology* 58(1):41-45.
- Vaughan, D., and R.E. Malcolm. 1985. Influence of humic substances on growth and physiological processes. p. 37-75. In D. Vaughan and R.E. Malcolm (Eds.) *Soil Organic Matter and Biological Activity*. Martinus Nijhoff/Dr. Junk Publishers, Dordrecht.
- Verdonck, O. 1984. Reviewing and evaluation of new materials used as substrates. *Acta Horticulturae* 150:467-473.
- Watanabe, A., K. Itoh, S. Arai, and S. KuWatsuka. 1994. Comparison of the composition of humic and fulvic acids prepared by the IHSS method and NAGOYA method. *Soil Sci. Plant Nutr.* 40:601-608.
- Wershaw, R.L. 1985. Application of nuclear magnetic resonance for determining functionality in humic substances. p. 561-584. In G.R. Aiken, D.M. McKnight, R.L. Wershaw and P. MacCarthy. (Eds.) *Humic Substances in Soil, Sediment and Water*. Wiley & Sons Inc., New York.
- Wershaw, R.L., K.A. Thorn, D.J. Pinckney, P. MacCarthy, J.A. Rice, and H.F. Hemond. 1986. Application of membrane model to the secondary structure of humic materials in peat. p. 133-157. In C. Fuchsman (Ed.) *Peat and Water: Aspects of Water Retention and Dewatering in Peat*. Elsevier Applied Science Publishers Ltd.

- Wershaw, R.L., K.A. Thorn, and D.J. Pinckney. 1988. Characterization of humic acid fractions by C-13 nuclear magnetic resonance spectroscopy. *Environ. Tech. Letters* 9:53-62.
- Wilson, M.A., S. Heng, K.M. Goh, R.J. Pugmire, and D.M. Grant. 1983. Studies of litter and acid-insoluble soil OM fractions using ^{13}C cross polarization nuclear magnetic resonance spectroscopy with magic angle spinning. *J. Soil Sci.* 34:83-97.
- Wilson, M.A. 1987. NMR techniques and applications in geochemistry and soil chemistry. Pergamon Press, Headington Hill Hall, Oxford, England.
- Wong, M.H., and L.M. Chu. 1985. Changes in properties of a fresh refuse compost in relation to root growth of *Brassica chinensis*. *Agricultural Wastes* 14:115-125.
- Zech, W.J.M.B., L. Hanmaier, and R.L. Malcolm. 1987. CPMAS ^{13}C NMR and IR spectra of spruce and pine litter and of Kalsen lignin fraction at different stages of decomposition. *Z. Pflanzenernahr. Bodenk.* 150:262-265.
- Zucconi, F., M. Forte, A. Monaco, and M. de Bertoldi. 1981a. Biological evaluation of compost maturity. *BioCycle* 22:27-29.
- Zucconi, F., A. Pera, M. Forte, and M. de Bertoldi. 1981b. Evaluating toxicity of immature compost. *BioCycle* 22:54-57.
- Zucconi, F., and M. de Bertoldi. 1987. Compost specifications for the production and characterization of compost from municipal solid waste. p. 30-50. In M. de Bertoldi, M.P. Ferranti, P. L'Hermite and F. Zucconi (Eds.) *Compost: Production, Quality and Use*. Elsevier Applied Science, London.

Figures legends

Fig. 1.

Simplified flowchart of the composting process (after Gray and Biddlestone, 1981).

Fig. 2.

A schematic temperature and processes vs. time of composting.

Fig. 3.

Changes and trends in the humification parameters (expressed as percentage of variation) during the composting of wastewater sludge and straw mixtures (after Saviozzi et al., 1988)

Fig 4.

Structural components of lignin (the numbers indicates typical ^{13}C -NMR assignments).

Fig. 5.

Structures of anhydroglucose repeating unit and cellobiose (4-O-(β -D-glucosepyranosyl) β -D-glucopyranose).

Table 1.

Effects of composting of the solid fraction of a slurry produced from methanogenic fermentation of cattle manure on tomato seedling response (following Hadar et al., 1985).

Table 1. Effects of composting of the solid fraction of a slurry produced from methanogenic fermentation of cattle manure on tomato seedling response (following Hadar et al., 1985).

Treatment	Unfertilized	Fertilized
	Dry weight (mg/plant)	
Raw material	166b	30c
Immature compost	319a	486b
Mature compost	355a	612a

**FORMATION AND PROPERTIES
OF HUMIC SUBSTANCE
ORIGINATING FROM
COMPOSTS**

ABSTRACT

Composting enables a detailed evaluation of the humification process of various organic wastes within a short period of time (3 to 6 months). Humic substances (HS) constitute a large fraction of the organic matter (OM) in compost, and they are its most active fraction due to their effects on soil ecology, structure, fertility and metal complexes, and plant growth. The formation and properties of HS extracted from various composts such as municipal solid waste (MSW), grape marc (GM), composted separated cattle manure (CSM), sewage sludge (SS), wood compost (WC) and other organic wastes were studied. Degradative and non-degradative techniques (FTIR, DRIFT, ^{13}C -NMR) were used to study the transformation of HS during composting of the various organic wastes. Time-course studies of composting some of these wastes showed an increase in the relative amount of humic acid (HA) (from 18% to 45% of OM in CSM compost and from 5% to 20% in MSW compost), whereas the formation rate of fulvic acid (FA) was inconsistent. A humification ratio (HR - the ratio of HA/FA) was used to evaluate compost maturity. Values of 0.9 to 3.4 and 3.0 to 9.2 were typical for immature and mature composts, respectively. Another humification index (HI) used to define compost maturity was calculated as the ratio between non-humified fraction (NHF) and the humified fraction (HA+FA). A HI decrease during composting represents the formation of HS. Elemental- and functional-group analyses indicated only minor differences between HA extracted from composts at various stages. Moreover, these values fell into a wide range, similar to that of soil HA. The ^{13}C -NMR spectra of the HA exhibited strong bands representing aliphatic structures in various composts (50% of total C in CSM, 30% in MSW and 61% in GM) and a lower level of aromatic components (37% in CSM, 22% in MSW and 22% in GM). The FTIR spectra showed similar trends of strong aliphatic and carbohydrate components. Both techniques provided more qualitative information indicating that HA extracted from mature compost exhibits more aromatic structures

and carboxyl groups and less carbohydrate components than that from immature compost.

Studies on the effects of HS on plant growth showed stimulative effects. Typical response curves indicated enhanced growth with increasing HS concentration in solution, followed by decreases in growth at higher concentrations. In soils, the addition of composts was found to stimulate growth beyond that provided by mineral nutrients, presumably because of the effects of HS.

Introduction

Composting can play useful role in global waste management due to the increasing amount of decomposed organic materials such as animal excreta, municipal solid waste (MSW) and sewage sludge (SS) in modern society. Composting has been defined as a controlled microbial aerobic decomposition of heterogeneous raw waste passing through a thermophilic phase with a temporary release of phytotoxins, followed by the production of CO₂, H₂O, minerals and stabilized organic matter (OM) containing a high amount of humic substances (HS). The importance of HS to soil ecology, fertility and structure and their beneficial effects on plant growth (Chen and Aviad, 1990) have caused an increase in the use of compost as an amendment to soils or as a substitute substrate for peat in container media (Avnimelech et al., 1993; Chen et al., 1992; Hoitink, et al., 1993). HS originating from compost form within a shorter period of time than soil HS, therefore better understanding of the chemical and physical characterization and transformation of the compost HS is necessary to study the differences in the formation period. Moreover due to the increase interest in compost as substitute to peat in container media it is essential to study the interactions of HS in the compost-plant system.

The objectives of this article were (1) to compare the rate of decomposition of OM in various composts; (2) to study the rates of formation and decomposition of HS-

humic acid (HA), fulvic acid (FA) and the non-humic fraction (NHF), in various composts; (3) to use humification parameters as maturity indexes; (4) to characterize compost-derived HA using elemental- and functional-group analyses and spectroscopic measurements; (5) to study the effects of compost-derived HS on nutrient solubility and plant growth.

Humic Substance Content

Among the chemical criteria used to characterize compost maturity, HS content is one of the most common. The amounts of HS fractions in various composts at various stages of composting are reviewed here. The main trends of several humification parameters over a period of 140 days of composting of wastewater sludge and straw mixtures are shown in Fig. 1 (Saviozzi et al., 1988). The following humification indexes (HI) were used to defined compost maturity: (1) HS- amount of total humic substances (HA, FA and NHF) as percent of OM; (2) HA- amount of humic acid as percent of OM; (3) FA- fulvic acid as percent of OM; (4) HR- the HA/FA ratio; (5) amount of the humic materials (HA+FA); (6) HI- the ratio between the NHF and the HA+FA fractions (Sequi et al., 1986); (7) HAp- humic acid percent of HS.

Total HS content follows different trends during the composting of different composts. HS content does not vary during composting of MSW (Sugahara and Inoko, 1981; Das, 1988; Jimenez and Garcia, 1992) These authors found that alkaline extraction of MSW compost at various stages of composting gives a steady value in the range of 19 to 22% of OM. A possible explanation for the presence of HS in the early stages of composting and for the stability of the HS content in MSW compost could be the relatively high content of NHF. Therefore HS in MSW compost cannot be used as an indicator of maturity. In contrast, during the composting of CSM, HS content doubled (from 377 to 710 g/kg OM). The different trends between the composts can be explained by the relatively high amount of HA in CSM (46% of OM),

whereas both HA and FA fractions in MSW compost amount to less than half of the total OM. The trend in HA levels during composting is similar in all composts, its content increasing during composting and reaching up to 46% and 48% of OM in CSM and poplar bark compost, respectively (Roletto et al., 1985; Inbar et al., 1990). In MSW compost, HA increased from 7% - 8% in the raw material to 12% - 14% in mature compost (Jimenez and Garcia, 1992). While the levels of HA increases the amount of FA usually decreases and the HA/FA ratio is therefore often used as an indicator of maturity. This trend of an increasing HA/FA ratio from fresh to composted OM has been reported for poplar bark (Roletto et al., 1985), farmyard manure (FYM), urban refuse (Saviozzi et al., 1988), CSM (Inbar et al., 1990) and MSW (Sugahara and Inoko, 1981; Das, 1988; Jimenez and Garcia, 1992; Gonzales et al., 1993). The value of the HR differs for various wastes but it generally increases by more than 80% during composting, indicating a high degree of OM polymerization, since more polymerized components are found in the mature compost.

Another parameter indicating the role of HS is the HI ($\text{NHF}/(\text{HA}+\text{FA})$) which decreases during composting due to humification of OM (Saviozzi et al., 1988; Inbar et al., 1990). The trend of decreasing HI from fresh to composted materials is similar in all wastes, but the rate of decrease changes according to the amount of NHF in the waste. The HI of grape marc (GM), containing a high level of sugars, drops sharply during composting, in contrast to the HI in CSM (Inbar, 1989). To summarize a large amount of HA forms while FA level usually decreases. The HI and the HR can only be used as maturity parameters when NHF is separated from the fulvic fraction, otherwise these parameters have no meaning.

Elemental- and Functional-Group Analyses:

The elemental composition, atomic ratios and functional groups of HA extracted from various composts as compared to "model" HA (Schnitzer, 1978) and Leonardite HA are presented in Table 1. The average compositional range of HA

extracted from various composts was similar to those of "model" and Leonardite HA. The C content of HA isolated from compost was lower than that of "model" HA, except in the cases of CSM and GM. This parameter does not change during composting (Roletto and Luda, 1987; Inbar, 1989; Garcia et al., 1992). H content was higher in compost HA than in Leonardite or "model" HA, although it was slightly reduced during composting of CSM and GM. N content was also higher in compost HA. The high N and H contents in the composts may be due to the presence of side chains containing those components. The highest contents of N and H were found in MSW compost (Garcia et al., 1989; Deiana et al., 1990; Garcia et al., 1992), which yielded a relatively low C/N ratio as compared to HA formed in soil or during composting of other stabilized wastes. The H/C ratio represents HA aromaticity (a low ratio indicates an aromatic structure). All compost HA had higher H/C ratios than the "model" or Leonardite HA, indicating that aromatic structures in compost HA are less pronounced and therefore compost HA has identified as "young" HA.

The major differences between "model" or Leonardite HA and compost HA were found in the functional-group analysis. Compared to soil and sediment HA, compost HA contained less fractional oxygen groups (carboxyl, carbonyl and phenolic carbon), due to the relatively short formation time during composting as compared to that required for soil OM transformation. During composting, the carboxyl group content increased (192 to 221 meq/100g in CSM, 176 to 224 meq/100g in MSW, 184 to 264 meq/100g in SS). The carbonyl group content showed the same trend, whereas that of the phenolic group decreased during composting. The total acidity (calculated as phenolic plus carboxyl groups) behaved differently in the various composts: it increased in composts in which the raw material had a stable structure (wood and cattle manure) but decreased in composts in which the raw materials contained more rapidly biodegradable fraction such as city refuse, yard waste and sludge (Roletto and Luda, 1987; Garcia et al., 1992; Kakezaw, 1992). There were no major differences between the elemental composition of HA from the various composts at various stages

of composting and that of "model" HA (Table 1). In general, the elemental and functional groups were similar to "young forms" of HA reported in the literature, even though HA amounts increased during composting.

The E_4/E_6 ratio was higher in compost HA, indicating a lower molecular weight than that of "model" HA (Chen et al., 1977). Elemental and functional group analyses of FA have been reported in only a few works (Roletto and Luda, 1987; Pandey, 1992). The compost FA had lower C and N contents but higher amounts of O and carboxyl groups than HA, and the overall elemental and functional groups extracted from compost were similar to "model" FA (Schnitzer, 1978; Roletto and Luda, 1987; Pandey, 1992).

Infrared Spectroscopy (FTIR, DRIFT):

Infrared spectroscopy was used to characterize the HS. FTIR and DRIFT spectroscopy served as qualitative tools to characterize the chemical groups and bands of the HA complex structures. Sugahara and Inoko (1981) investigated HA extracted from city refuse at various stages of composting by IR spectroscopy and found that the aliphatic C-H and C-O stretching decreased while the carboxyl COO⁻ increased during composting. The main absorbance bands were: a broad band at 3400-3300 cm⁻¹ (H-bonded OH groups); a slight shoulder at 3085-3075 cm⁻¹ (aromatic C-H stretching); two peaks at 2930 and 2850 cm⁻¹ (aliphatic C-H stretching); a slight shoulder at 1730-1715 cm⁻¹ (C=O of carboxyl, C=O of ketonic carbonyl); a well-pronounced peak at 1610-1655 cm⁻¹ (aromatic C=C stretching); a sharp peak at 1540-1560 (amide II); a peak at 1460-1450 cm⁻¹ (-CH deformation of -CH₃ and -CH bending of CH₂); a peak at 1420 cm⁻¹ (aromatic ring stretch); a peak at 1400-1350 cm⁻¹ (aliphatic C-H bending); a peak at 1270-1220 cm⁻¹ (aromatic C-OH and -C-OH stretch and OH deformation of -COOH); a peak around 1100-1020 cm⁻¹ (C-O stretch of polysaccharides). The interpretations of the IR spectra are based on Inbar et al. (1990), Niemeyer et al.

(1992), Deiana et al. (1990), Sugahara and Inoko (1981), Garcia et al. (1992), Roletto and Luda (1987) and Stevenson (1982).

HA spectra from the various mature composts studied here exhibited similar features: all showed a strong aliphatic nature with aromatic structure and peptide bonds, along with relatively low carboxyl content, similar to young type III HA extracted from fresh soil OM (Stevenson and Goh, 1971). HA extracted at several stages during the composting of various wastes exhibited only minor changes. The main ones consisted of slight reductions of the aliphatic and polysaccharide bands during composting of SS and MSW (Sugahara and Inoko, 1981; Garcia et al., 1989, 1992), whereas during composting of CSM, no changes were observed; during the composting of GM there was a relative increase in aliphatic carbon and a decrease in polysaccharides. The minor changes in HA extracted from CSM during composting are presented in Fig. 2.

CPMAS ^{13}C -NMR Spectroscopy

The use of NMR spectroscopy to study HA has gained prominence over the last few years. Wershaw (1985) showed that although NMR analysis is only applied for a short time, it can provide more definitive structural information than other methods. The ^{13}C -NMR spectra were divided into the following regions (Hatcher et al., 1983; Malcolm, 1989): 0-50 ppm (alkyl C-methyl, methylene and methine groups), 50-60 ppm (methoxyl C), 60-70 ppm (alcohol), 70-98 ppm (C-O, C-N), 98-112 ppm (anomeric C), 112-145 ppm (aromatic C), 145-163 ppm (phenolic C), 163-190 ppm (carboxyl groups), 190-215 ppm (carbonyl C). The 0-112 ppm region was calculated as aliphatic C, 60-112 ppm as polysaccharide C, 112-163 ppm as aromatic C. Total aromaticity was calculated by expressing aromatic C (112-165 ppm) as percentage of aliphatic C (0-112 ppm) plus aromatic C. CPMAS ^{13}C -NMR spectra of HA extracted from CSM is presented in Fig. 3. There were only minor differences between the spectra of HA from the fresh and composted manure (12% increase in carboxyl

content; 8% increase in aromatic carbon and 8% decrease in alkyl carbon), similar to the FTIR spectra (Inbar, 1989). During the composting of MSW, changes in HA structure were observed: the HA aromatic carbon increased by 39% while the aliphatic carbon content decreased by 19% (Garcia et al., 1992). Opposite changes occurred in GM HA: while the alkyl region increased, the aromatic carbons decreased, indicating highly oxidizing conditions (Inbar, 1989). The main differences between HA extracted from various composts are present in Table 2. CSM HA seemed to be more aromatic and less aliphatic than MSW and GM HA. The higher aromaticity of the CSM HA can be explained by the high lignin content in this compost (61.5% of dry weight), whereas GM and MSW contain more easily biodegradable OM. These differences in HA structure lead to the conclusion that the raw materials components influence not only the time of composting but also the formation of HS.

Humic Substance Effects on Plant Growth

Applying OM as a fertilizer increased plant yield to a higher level than that resulting from the application of the same level of inorganic fertilizer, presumably because of the effects of the HS. In soils, the addition of compost can stimulate growth beyond that provided by mineral nutrient supplements alone (Senesi, 1989; Chen et al., 1994). This improvement can be explained by direct short-term fertilizer effects and some indirect effects. Table 3 presents the typical effects of HA stimulation on wheat growth when given in water or nutrient solution. Addition of FA to the nutrient solution yielded an atypical response curve whereby cucumber root and shoot weights increased with increasing FA concentration up to a FA concentration where shoot and root weights decrease. Chen et al. (1994) found the same trend in melon shoots and roots with increasing concentrations of HS originating from a water extract of CSM. Both experiments showed that higher OM levels result in decreased growth, possibly resulting from a suboptimal concentration of free micronutrients in solution due to an excessive amount of chelators (HS). Complexation of transition metals, such

as Cu, Zn, Fe and Mn by HS has been the focus of numerous publications. The solubility of most of the micronutrients in their inorganic forms is limited in soil or nutrient solution. The presence of HS may contribute to improved availability of these elements (Chen and Aviad, 1990). HS not only increased the solubility of Fe in solution but they also affected Fe transport from roots to shoots. Chen et al's (1994) work on HS extracted from CSM indicates that residual concentrations of Fe, Mn and Cu in nutrient solutions that were equilibrated at pH 6.5 and 7.0 increased with increasing concentrations of added OM. With regard to plant nutrition, this data suggests that at low pH, HS may not exhibit beneficial effects, but at pH 7.5 or higher, a high concentration of HS is required to maintain micronutrient availability. The stimulation of ion uptake by the addition of HS led to the proposal that these materials affect membrane permeability related to surface activity of the HS. Another phenomenon related to CSM and GM HS is hormone-like activity (Chen et al., 1994). When mung-bean cuttings were placed in nutrient solutions containing OM extracted from CSM, GM and Leonardite, root weight and number were significantly higher as compared to the control. This result supports those of other plant growth assays, in which HS added to a nutrient solution enhanced root formation activity, which could be considered an auxin-like activity. To summarize, the effects of HS on plant growth can be divided into direct and indirect, as follows (Chen and Aviad, 1990):

Direct effects:

1. Effects on membranes resulting in improved transport of nutritional elements
2. Enhanced protein synthesis
3. Hormone-like activity
4. Enhanced photosynthesis
5. Effects on enzyme activity

Indirect effects:

1. Solubilization of microelements (Fe, Zn, Cu and Mn) and some macroelements (K, Ca, P)

2. Reduction of active levels of toxic elements
3. Enhancement of microbial populations

Conclusions

1. The level of HS formation reached a maximum within 90 to 120 days of composting without forced aeration.
2. HS level increased with composting time in CSM and wood compost (WC), but remained steady in MSW compost.
3. HA level increased in CSM, WC, MSW and SS composts.
4. FA level remained steady in CSM and decreased in MSW composts.
5. NHF level decreased with composting time.
6. Elemental- and functional-group analyses indicated only minor differences between HA extracted from composts at various stages of composting.
7. Values of C, H and S in HA derived from various composts were similar to those found in soils. However, the N levels were higher in compost-derived HA and the C/N ratio was therefore lower.
8. Total acidity, -COOH and phenolic OH levels were lower in HA derived from composts than the average numbers from soil HA.
9. FTIR and NMR spectra of compost-derived HA were similar, regardless of composting time.
10. FTIR and NMR spectra of compost-derived HA exhibited high levels of polysaccharides and aliphatic structures.
11. Biological activity of compost-derived HS resembled that of soil-derived HS.
12. Plant-hormone-like activity attributed to HS does not involve auxins, cytokinins or abscisic acid.
13. Growth enhancement of plants in nutrient solution and soils by HS may be attributed to the maintenance of Fe, Mn, Cu and Zn in solution at sufficient levels. This effect is pH-dependent, its effectiveness decreasing with increasing pH.

14. The commonly cited conclusion (including 1992 publications) stating that:
"Stimulatory effects in nutrient solution of HS can't be ascribed to a supply of
elements" may be incorrect.

REFERENCES

- Avnimelech, Y., A. Cohen and D. Shkedy. 1993. Can we expect a consistent efficiency of municipal waste compost application? *Compost Science and Utilization* 1:7-15.
- Chen, Y., Senesi, S., and Schnitzer, M., 1977. Information provided on humic substances by E_4/E_6 ratios. *Soil Sci. Soc. Amer. J.* 41, 352-358.
- Chen, Y., and Aviad, T., 1990. Effects of humic substances on plant growth. In: P. MacCarthy, C.E. Clapp, R.L. Malcolm and P.R. Bloom (eds.), *Humic Substances in Soil and Crop Sciences: Selected Readings. Proceedings of a symposium cosponsored by the International Humic Substances Society, Chicago, Illinois*, 161-186.
- Chen, Y., Y. Inbar and Y. Hadar. 1992. Composted residues reduce peat and pesticide use. *BioCycle* 6:48-51.
- Chen, Y., Magen, H., and Riov, J., 1994. Humic substances originating from rapidly decomposing organic matter: properties and effects on plant growth. In: N. Senesi, and T.M. Miano (eds.), *Humic Substances in the Global Environment and Implications on Human Health*. Elsevier Science, Amsterdam, 427-445.
- Das, A., 1988. City garbage compost as a source of humus. *Biological Wastes* 26, 65-69.
- Deiana, S., Gessa, C., Manunza, B., Rausa, R., and Seeber, R., 1990. Analytical and spectroscopic characterization of humic acids extracted from sewage sludge, manure, and worm compost. *Soil Science* 150, 419-424.
- Garcia, C., Hernandez, T., Costa, F., and Rio, J.C., 1989. Study of the lipid and humic fractions from organic wastes before and after the composting process. *Sci. Tot. Environ.* 81 & 82, 551-560.
- Garcia, C., Hernandez, T., and Costa, F., 1992. Comparison of humic acids derived from city refuse with more developed humic acids. *Soil Sci. Plant Nutr.* 38, 339-344.

- Gonzales, S.J., Carballas, M., Villar, M.C., Beloso, M.C., Cabaneiro, A., and Carballas, T., 1993. Carbon and nitrogen containing compounds in composted urban refuses. *Biores. Technol.* 45, 115-121.
- Hatcher, P.G., Breger, I. A., Dennis, L.W., and Maciel, G.E., 1983. Solid-state ^{13}C -NMR of sedimentary humic substances: new revelation on their chemical composition. In: R.F. Christman and E.T. Gjessing (eds.), *Aquatic and Terrestrial Humic Materials*. Ann Arbor Science Publications Inc., Ann Arbor, MI, p. 37-81.
- Hoitink, H.A.J., M.J. Boehm, and Y. Hadar. 1993. Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. p. 601-622. *In Science and Engineering of Composting*. H.A.J. Hoitink and H.M. Keener, (ed.). Renaissance Publications, Worthington, OH.
- Inbar, Y., 1989. Formation of humic substances during composting of agricultural wastes and characterization of their physico-chemical properties. Ph.D. Thesis, The Faculty of Agriculture, The Hebrew University of Jerusalem.
- Inbar, Y., Chen, Y., and Hadar, Y., 1990. Humic substances formed during the composting of organic matter. *Soil Sci. Soc. Amer. J.* 54, 1316-1323.
- Jimenez, E., and Garcia, P., 1992. Determination of maturity indices for city refuse composts. *Agric. Ecosyst. Environ.* 38, 331-343.
- Takezawa, M., Nishida, T., and Takahara, Y., 1992. Structural characteristics of humic acids extracted from woody composts by two-steps composting process. *Soil Sci. Plant Nutr.* 38, 85-92.
- Malcolm, R.L., 1989. Applications of solid-state ^{13}C -NMR to geochemical studies. In: M.H.B. Hayes, P. MacCarthy, R. Swift and R.L. Malcolm (eds.), *Humic Substances in Soil, Sediments and Water: II. In Search of Structure*. John Wiley & Sons, New York. p. 339-372.

- Niemeyer, J., Chen, Y., and Bollag, J.M., 1992. Characterization of humic acids, composts, and peat by diffuse reflectance Fourier-transform infrared spectroscopy. *Soil Sci. Soc. Amer. J.* 56, 135-140.
- Pandey, S.B., 1992. Characterization of fulvic acids extracted from some organic manures and wastes by potentiometric titration. *Biores. Technol.* 39, 77-83.
- Roletto, E., Chiono, R., and Barberis, E., 1985. Investigation on humic matter from decomposing poplar bark. *Agricultural Wastes* 12, 261-272.
- Roletto, E., and Luda, M.P., 1987. Chemical and infra-red spectroscopic study of humic and fulvic acids from composted poplar bark. *Biological Wastes* 19, 205-214.
- Saviozzi, A., Levi-Minzi, R., and Riffaldi, R., 1988. Maturity evaluation of organic waste. *BioCycle* 29, 54-56.
- Schnitzer, M., 1978. Humic substances: chemistry and reactions. In: M. Schnitzer and S.U. Khan (eds.), *Soil Organic Matter*. Elsevier, Amsterdam, p. 1-64.
- Senesi, N., 1989. Composted materials as organic fertilizers. *Sci. Tot. Environ.* 81 & 82, 521-542.
- Sequi, P., Nobili, M., Leita, L., and Cercignani, G., 1986. A new index of humification. *Agrochimica* 30, 175-179.
- Stevenson, F.J., and Goh, K.M. 1971. Infrared spectra of humic acids and related substances. *Geochim. Cosmochim. Acta* 35, 471-783.
- Stevenson, F.J., 1982. *Humus Chemistry*. John Wiley & Sons Inc., New York.
- Sugahara, K., and Inoko, A., 1981. Composition analysis of humus and characterization of humic acid obtained from city refuse compost. *Soil Sci. Plant Nutr.* 27, 213-224.
- Vaughan, D., and Malcolm, R.E. 1985. Influence of humic substances on growth and physiological processes. In: D. Vaughan and R.E. Malcolm (eds.), *Soil Organic Matter and Biological Activity*. Martinus Nijhoff, Dordrecht, p. 37-75.

Wershaw, R.L., 1985. Application of nuclear magnetic resonance for determining functionality in humic substances. In: G.R. Aiken, D.M. MuKnight, R.L. Wershaw, and P. MacCarthy (eds.), Humic Substances in Soil, Sediments and Water: II. In Search of Structure. John Wiley & Sons, New York, p. 561-584.

Figures legends:

Fig 1: The major relative trends of several humification parameters (expressed as percentage of variation) during 140 days of sludge-straw composting (after Saviozzi et al., 1988).

Fig 2: FT-IR spectra of HA extracted from separated cattle manure at three stages during composting (after Inbar, 1989).

Fig 3: CPMAS ^{13}C -NMR spectra of HA extracted from separated cattle manure at the beginning and after 147 days of composting (after Inbar, 1989).

Table 1. Elemental analysis, H/C and C/N ratios, functional groups and E4/E6 analysis of compost-derived humic acid (HA).

	CSM ¹	MSW ²	SS ³	WC ⁴	GM ¹	"model" HA ⁵	Leonardite ²
C	57.8	52	53	50.58	58.2	56.2	61-55.1
H	5.3	6	5	5.66	5.9	4.7	3.34-3.86
N	3.8	6.3	4.12	3.53	5.8	3.2	1.56-2.43
S	1.1	0	nd	nd	0.66	0.8	0
H/C	1.11	1.4	1.17	1.34	1.21	1	0.66-0.88
C/N	16.6	11.7	13	16.9	14.2	20	52
Total acidity (meq/100 g)	346	498	626	439	413	670	653
COOH (meq/100 g)	221	225	264	247	271	360	279
Phenol OH (meq/100 g)	126	273	362	192	142	390	374
C=O (meq/100 g)	NA	252	288	276	nd	nd	510
E4/E6	7.8	7.1			9.4	4.8	

¹ Inbar, 1989

² Garcia et al., 1992

³ Garcia et al., 1989

⁴ Kakezawa et al., 1992

⁵ Schnitzer, 1978

**Table 2. Carbon distribution in humic acids (HA) extracted from CSM, GM and MSW
(as determined by ^{13}C -NMR).**

		CSM ¹	GM ¹	MSW ²
Aliphatic C	% of total C	50.4	61	67.3
Aromatic C	% of total C	37.1	22	18.8
Carboxylic C	% of total C	10.6	15	10.2
Aromaticity	% of total C	42.4	27	21.8

¹ Inbar, 1989

² Garcia et al., 1992

Table 3: The effects of 50 mg L⁻¹ humic acid (HA) in water or Hoagland's nutrient solution on wheat growth (after Vaughan and Malcom, 1985).

Culture medium	Plant organ	Fresh weight, mg/plant	Stimulation, %
Water	Root	93	0
	Shoot	185	0
Water + HA	Root	146	57.5
	Shoot	252	36.2
Hoagland's	Root	182	96.3
	Shoot	342	84.9
Hoagland's + HA	Root	203	119.0
	Shoot	390	110.8

**CHEMICAL AND BIOLOGICAL
CHARACTERIZATION OF
ORGANIC MATTER DURING
COMPOSTING OF MUNICIPAL
SOLID WASTE**

ABSTRACT

Composting of municipal solid waste (MSW) was studied in an attempt to elaborate transformations of organic matter (OM) during the process and define parameters for the degree of maturity of the product. Composting was performed in 1-m³ plastic boxes and the following parameters were measured in 13 samples during 132 days of composting: temperature, C/N ratio, ash content, humic substance contents and fractions (humic acid, fulvic acid and non humic fraction - HA, FA and NHF respectively). Spectroscopic methods (CPMAS ¹³C-NMR, DRIFT) were used to study the chemical composition of the OM. A bioassay based on growth of cucumber plants was correlated to other parameters. The C/N ratio and ash content showed a typical high rate of change during the first 60 days and reached a plateau thereafter. The HA content increased to a maximum at 112 days, corresponding to the highest plant dry weight and the highest 1650/1560 (cm⁻¹/cm⁻¹) peak ratios calculated from DRIFT spectra. ¹³C-NMR and DRIFT spectra of samples taken from the composting MSW during the process showed that the residual OM contained an increasing levels of aromatic structures. Plant-growth bioassay, HA content and the DRIFT spectra indicated that MSW compost described in this study, stabilized and achieved maturity after about 110 days.

Problems caused to modern society by municipal solid waste (MSW) have become more severe over the last decade due to increasing amounts of waste and decreasing availability of landfill space (Alter, 1991; Finstein, 1992). Recycling has become an attractive option for MSW management. For the organic fraction composting seems to be a desirable treatment having the capacity of reducing the volume and weight by approximately 50% and resulting a stable product which can be beneficial to agriculture (He et al., 1992). The main obstacles facing the production of MSW compost and its utilization stem from the heterogeneous nature of the raw material. The composition of MSW can vary with climate, standard of living, type of garbage collection systems and season.

The composting process consists of a biological treatment in which aerobic thermophilic microorganisms utilize organic matter (OM) as a substrate. The main products of the composting process are fully mineralized materials such as CO_2 , H_2O , NH_4^+ stabilized OM (mostly humic substances) heavily populated with competitive microbial biomass and ash. Compost has the potential of improving soil structure, increasing cation exchange capacity, suppressing soil borne plant pathogens and enhancing plant growth (Saviozzi et al., 1988; Chen et al., 1989; Hoitink et al., 1993). The chemical and microbiological properties of MSW compost have been widely studied and the following parameters have been proposed as maturity and stability indexes: C/N in solid and in water phase (Chanyasak et al., 1982; Jimenez and Garcia, 1992), cation exchange capacity (Harada et al., 1981), humification indexes (Jimenez and Garcia, 1992) and oxygen and CO_2 respirometry (Iannotti et al., 1994). Chemical and spectroscopic analyses of OM transformations during composting in relation to compost maturity of various raw materials have been recently reviewed (Chen and Inbar, 1993).

One of the main obstacles to successful utilization of MSW compost in agriculture is the lack of reliable quality criteria and understanding the OM transformations throughout the process. Proper evaluation of compost maturity is essential for the

establishment of such criteria. Therefore, the objective of this research was to study the transformation of OM during composting of MSW and to correlate chemical analyses and ^{13}C -NMR and DRIFT spectra, with plant performance bioassays.

MATERIALS AND METHODS

Composting of MSW

MSW was collected over a 24 h period from the town of Afula in Israel, and mechanically mixed in a MSW classifying facility. After sieving and eliminating the >5-cm fraction, the raw material was divided into three perforated plastic boxes (about 1-m³ each) and composted for 132 days. The compost was turned, mixed, and sampled after removal of glass, plastics and metals by hand-picking at 0, 6, 13, 19, 27, 34, 41, 48, 62, 76, 90, 112 and 132 days. Samples (10 L each) were stored at 4°C. Smaller subsamples were dried at 65°C, then ground and sieved. Particles smaller than 1-mm in diameter were used for analyses. Temperature was measured at 4 h intervals by thermocouples connected to a data logger until reaching ambient temperature (~ 15 to 20°C), then, larger intervals (4 d) were applied between temperature readings.

Chemical analyses

Total organic carbon was measured by loss of weight on ignition at 550°C for 8 h (C organic calculated as 58% of the OM). Organic nitrogen was measured by a Tecator Automatic Kjeldahl System (Kjeldahl System 1030, Tecator, Hsiganas, Sweden).

For studies on humic substances (HS) 20 g of compost were extracted with 200 ml 0.1 N NaOH under N₂. The supernatant solution containing the soluble HS was separated by centrifugation at 15,000×g and the residue was resuspended in 0.1 N NaOH. This procedure was repeated eight times. The combined solutions were acidified with 6 M HCl to pH~1 and allowed to stand at room temperature for 24 h. The supernatant solution

containing the extracted fulvic fraction (FF) was separated from the humic acid (HA) by centrifugation (2,500×g for 30 min.). The fulvic fraction was separated into fulvic acid (FA) and non-humic fraction (NHF) by adsorption of the FA onto Amberlite XAD-8 resin (mesh size 20-60, Sigma, St. Louis, USA). The Amberlite XAD-8 resin was cleaned before use according to the method described by Thurman and Malcolm (1981). The FA was passed through the column at a flow rate of about 15 pore volumes per hour. The column was then washed by 0.65 pore volumes of distilled water. At this stage the FA adsorbed to the resin while the NHF was eluted. Carbon concentrations of the FF and NHF were determined by a total carbon monitor (TCM 480, Carlo Erba Instruments, Millano, Italy). FA content was calculated as the FF minus NHF. The following humification parameters were calculated: (i) humification ratio (HR)=HA/FF; (ii) humification index (HI)=HA/FA; (iii) HA percentage (HP)=100×HA/HS.

¹³C Nuclear Magnetic Resonance Spectroscopy

Solid state C-13 NMR spectra with CPMAS (Cross Polarization Magic Angle Spinning) were obtained using a Chemagnetics M-100 NMR-spectrometer, located at Pennsylvania State University. The spectrometer operates at a ¹H frequency of 100 MHz and a ¹³C frequency of 25.2 MHz. Pertinent experimental parameters were the following: contact time of 1 ms; recycle delay time of 1 s; sweep width of 14 kHz (562.5 ppm) and line broadening of 30 Hz. Dried sieved compost samples were placed into a rotor and spun at a frequency of 3.5 kHz at the magic angle (54.7° to the magnetic field).

The ¹³C-NMR spectra were divided into five regions (Hatcher et al., 1983). Region I (0-50 ppm) is assigned to paraffinic carbon or carbons that are bonded only to other carbons. Region II (50-112 ppm) is assigned to C-O, C-N bonds as in carbohydrates, alcohols, esters and amines. Region III (112-163 ppm) is assigned to aromatic and phenolic carbons. Region IV (163-190 ppm) represents carboxyl, ester and amide carbons groups. Region V (190-215 ppm) represents the carbonyl carbons. The

region 0-112 ppm was calculated as aliphatic carbon, 60-112 ppm as polysaccharide carbon and 112-163 ppm as aromatic carbon. Total aromaticity was calculated by expressing aromatic C (112-163 ppm) as percentage of the aliphatic C (0-112 ppm) plus aromatic C (Hatcher et al., 1981).

Diffuse Reflectance Infrared Fourier-Transform (DRIFT) Spectroscopy

The DRIFT spectra of MSW compost at twelve stages of the process were obtained for a wavenumber range of 4000 to 400 cm^{-1} in a Nicolet 550 Magna-IRTM spectrometer (Nicolet Instruments Corporation, Madison, WI, USA). Bulk samples were prepared for analysis according to the methods of Niemeyer et al. (1992), using 15 mg of dried bulk compost mixed with KBr for a total sample weight of 310 mg. The samples were transferred to a macro-sample holder and smoothed with a glass slide. To obtain DRIFT spectra, 400 scans were collected. Spectra of the residual ash obtained after ignition at 550°C for 8 h and loss of the OM were recorded using the same methodology. To compare one spectrum to another a linear baseline correction was applied using 4000 cm^{-1} , 2000 cm^{-1} and 860 cm^{-1} as zero absorbance points. The major peaks data (intensity and wavenumber) were found by using the OMNIC software (Nicolet Instruments Corporation). The spectra of the ash free OM was calculated by subtracting the spectrum of the ash from that of the bulk materials, using the OMNIC software.

Plant-growth bioassay

Ten cucumber seeds (*Cucumis Sativus* L., cv. dlila, Zeraim-Gedera, Israel) were placed in 0.5-L pot containing 50% MSW compost (various stages of composting) and 50% of perlite (Habonim, Israel, no. 4). Each treatment consisted of five replicates. The plants were irrigated every two days with equal amounts of tap water without fertilizer. After one week the seedlings were thinned, leaving the best three plants per pot. These

plants were grown for one month. Wet and dry weights of the plants in each pot were determined.

Statistical analysis

Ash content, C/N ratio and temperature results are presented as the average value of three replicates of the three composting boxes, while the data for each box is the average of three replicates with standard errors. HA, FA and NHF content and spectroscopic analyses were determined on a mixed sample from the three composting boxes. The plant bioassay statistical analysis was performed using the Duncan test at $p=0.05$.

RESULTS

Composting of MSW

As reported earlier (Harada et al., 1981; Jimenez and Garcia, 1992; Inbar et al., 1993) three typical phases of composting were observed by temperature measurements during the process (Fig. 1): (i) a mesophilic phase, during the first 2 d in which temperature rose to 45°C; (ii) a thermophilic phase lasting 4 weeks in which the temperature increased to a maximum of 72°C; further temperature increases were avoided by turning and mixing the compost once a week; (iii) after 60 d of composting the compost temperature equaled that of the ambient. It is interesting to note that the highest temperature during thermophilic phase was over 70°C while the mean and minimum ambient temperatures were only 10-15°C and 0°C respectively.

The change in the C/N ratio and ash content reflects OM decomposition and stabilization during composting (Fig. 2). The C/N ratio decreased rapidly from an initial value of 28 in the raw material to 18 after only 20 days. The ratio continued to decrease, albeit less sharply, to 12.2 after 60 days. From this point on the C/N ratio stabilized at a

value of about 12 (11.8-12.8) for the remainder of process. The initial ash content of the compost was relatively high (~45%) since the raw matter had been collected from a residential area without on-site separation. The change in ash content followed a trend similar to that of the C/N ratio, exhibiting three phases: (i) lasting over the first 20 d, when most of the OM decomposed; (ii) days 20-80; (iii) days 80 to the end of the experiment, the period at which the curing stage began and the rate of decomposition of OM was extremely low.

Humic Substance Content

Humic substances (HS) comprise the most important fraction of OM due to their effects on soil ecology, structure, fertility and plant growth. NaOH extracted HS from composts can be separated into humic acid (HA), fulvic acid (FA) and a non-humic fraction (NHF) (Chen and Inbar, 1993). The levels of HA, FA and NHF in MSW compost at various stages of the process represent the humification process (Fig. 3). Total levels of HS did not change during the composting process, maintaining values of 14.2% to 18% of the OM. The FA level remained steady at 7.5% of OM for the first 20 days, then decreased to 4.5% in the mature compost. The HA level increased during the composting process, reaching 14% of the OM after 112 days maintaining this value till the end of the process. The NHF decreased rapidly from 8.7% to 4.4% at 60 days of composting probably since it consisted of polysaccharides and fats which are readily decomposed. All three humification parameters (HR, HI and HP) increased during the composting as follows (Table 1): (i) HR increased from 0.42 in the raw MSW to 1 after 30 days, stabilized at 1-1.18 until day 90, then sharply increased to 1.35 in the mature compost; (ii) HI increased rapidly from 0.91 in the raw material to 2 within 30 days, then gradually increased to a final value of 3 at the end of the process; (iii) HP values exhibited a similar trend. The increasing trend of these parameters indicated that during composting HA became the main fraction of HS.

Spectroscopic Methods

DRIFT Spectra

The use of DRIFT spectra represents a new approach in studies of the structure of heterogeneous materials such as plant tissue and bulk compost samples. The development of an attachment mounted onto FTIR spectrophotometers has allowed the determination of DRIFT spectra. Sample preparation for DRIFT spectroscopy is much simpler than for FTIR spectroscopy, interferences due to water adsorption are reduced, and resolution is improved. The spectra obtained using DRIFT had a higher degree of resolution as compared with FTIR. Moreover, the bands indicative of aliphatic C-H, carboxyl and carboxylate functional groups, aromatic C=C, and the C-O stretch of polysaccharides were prominent and very well resolved. The DRIFT spectra can also be used to fingerprint OM acquired from various sources. Relative concentrations of functional groups were found to be fairly constant regardless of sample concentration. The DRIFT spectra peak intensity ratio can be used to determine the humification process (Niemeyer et al., 1992).

All five bulk MSW compost spectra samples from different stages (raw, 34, 76, 90 and 132d) of composting exhibited similar peaks (Fig. 4). The main absorbance bands were: a broad band at 3300-3400 cm^{-1} (H-bonds, OH groups), two distinct peaks at 2930 and 2850 cm^{-1} (C-H asymmetric, C-H stretch of -CH), a peak at 2520 cm^{-1} , a sharp peak at 1800 cm^{-1} , a slight shoulder around 1740 cm^{-1} , a wide peak at 1650 cm^{-1} (C=C in aromatic structure), a shoulder at 1560 cm^{-1} (amide II bonds), a strong peak at 1450 cm^{-1} (C-H deformation of CH_2 or CH_3 groups), a peak in the 1100-1000 cm^{-1} range (C-O stretch of polysaccharide, Si-O stretch) and a sharp peak at 870 cm^{-1} (aromatic CH out of plane bending) (Baes and Bloom, 1989; Inbar et al., 1989).

The distinct change in the spectra resulting from the composting process was a reduction of the 1560 cm^{-1} peak with time probably due to the relatively rapid

biodegradation of the amino chain. The aromatic region and the 1800 cm^{-1} peak became sharper during composting. Peaks in the aliphatic region at 2930 and 2850 cm^{-1} decreased, while the 1450 cm^{-1} peak increased. The spectra of MSW compost (wavenumbers $4000\text{--}400\text{ cm}^{-1}$) were different from spectra of composts from other sources that were reported earlier (Niemeyer et al., 1992; Inbar et al., 1989). The main differences between the spectra of MSW compost and the other composts were in the following regions: (i) a sharp peak and a shoulder at 2520 cm^{-1} and 2580 cm^{-1} respectively which does not appear in spectra reported for other composts; (ii) a narrow sharp peak at 1800 cm^{-1} that appears as a small shoulder at 1710 to 1720 cm^{-1} in spectra of other composts; (iii) the peak at 1450 cm^{-1} had become the main peak in the spectra of MSW compost instead of the aromatic or the polysaccharide peaks in other composts; (iv) an intense and sharp peak at 870 cm^{-1} that does not appear in spectra of other compost. According to Pouchert (1981) the peak at 2520 cm^{-1} can be attributed to S-H stretch of aromatic or nonaromatic mercaptans and sulfides, the absorbance at 1800 cm^{-1} can be attributed to the C=O stretch of the conjugated carbonyl group of aromatic esters and lactones. Both these organic compounds should not appear in composts for the following reasons: (i) S-H bonds can be related to proteins which are rapidly biodegraded; (ii) sulfides form only under anaerobic conditions; (iii) esters and lactones are biodegraded relatively rapidly by many microorganisms.

Spectra of residual ash samples were recorded in order to verify whether these peaks are related to OM or to minerals forms in the compost. The spectra of the residual ash were in similar to the spectrum of calcite (CaCO_3) exhibiting major absorption bands at 2983 cm^{-1} , 2874 cm^{-1} , 2580 cm^{-1} , 2512 cm^{-1} , 1800 cm^{-1} , 1420 cm^{-1} and 874 cm^{-1} . The ash also exhibited peaks in the $1100\text{--}950\text{ cm}^{-1}$ region and sharp peaks at 680 cm^{-1} and 620 cm^{-1} which can be attributed to phosphate minerals or to Si-O components (Kodama, 1985). Thus, it is concluded that the mineral matter is responsible for the unusual peaks observed in the spectra of the MSW compost. To correct and reduce the

contribution of the mineral component, subtraction of the residual ash spectrum from the bulk spectrum was performed. The spectra of the ash free MSW compost at two stages of composting are presented in Fig. 5. The spectrum of the mature compost is generally similar to spectra of other composts (Niemeyer et al., 1992) with specific differences at: (i) the polysaccharide region peak ($1000\text{--}1100\text{ cm}^{-1}$) which was smaller in MSW than in other composts; and (ii) the OM in MSW compost was more aliphatic (and less aromatic) than pine bark compost, hardwood bark compost and compost from separated cattle manure (CSM) reported by Niemeyer et al. (1992). Various differences between the raw materials and mature compost as revealed by comparing the subtracted spectra can be pointed out: (i) reduction of the 1560 cm^{-1} peak; (ii) reduction and transition to a small shoulder of the 1738 cm^{-1} peak (CO_2H groups) in the mature compost; (iii) a sharp decrease in the aliphatic region (2930 cm^{-1} and 2850 cm^{-1}); and (iv) an increase in the aromatic region (1650 cm^{-1}).

Following Inbar et al. (1989) we monitored the changes in the IR spectra by calculation of the ratio between the intensity of major peaks. The peaks at 2930 cm^{-1} , 2850 cm^{-1} , 1650 cm^{-1} , 1560 cm^{-1} and 1050 cm^{-1} were chosen for these calculations. The ratio $1650/2930$ (aromatic C / aliphatic C) increased from 0.88 to 1.10, the ratio $1650/2850$ (aromatic C / aliphatic C) increased from 0.79 to 1.54, the $1650/1050$ ratio (aromatic C / polysaccharide) increased from 2.39 to 2.80, and the $1650/1560$ ratio (aromatic C / amide II bond) increased from 0.94 to 1.52. These ratios were calculated for spectra of ash free OM. The changes reflect a decrease in levels of polysaccharides, aliphatic and amide components, and an increase in levels of aromatic structures in mature compost. We believe that the increase in aromatic structures is relative and it resulted from decomposition of aliphatic moieties. The linear correlation between the aromatic to aliphatic peak ratio ($1650/2930$) and the C/N ratio (Fig. 6) exhibited $R^2=0.936$ calculated for the equation $Y = 1.263 - 0.014X$. This correlation indicates that DRIFT is a useful and reliable tool in the analysis of OM decomposition during composting.

CPMAS ^{13}C -NMR

During the process of compost production, organic materials that are chemically complex and difficult to fractionate are formed. Extraction procedures that are commonly applied to soils and composts remove only a part of the C in the system. To profile the organic components of composts during the course of their formation, a method that can be applied to the intact sample is preferable. The only physical method holding some promise for the reliable characterization of such organic materials appears to be nuclear magnetic resonance (NMR) (Wershaw, 1985). The CPMAS ^{13}C -NMR technique has been reported to provide useful information on transformation of OM during composting (Chen and Inbar, 1993). The major peaks exhibited in the CPMAS ^{13}C -NMR spectra of samples representing five stages in the composting process (0, 34, 76, 90 and 132 d) (Fig. 7) are at: 30 ppm (methylene C), 55 ppm (methoxyl C), 72 ppm (carbohydrate or aliphatic alcohols), 105 ppm (anomeric carbons of polysaccharides), 130 ppm (C-substituted aromatic C), 150 ppm (O- substituted aromatic C) and at 175 ppm (carboxyl carbon) (Kogel-Knabner et al., 1991).

The distinct changes in the spectra resulting from decomposition of OM were: reduction of the 30 ppm peak; appearance of the 55 ppm peak as a shoulder in the mature compost and an increase of the aromatic region peak (130 and 150 ppm peaks). The relative changes in the level of carbon containing groups are presented in Table 2. The following changes should be noted: (i) total aliphatic carbon (0-112 ppm) decreased by 14.4% from 74.8% in the raw material to 64.0% in the mature compost; polysaccharides (60-112 ppm) decreased by 17.0% from 42.3% to 35.1% in the mature compost; and alkyl groups (0-50 ppm) decreased by 11.0% during the composting process; (ii) aromatic components, including aromatic and phenolic carbons, increased by 54%, from 15.9% in the raw material up to 24.5% of the total carbon in the mature compost; total aromaticity increased by 58.2%, from 17.5% in the raw material to 27.7% in the mature compost; (iii) carboxyl carbon increased by only 9.5%; (iv) the C=O carbonyl carbon increased by

73.7%, from 1.9% in the raw material to 3.3 % in the mature compost. It should be noted however, that this carbonyl C represents a small fraction of total C and the peaks were poorly resolved.

Plant-growth bioassay

To establish that a compost is mature, devoid of phytotoxic compounds and ready for use for plant production, several chemical parameters have been suggested as maturity indices. However, so far only a plant bioassay can provide reliable results (Hoitink and Kuter, 1986; Inbar et al., 1993). The dry weight of cucumber plants grown in composts of various ages are presented in Fig. 8. Plants grown in media containing fresh (14 d) compost exhibited inhibited growth compared to plants grown on older compost. The dry weight of plants grown on samples taken after 28 to 84 d of composting were statistically similar. However, the dry weight of plants grown in 112 and 132 d old composts were significantly higher than all the others. These results show that as OM decomposed the compost became a better substrate for plant growth. Only mature compost supported better plant growth.

The highest plant dry weight was achieved after 112 days of composting. This corresponds to the sample in which maximum amount of HA were present, and to the time at which the 1650/1560 DRIFT peak ratio has stabilized at a maximum height. The linear correlation between the 1650/1560 ratio and plant dry weight exhibited a $R^2=0.81$ calculated for the equation $Y = 0.81 + 0.54X$. The linear correlation between HA content in OM and plant dry weight exhibited a $R^2=0.70$ calculated for the equation $Y = 3.59 + 7.78X$.

DISCUSSION

In this article we studied transformations of OM during composting of MSW, aiming to define compost maturity and to correlate the chemical and spectroscopic analyses to plant growth. Although the C/N ratio is the most common parameter used to define decomposition levels of OM it appears to be a non reliable indicator of compost maturity (Harada et al., 1981). This ratio changed dramatically during the second phase of composting performed in our study (from 28 to 12 within 60 d), it did not change significantly during the curing phase. Only minor changes in the C/N ratio occurred during the curing stage whereas instability of the OM was observed in spectra obtained by DRIFT and NMR. A C/N ratio of 10-12 is usually considered to be an indicator of stable and decomposed OM (Jimenez and Garcia, 1992). The fact that no changes took place during the curing phase challenges the use this ratio as a ultimate parameter. The level of HS in the compost did not change during composting of MSW. This observation agrees with that of Sugahara and Inoko (1981), Das (1988), and Jimenez and Garcia (1992), but contradicts the report of Inbar et al. (1989) who showed that HS levels increased significantly with time of composting in CSM. This suggests that the type of raw material is of major importance to the humification process during composting. In this study the HA content increased during composting, reaching a stable value of 13.5% of the OM after 112 days, whereas the FF level decreased. The increasing level of HA represents the degree of humification and maturity of the compost. HI and HR are commonly used to analyze the humic fraction (Jimenez and Garcia, 1992; Inbar et al., 1989). We found that the HI increased to a ratio of 3, and the HR increased to 1.35. The trend of change of these values is similar in various wastes although specific values differ in other wastes. In general as well as in our present study, fresh composts contain low levels of HA and higher levels of FA (Harada et al., 1981; Saviozzi et al., 1988; Inbar et al., 1989; Gonzalez et al., 1993; Ciavatta et al., 1993a; Ciavatta et al., 1993b). The HP increased to a level

higher than 73% in the mature compost indicating that the fraction defined as FA, extracted from heterogeneous materials such as MSW, contains relatively high levels of biodegradable OM that was decomposed mainly during days 20 to 40 (Fig. 3).

DRIFT spectra provide useful information about the transformation of OM. The main absorbance bands indicate changes in OM structure are at 2930 cm^{-1} and 2850 cm^{-1} (aliphatic groups), 1650 cm^{-1} (aromatic structure), and $1100\text{-}1000\text{ cm}^{-1}$ (polysaccharide). The relative height of the aromatic peak at 1650 cm^{-1} increased with composting time and stabilized at a later stage of the process similar to the level of HA. The increase in the $1650/2930$, $1650/2850$ and $1650/1560$ ratios also indicates that the easily degradable OM constituents such as aliphatic chains, polysaccharides and alcohols are chemically or biologically oxidized, and therefore the mature compost contained more aromatic structures of higher stability. The DRIFT spectra of the bulk MSW compost provide information not only on the OM bonds but also on the minerals in the raw material. The relative level of these materials increased (1800 cm^{-1} and 1450 cm^{-1}) during composting in parallel to the increase in ash. Based on our measurements with MSW compost one should bear in mind that when analyzing high ash containing wastes, spectra of the ash should be subtracted from those of the bulk material.

CPMAS ^{13}C -NMR spectroscopy is a powerful technique for the analysis of chemical transformations of the OM during composting. ^{13}C -NMR spectra determined in this research also indicate that the OM structure become more aromatic while carbohydrates, polysaccharides, alcohols, lipids and protein decomposed, along with the relative increase of all other components (HA). These findings are in general agreement with the data obtained from the DRIFT spectra except for the carbonyl and carboxyl regions that, according to the NMR data, increased during composting by 73.0% and 9.5%, respectively. In the DRIFT spectra the carbonyl and carboxyl carbon ($1740\text{-}1720\text{ cm}^{-1}$) peak intensities were smaller or subjected to a shift. A comparison between ^{13}C NMR spectra of CSM (Inbar et al., 1989; Inbar et al., 1991) and those of the MSW

compost reveals consistent trends observed in the various carbon-containing functional groups. The only exception was the carboxyl region that increased by 90% during composting of CSM whereas in MSW compost, the carboxyl region increased by 9.5% only. The general similarity in the transformation of OM during composting of these two different raw materials as analyzed by DRIFT and NMR suggest that the composting process transforms heterogeneous raw OM to a compositionally uniform product at the end of the process.

The aim of the plant-growth bioassay (Fig. 8) was to obtain biological criteria for compost maturity. We hypothesized that plant growth may serve as an integrative single parameter for compost maturity. The results of the plant-growth bioassay were similar to those of Zucconi et al. (1981), Chanyasak et al. (1983), and Inbar et al. (1993). In all of these studies plant growth in immature compost was inhibited and it generally improved in the mature compost. These authors suggested that in immature composts low molecular weight organic acids induce phytotoxicity in addition to competition for oxygen and nutrients, due to high rates of OM decomposition. The correlation between the HA level, the 1650/1560 DRIFT peak ratio and plant biomass suggested that these parameters could indicate compost maturity.

CONCLUSIONS

Compost maturity is very difficult to define by only one parameter usually several parameters are needed to be crossbred. This article demonstrates that ¹³C-NMR and DRIFT spectroscopy, together with data on HS provide useful information on the OM transformations occurring during the composting process. Plant-growth bioassay, level of HA and the 1650/1560 DRIFT peak ratio exhibited the same trend during composting and can therefore be used as maturity indices. All these parameters exhibited three phases: (i) rapid decomposition during the first 30 d; (ii) the stabilization stage till day 90; and (iii)

the maturation phase from day 100 on. The MSW compost, described in this study, was mature and ready for use as an agricultural substrate after about 110 days of composting. The understanding of compost maturity has to be improved by investigations of additional parameters for maturity such as the biochemistry of humic components and reactions of the OM occurring in the interface between solid and aqueous phases (Chanyasak et al., 1983; Chen and Inbar, 1993).

REFERENCES

- Alter, H. 1991. The future course of solid waste management in the US. *Waste Management & Research* 3:3-20.
- Baes, A.U., and P.R. Bloom. 1989. Diffuse reflectance and transmission fourier transform infrared (DRIFT) spectroscopy of humic and fulvic acids. *Soil Sci. Soc. Am. J.* 53:695-700.
- Chanyasak, V., A. Katayama, M. Hirai, S. Mori, and H. Kubota. 1983. Effects of compost maturity on growth of komatsuna (*Brasica rapa* var. *peridis*) in neubauer pot. II. Growth inhibitory factors and assessment of degree of maturity by org-C/org-N ratio of water extract. *Soil Sci. Plant Nutr.* 29:251-259.
- Chanyasak, V., M. Hirai, and H. Kubota. 1982. Changes of chemical components and nitrogen transformation in water extracts during composting of garbage. *J. Ferment. Tech.* 60:439- 446.
- Chen, Y., and Y. Inbar. 1993. Chemical and spectroscopic analysis of organic matter transformations during composting in relation to compost maturity. p. 551-600. *In* H.A.J. Hoitink and H.M. Keener (ed.) *Science and engineering of composting: design, environmental, microbiological and utilization aspects*. Renaissance Publications, Worthington, OH.
- Chen, Y., Y. Inbar, Y. Hadar, and R.L. Malcolm. 1989. Chemical properties and solid-state CPMAS ^{13}C -NMR of composted organic matter. *Sci. Total Environ.* 81/82:201-208.
- Ciavatta, C., M. Govi, L. Pasotti, and P. Sequi. 1993a. Changes in organic matter during stabilization of compost from municipal solid waste. *Bioresource Tech.* 43:141-145.
- Ciavatta, C., M. Govi, and P. Sequi. 1993b. Characterization of organic matter in compost with municipal solid wastes: an italian approach. *Compost Sci. & Utilization* 1:75-81.

- Das, A. 1988. City garbage compost as a source of humus. *Biol. Wastes* 26:65-69.
- Finstein, M.S. 1992. Composting in the context of municipal solid waste management. p. 355-374. *In* R. Mitcell (ed.) *Environmental microbiology*. Wiley-Liss, New-York.
- Gonzalez, S.J., M. Carballas, M.C. Villar, M.C. Beloso, A. Cabaneiro, and T. Carballas. 1993. Carbon-and nitrogen-containing compounds in composted urban refuses. *Bioresource Tech.* 45:115-121.
- Hatcher, P.G., M. Schnitzer, L.W. Dennis, and G.E. Maciel. 1983. Solid state ^{13}C -NMR of sedimentary humic substances: new revelation on their chemical composition. p. 37-81. *In* R.F. Christman and E.T. Gjessing (ed.) *Aquatic and terrestrial humic materials*. Ann Arbor Science Publishers, MI.
- Hatcher, P.G., M. Schnitzer, L.W. Dennis, and G.E. Maciel. 1981. Aromaticity of humic substances in soil. *Soil Sci. Soc. Am. J.* 45:1089-1094.
- Harada, Y., A. Inoko, M. Tadaki, and T. Izadaki. 1981. Maturing process of city refuse compost during piling. *Soil Sci. Plant Nutr.* 27:357-364.
- He, X.T., S. J. Traina, and T. Logan. 1992. Chemical properties of municipal solid waste composts. *J. Environ. Qual.* 21:318-329.
- Hoitink, H.A.J., M.J. Boehn, and Y. Hadar. 1993. Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. p. 601-621. *In* H.A.J. Hoitink and H.M. Keener (ed.) *Science and engineering of composting: design, environmental, microbiological and utilization aspects*. Renaissance Publications, Worthington, OH.
- Hoitink, H.A.J., and G.A. Kuter. 1986. Effects of composts in growth on soilborne pathogens. p. 289-306. *In* Y. Chen and Y. Avnimelech (ed.) *The role of organic matter in modern agriculture*. Martinus Nijhoff Publishers, Netherlands.
- Iannotti, D.A., M.E. Grebus, B.L. Toth, L.V. Madden and H.A.J. Hoitink. 1994. Oxygen respirometry to assess stability and maturity of composted municipal solid waste. *J. Environ. Qual.* 23:1177-1183.

- Inbar, Y., Y. Hadar, and Y. Chen. 1993. Recycling of cattle manure: the composting process and characterization of maturity. *J. Environ. Qual.* 22:857-863.
- Inbar, Y., Y. Chen, and Y. Hadar. 1991. Carbon-13 CPMAS NMR and FTIR spectroscopic analysis of organic matter transformations during composting of solid wastes from wineries. *Soil Sci.* 152:272-282.
- Inbar, Y., Y. Chen, and Y. Hadar. 1989. Solid state carbon-13 nuclear magnetic resonance infrared spectroscopy of composted organic matter. *Soil Sci. Am. J.* 53:1695-1701.
- Jimenez, E.I., and V.P. Garcia. 1992. Determination of maturity indices for city refuse composts. *Agric. Ecosystems Environ.* 38:331-343.
- Kodama, H. (ed.). 1985. Infrared spectra of minerals, reference guide to identification and characterization of minerals for the study of soils. Ministry of Supply and Services Canada.
- Kogel-Knabner, I., P.G. Hatcher, and W. Zech. 1991. Chemical structural studies of forest soil humic acids: aromatic carbon fraction. *Soil Sci. Am. J.* 55:241-247.
- Niemeyer, J., Y. Chen, and J.M. Bollag. 1992. Characterization of humic-acids, compost, and peat by diffuse reflectance fourier-transform infrared spectroscopy. *Soil Sci.* 56:135- 140.
- Pouchert, C.J. (ed.). 1981. The aldrich library of infrared spectra, edition III. Aldrich Chemical Company Inc., Milwaukee, Wisconsin.
- Saviozzi, A., R. Levi-Minzi, and R. Riffaldi. 1988. Maturity evaluation of organic waste. *BioCycle* 29:54-56.
- Sugahara, K., and A. Inoko. 1981. Composition analysis of humus and characterization of humic acid obtained from city refuse compost. *Soil Sci. Plant Nutr.* 27:213-224.
- Thurman, E.M., and R. Malcolm. 1981. Preparative isolation of aquatic humic substances. *Environ. Sci. Tech.* 15:463-466.

- Wershaw, R.L. 1985. Application of nuclear magnetic resonance for determining functionality in humic substances. p. 561-584. *In* G.R. Aiken et al (ed.) Humic substances in soil, sediment and water. John Wiley & Sons, Inc., New York.
- Zucconi, F., M. Forte, A. Monaco, and M. De Bertoldi. 1981. Biological evaluation of compost maturity. *BioCycle* 22:27-29.

Table 1. Humification index ($HI=HA/FA$), humification ratio ($HR=HA/FF$) and percentage of humic acid ($HP=100 \times HA/HS$) during composting of MSW.

Composting Time (days)	HI	HR	HP
0	0.91	0.42	47.57
13	1.25	0.67	55.51
19	1.36	0.76	57.66
27	2.01	1.02	66.80
34	2.07	1.01	67.47
41	2.14	1.13	68.14
48	2.18	1.17	68.54
62	1.95	1.08	66.07
76	2.59	1.18	72.12
90	2.77	1.12	73.50
112	2.82	1.35	73.84
132	2.99	1.32	74.96

Table 2. Distribution of carbon containing groups during composting of MSW as determined by CPMAS ^{13}C -NMR (percent of total carbon).

Composting Time (days)		0	13	34	76	132	Change from raw material
Chemical shift (ppm)	Carbon containing group	Percent of total carbon					
0-50	Alkyl	26.9	25.5	24.7	23.6	23.6	-12.3%
50-112	O-alkyl	47.9	52.1	46.5	42.4	40.4	-16.0%
112-145	Aromatic	11.6	10.5	13.6	19.0	16.9	+45.6
145-163	Phenolic	4.3	3.6	4.9	6.6	7.6	+76.7
163-190	Carboxyl	7.4	6.6	8.2	8.0	8.1	+9.5%
190-215	Carbonyl	1.9	1.6	2.1	3.4	3.3	+73.0%

Figures legends

Fig. 1. Ambient and compost temperature vs. composting time of MSW.

Fig 2. C/N ratio and ash content during composting of MSW.

Fig. 3. Relative content of the humic acid (HA), fulvic acid (FA) and non humic fraction (NHF) as percent of organic matter vs. composting time of MSW.

Fig. 4. DRIFT spectra of bulk MSW compost at five stages of composting (raw, 34, 76, 90, 132 d).

Fig. 5. DRIFT spectra of the ash free OM in MSW compost at two stages of composting (raw and 132 d). Ash spectra were subtracted from those of the bulk materials.

Fig. 6. DRIFT peak intensity ratio ($1650/2930\text{ cm}^{-1}/\text{cm}^{-1}$) vs. C/N during composting of MSW. Peak ratio was calculated from twelve subtracted spectra at various stages of composting, using the same method as in Fig. 5 (dotted lines represent the 95 confidence interval).

Fig. 7. CPMAS ^{13}C -NMR spectra of MSW compost samples at five stages during composting of MSW (raw, 13, 34, 76, 132 d).

Fig. 8. Dry weight of cucumber plants grown in MSW compost samples at eight stages during composting time. Data followed by the same letter are not significantly different ($p = 0.05$).

Suppression of Soil-Borne Pathogens
by Composted
Municipal Solid Waste

Suppression of Soil-Borne Pathogens by Composted Municipal Solid Waste

Abstract

This study was aimed at determining the level of suppressiveness of composted municipal solid waste (CMSW) against soil-borne plant pathogens and investigating the microbial mechanisms involved in this suppression. The suppressiveness of CMSW to *Rhizoctonia solani*, *Fusarium oxysporum* and *Pythium aphanidermatum* was studied in greenhouse experiments in 500 mL polypropylene pots. The pots contained sandy soil or peat amended with different amount of CMSW. *R. solani* and *P. aphanidermatum* were reduced by about 80 and 33%, respectively. *F. oxysporum*, which causes wilt-disease of cotton, was reduced by 95%. Compost age influenced its suppression ability. Moreover, compost which had been composting for 1.5 months of composting did not suppress *R. solani*, whereas 2-week-old compost suppressed *F.oxysporum*, as did mature compost. Reducing the CMSW's microbial population by autoclaving, gamma irradiation or heat treatment negated *R. solani* suppression, indicating that the mechanism involved in this suppression is biological. This conclusion was supported by the fact that the addition of 5% (v/v) CMSW to heat-treated CMSW caused the recolonization of antagonistic microorganisms and restored the compost's suppressive ability. Heat treatment of CMSW did not eliminate its ability to suppress *F. oxysporum*, suggesting the involvement of a thermophilic population in this particular case. In addition, physical and chemical factors may be involved in this suppression. The mechanism governing in the suppression of *R. solani* is suggested to involve lysis of fungal hyphae and bacterial activity. Some bacteria isolated from the CMSW showed chitin-degradation ability and the capacity to inhibit *R. solani* growth. Two of the isolated bacteria exhibited suppression of *R. solani* in cotton plants in greenhouse experiments. One of these was identified as *Enterobacter agglomerans*. Inoculation of heated compost with *Trichoderma harzianum* suppressed *R.*

solani significantly. However, the addition of *T. harzianum* did not improve the suppressive ability of the mature compost despite the fact that this compost was successfully colonized.

Introduction

Compost is the stable product of aerobic microbial decomposition of heterogenic organic matter (12). Various organic wastes such as grape marc, cattle manure, hardwood bark, yard waste, municipal solid waste (MSW) and sewage sludge can be used as raw materials for composting (2, 3, 11, 13). Composting of the organic fraction of MSW can reduce up to 50% of the waste volume reaching at the landfills and produces a stable product that can be used in agriculture (4, 5, 14).

The extent of organic matter decomposition and compost maturity affect the latter's ability to suppress soil-borne plant pathogens (12, 14). In recent years, compost amendments in agriculture have been investigated as part of an integrated system of biological control due to their ability to suppress soil-borne plant pathogens such as *Pythium aphanidermatum*, *Rhizoctonia solani* and *Fusarium oxysporum* (6, 7, 8, 10, 15, 22, 23, 24).

Suppressive compost can be defined as providing an environment in which disease development is reduced, even when the pathogen is introduced in the presence of a susceptible plant (1). Plant-disease suppression is a direct result of the activity of antagonistic microorganisms, which naturally recolonize compost during the cooling phase of the composting process. Thus, compost sterilization most of the time negates suppression. Although the mechanism of suppression is biological in all cases, different specific mechanisms have been observed in relation to various plant diseases (12). The mechanism involved in suppression of *P. aphanidermatum*, which causes damping-off in cucumber plants, is general (17). Competition for nutrients was found to be responsible for this type of suppression and to be affected by carbon-source concentration and microbial activity (18).

Media amended with CGM (composted grape marc) or CSM (composted separated manure) and other composts were also effective in suppressing diseases caused by *Sclerotium rolfsii* and *R. solani* (9). The mechanism involved in this type of suppression is a specific one in which mycoparasitism is observed. *S. rolfsii* sclerotia lost their viability within 24 h of incubation in the compost. Scanning electron microscopy revealed the sclerotia to be heavily colonized by fungi, mainly *Penicillium* spp. (9, 17). In other composts, specific suppression has been found to be related to the a large *Trichoderma* population as well as the development of antagonistic bacteria (16, 19). The use of suppressive compost provides effective biological control of plant pathogens. The use of mature compost colonized by antagonistic microorganisms is essential for its suppressiveness and for enhanced plant growth. Therefore, it is crucial to define parameters for compost maturity and quality control measures for the application of compost in agriculture.

Results

Effect of Composted MSW (CMSW) on cotton-plant development

Different amounts of CMSW (Amnir Recycling Industry, Afula, Israel) were added to the soil in order to study its influence on plant growth (Figs. 1 and 2). All experiments were conducted under controlled conditions in the greenhouse, using 500ml polypropylene pots containing sandy soil with 0-80% CMSW. Cotton plants were grown for 18 days, and percent emergence was determined daily. At the end of the experiment, plant height was measured. The addition of 5% CMSW to the soil did not affect emergence rate relative to controls (100% soil). However, soil amended with 20, 40 and 80% CMSW exhibited a 2-, 14- and 15-day delay in emergence, respectively.

Fig. 2 shows the effect of CMSW soil amendments on plant growth. The addition of 2 and 5% CMSW enhanced plant growth relative to the control. However, the addition of 10% or more CMSW resulted in a significant reduction in cotton-plant height. The same trend was found when cotton plants dry weight was determined (data not shown).

Fig. 1: Effect of CMSW on cotton plant emergence. 0-80 : percentage of CMSW added to the soil.

Fig. 2: Effect of CMSW on cotton-plant height. 0-100 : percentage of CMSW added to the soil. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Suppression of soil-borne pathogens by CMSW

The ability of CMSW to suppress soil-borne plant pathogens was studied using 1) cotton plants inoculated with *R. solani* which causes damping-off, and *F. oxysporum* which causes wilt disease; and 2) cucumber seedlings inoculated with *P. aphanidermatum* which causes damping-off. Plants (cotton or cucumber) were grown for 7, 14 or 21 days with *P. aphanidermatum*, *R. solani* or *F. oxysporum*, respectively. Experiments were conducted in controlled conditions in the greenhouse, using 500ml polypropylene pots containing sandy soil or peat with different amount of CMSW.

Suppression of *R. solani* by CMSW

R. solani which causes damping-off of plants was added to the soil. The inoculation was produced in a chopped potato-soil mixture (19). At the end of each experiment, the percentage of diseased plants and disease index (0-no disease, 5-dead plant) were determined. Fig. 3 shows *R. solani* suppression with increasing amount of CMSW in the soil: 2-40% CMSW suppressed the disease by about 80% relative to the control.

Fig. 3: Suppression of *R. solani* in cotton plants by CMSW. 0-40 : percentage of CMSW added to the soil. A) percent of diseased plants; B) disease index: disease severity index between 0-5. 0--healthy plants; 5--dead plants. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Suppression of *F. oxysporum* by CMSW

Wilt disease caused by *F. oxysporum* was tested over two growth-cycle. During the first growth cycle, 93 and 97% disease reduction was observed with 5 and 10% CMSW, respectively (Fig. 4). During the second growth-cycle disease incidence and severity increased significantly in the 5% CMSW treatment, whereas this increase was not significant in the 10% CMSW treatment.

Fig. 4: Suppression of *F. oxysporum* over two cycles of growth of cotton plants by CMSW (5 and 10% CMSW in the soil). Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Suppression of *P. aphanidermatum* by CMSW in container media

Suppression of *P. aphanidermatum* by CMSW was studied over two cycles of growth using cucumber plants in two media: 1) sandy soil contain 10% CMSW; 2) peat contain 50% CMSW.

In container media damping-off was tested in different amount of *P. aphanidermatum* inoculum, 0-7000 cfu (colony forming unit) per liter of dry container media (Fig. 5). At inoculum level of 5000 cfu and greater, *P. aphanidermatum* was reduced by 64% by CMSW, container media which consists 50% CMSW, 78% of the plants were healthy compared to 38% in the control.

Fig. 5: Suppression of *P. aphanidermatum* in container media in different inoculation densities. Squares - 50% CMSW in peat; Circles - peat. Lines indicates standard-error ($p=0.05$).

The influence of CMSW in container media on *P. aphanidermatum* cfu was tested in two cycles of growth, two weeks each cycle (Fig 6). Cucumbers plants were grown on container media which contain 50% CMSW. Different densities of the pathogen were added to the media, between 0 and 5000 cfu/L media.

In the media which contain CMSW disease severiry was low (90% healthy plants) in the two growth cycles, relative to the peat media (control) which resulted in 50% diseased plants in the first growth-cycle and 100% disease plants in the second growth-cycle.

Fig. 6: Suppression of *P. aphanidermatum* in container media in different inoculation densities, in two growth-cycles. Black circles - peat in the first growth-cycle; White circles - peat in the second growth cycle; Black triangles - peat:CMSW (1:1) in the first growth-cycle; White triangles - peat:CMSW (1:1) in the second growth-cycle. Lines indicates standard -error ($p=0.05$).

In order to find correlation between inoculum density and the percentage of diseased plants in container media, samples of the media were taken from the treatment of 5000 cfu, in the end of the first cycle. The cfu of *P. aphanidermatum* was determined on selective medium (20), significant correlation ($r=0.993954$) was found between disease percentage and *P. aphanidermatum*-cfu (Fig. 7). Significant correlation was found between media which contain peat or peat and CMSW and between the two growth-cycles of the different media.

Fig. 7: Suppression of *P. aphanidermatum* in container media, A) Percent of diseased plants; B) final inoculum density of the pathogen, in two cycle growth (I, II). C1=CMSW; Inoculum density in all treatments in the beginning was 5000 cfu/L media. Lines indicates standard-error ($p=0.05$).

Suppression of *P. aphanidermatum* by CMSW in sandy soil

Damping-off caused by *P. aphanidermatum* was tested in sandy soil contain 10% of CMSW in two growth cycles. The initial density of *P. aphanidermatum* was 2000 cfu/L media. Disease was reduced by 33% following the addition of 10% CMSW (Fig. 8), in the first growth cycle. In the second growth cycle there was 84% disease reduction.

Fig. 8: Suppression of *P. aphanidermatum* over two cycles of growth of cucumber plants by CMSW in soil. Squares -- 10% CMSW in soil; Circles -- soil (control). The initial density of *P. aphanidermatum* was 2000 cfu/L media. Lines indicates standard error (p=0.05).

Influence of compost maturity on the suppression of soil-borne pathogen

Controlled composting was performed to study the influence of composting time on the suppression of soil-borne pathogens.

MSW was collected over a 24 h period from the city of Afula in Israel, and mechanically mixed in a MSW classifying facility. After sieving and eliminating the >5-cm fraction, the raw material was divided into three plastic boxes (about 1 m³ each) and composted for 120 days. The compost was turned, mixed, and sampled after manually removing glass, plastics and metals at 14, 30, 45, 75, 120, 240 days. Samples (10 L) were stored at 4°C (4).

CMSW samples taken at the various stages of composting were added to soil inoculated with *R. solani* or *F. oxysporum*. *R. solani* was suppressed only when the CMSW was about 2.5 months old (Fig. 9): composts from 0.5-, 1- and 1.5-month old batches did not suppress the disease, which infected 80% of the plants; compost after 2.5, 4, 8 and 24 months suppressed *R. solani* by 50%. Different results were obtained for *F. oxysporum* (Fig. 10), which was suppressed by 0.5-, 1- and 1.5-month-old compost by 80%. 93% suppression was found with compost after 2.5 or more months of composting.

Fig. 9: Effect of CMSW age on its suppression of *R. solani* in cotton plants. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Fig. 10: Effect of CMSW age on its suppression of *F. oxysporum* in cotton plants. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

It was concluded that thermophilic microorganisms are involved in the suppression of *F. oxysporum*. The temperature of the 0.5-month-old compost pile was around 70°C: from the literature, *Bacillus* spp. are known to survive at such temperatures, and to be antagonist to some soil-borne pathogens. Suppression of *F. oxysporum* could be also, due to chemicals inhibitors exist in the CMSW. This phenomena needs further study.

Mechanism of *R. solani* suppression by CMSW

To understand the mechanism involved in the suppression of *R. solani* by CMSW, four treatments were applied to reduce the microbial population of the compost. Autoclaving (two times, 1 h each, 121°C) and gamma-irradiation (5 Mrad/h) negated suppression (Fig. 11). Disease severity following autoclaving and gamma-irradiation treatments were 92 and 86%, respectively, in comparison to the soil (control) and mature CMSW treatments resulted in 86 and 15% disease severity, respectively.

The autoclaved CMSW was washed with four volumes of water to remove the liquid chemical fractions resulting from the autoclave procedure: this treatment also resulted in the negation of CMSW's suppressive ability.

Fig. 11: Effect of autoclaving and gamma-irradiating CMSW on its suppression of *R. solani* in cotton plants. CMSW-- 10% CMSW in soil; Autoclave-- autoclaving of CMSW twice for 1 h each, at 121°C; Autoclave wash-- autoclaved CMSW washed with four volumes of water; Gamma-- gamma-irradiation (5 Mrad/h) of CMSW. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Heat treatment (5 days at 60°C), a moderate sterilization, also eliminated CMSW suppression of *R. solani* (Fig. 12). Mature CMSW reduced the disease by about 89%, whereas following the heat treatment the percentage of diseased plants reached 90. It is assumed that the biological vacuum created by the heat treatment enabled the pathogen to colonize the media more aggressively.

Fig. 12: Effect of heat-treating CMSW on its suppression of *R. solani* in cotton plants. CMSW-- 10% CMSW in soil; CMSW Heated-- heat treatment for 5 days at 60°C. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Another proof of microbial involvement in the suppression of *R. solani* was provided by adding 5% (v/v) mature CMSW to the heat-treated CMSW. This treatment restored suppression to levels observed with mature CMSW (Fig. 13). The incubation of heat-treated CMSW with 5% mature CMSW for 5 days resulted in a damping-off level of 43%, as compared to 95% with heat-treated CMSW alone.

Fig. 13: Effect of adding mature CMSW to heat-treated CMSW on its suppression of *R. solani* in cotton plants. CMSW-- 10% CMSW in soil; CMSW heated-- heat treatment for 5 days at 60°C; CMSW-heated + raw CMSW-- heated CMSW inoculated with 5% mature CMSW. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Mechanism of suppression *F. oxysporum* by CMSW

Heat treatment of CMSW did not affect the suppression of *F. oxysporum* (Fig. 14). Disease level with the heat-treated CMSW was the same as with mature CMSW: 0 and 10%, respectively during the first growth cycle and about 30% for each during the second growth cycle. This results suggest that thermophilic microorganisms involved in the suppression of *F. oxysporum*, survived the heat treatment.

Fig. 14: Effect of heat treatment of CMSW on its suppression of *F. oxysporum* in cotton plants. CMSW-- 10% CMSW in soil; CMSW heat -- heat treatment for 5 days at 60°C. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Microbial population and activity in CMSW

To elucidate the involvement of the CMSW's microbial population in CMSW suppressive activity, its identity and activity were determined in three CMSW media: mature CMSW (10% CMSW in soil), heated CMSW (mature CMSW treated for 5 days at 60°C), and heated CMSW inoculated with mature CMSW (5%, v/v). Three methods were used: 1) fungi and bacteria were determined on agar plates (Table 1); 2) microbial activity was monitored by measuring the rate of hydrolysis of fluorescein diacetate (FDA), FDA is hydrolyzed by a number of enzymes in living cells, such as proteases, lipases, and esterases (Table1); 3) the level of CO₂ emission was determined by gas chromatograph (Fig.15).

Table 1: Microbial population and activity of CMSW

	Mature CMSW	CMSW heat	CMSW heat+ CMSW
Fungi (cfu/g) ⁽¹⁾	$1.5 * 10^5$	$< 10^2$	$0.6 * 10^3$
Bacteria (cfu/g) ⁽²⁾	$4.5 * 10^8$	$1.4 * 10^3$	$1.5 * 10^8$
FDA (μ gr flourescein/ g) ₍₃₎	1.6a ⁽⁴⁾	1.03b	1.9a

(1) Fungi were determined on Martin agar.

(2) Bacteria were determined on Difco nutrient agar.

(3) FDA (Sigma Chemical Co) was measured as proposed by Schnurer and Rosswall (21), after a 2 h incubation.

(4) Different letters indicate significant differences between treatments (p=0.05).

Fig. 15: Accumulated CO₂ emission from CMSW media. CMSW-- mature CMSW; CMSW heat-- mature CMSW treated for 5 days at 60°C; CMSW-heat+CMSW- - heated CMSW inoculated with mature CMSW (5%, v/v). CO₂ level was measured with a GC model 580, GOW-MAC Instruments Co., USA equipped with Propak Q column)

Microscopic observation of *R. solani* in CMSW

Microscopic observation was performed to understand the mechanism involved in the suppression of *R. solani* by CMSW. The fungus was grown on a nylon 100µm mesh and then transferred to soil with CMSW after 1 h (as seen in a light microscope *400), there were many bacteria along the fungal hyphae in the CMSW treatment. In the control treatment (soil), some bacteria were observed, but not near the hyphae. After 12 h the fungal hyphae in the CMSW treatment changed shape and after 24 h complete lysis had occurred.

Isolation of antagonistic bacteria from CMSW

Bacteria were isolated from CMSW-meshes to pinpoint antagonistic activity against *R. solani*. Some of the bacteria exhibited chitinolytic activity on chitin-agar plates and growth inhibition of *R. solani*. Eight strains were tested in a bioassay in the greenhouse: upon addition to the soil, two of the isolates suppressed *R. solani* by about 50% (Fig. 16). One of these strains was identified by fatty-acid-profile analysis as *Enterobacter agglomerans*, a strain which is known as an efficient antagonistic bacterium.

Fig. 16: Effect of bacteria isolated from CMSW on suppression of *R. solani* in cotton plants. Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Addition of *Trichoderma harzianum* to CMSW

T. harzianum, a known *R. solani* antagonist, was not found in the mature CMSW, as determined by growth on Trichoderma-selective medium. To improve the suppressiveness of CMSW against *R. solani*, soil, mature CMSW and heated CMSW were inoculated with *T. harzianum* (Fig. 17). The antagonistic fungus successfully colonized all three media, reaching a level of 10^6 cfu/g soil, but the resultant level of suppression was not higher than with mature CMSW alone.

Fig. 17: Effect of adding *Trichoderma harzianum* to CMSW on suppression of *R. solani* in cotton plants. Soil Tr-- soil inoculated with *T. harzianum* (10^6 cfu/g soil); CMSW-Tr-- CMSW inoculated with *T. harzianum* (10^6 cfu/g soil); CMSW-Heated-- ; heat treatment of CMSW for 5 days at 60°C; CMSW-Heated-Tr-- heated CMSW inoculated with *T. harzianum* (10^6 cfu/g soil). Different letters heading the columns indicate significant differences between treatments ($p=0.05$).

Conclusions

CMSW suppressed the soil-borne pathogens *R. solani*, *P. aphanidermatum* and *F. oxysporum*. In cotton, CMSW reduced the damping-off caused by *R. Solani* by about 80% relative to the control (soil). *F. oxysporum*, which causes wilt disease of cotton, was reduced by 98% relative to the control, and *P. aphanidermatum*, which causes damping-off in cucumber, was reduced by 33%.

In the experimental system with *R. solani*, autoclaving (two times, 1 h each, 121°C), gamma-irradiation (5 Mrad/h) and heat treatment (5 days at 60°C) all negated suppression. Thus, it can be concluded that the mechanism involved in *R. solani* suppression is biological. Further evidence of microorganisms involvement was provided when 5% (v/v) CMSW was added to the heat-treated CMSW. This addition caused the recolonization of antagonistic microorganisms and restoration of the compost's suppressive ability.

In the experimental system with *F. oxysporum* heat treatment (5 days at 60°C) did not eliminate suppressive ability, suggesting that either a thermophilic population is responsible for this particular suppression or that induce resistance or physical and chemical factors are involved.

In the case of *R. solani*, the suppressive mechanism is presumed to be lysis of fungal hyphae. Microscopic examination of *R. solani* grown on nylon meshes and incubated in CMSW-amended soil indicated the possible involvement of bacteria. Some of the bacteria were isolated from the compost meshes. Eight isolates showed the ability to degrade chitin and to inhibit *R. solani* growth in Petri dishes. Two of these isolates suppressed *R. solani* in cotton in greenhouse experiments. One of them was identified by fatty-acid-profile analysis as *E. agglomerans*; the other could not be identified by this method.

T. harzianum was added to mature or heated compost. The fungi colonized the heated compost and significantly suppressed *R. solani*. In contrast, *T. harzianum* did not improve the suppressive ability of the mature compost despite the fact that it colonized the compost successfully.

Understanding the microbial mechanisms involved in the suppression of soil-borne pathogens could contribute to the quality improvement of compost via its enrichment with beneficial microorganisms. The key to large-scale utilization of disease-suppressive compost is the development of reproducible products with defined, consistent properties.

References

- 1) Baker, K. F. 1987. Evolving concepts of biological control of plant pathogens. *Annu. Rev. Phytopathol.*, 25: 67-85.
- 2) Bruns, C., Ahlers, S., Gatinger, A., Schöler, C., Vogtman, H. and Wolf, G. 1996. The suppressive effects of composted separately collected organic waste and yard waste compost on two important soilborne plant pathogens. *In: The Science of Composting*. pp. 1094-1095. M. Bertoldi, P. Sequi, B. Lemmes and T. Papi, (Eds.). Blackie A & P: London.
- 3) Chef, D. G., Hoitink, H. A. J. and Madden, L. V. 1983. Effects of organic components in container media on suppression of *Fusarium* wilt of chrysanthemum and flax. *Phytopathology*, 73: 279-281.
- 4) Chefetz, B., Hatcher, P. G., Hadar, Y. and Chen, Y. 1996. Chemical and biological characterization of organic matter during composting of municipal solid waste. *J. Environ. Qual.*, 25: 776-785.
- 5) Chen, Y. and Hadar, Y. 1987. Composting and use of agricultural wastes in container media. *In: Composts: Production Quality and Use*. pp. 71-77. M. P. Ferranti, P. L'Hermite and F. Zucchini, (Eds.). Elsevier Science: London.

- 6) Chung, Y. R., Hoitink, H. A. J., Dick, W. A. and Herr, L. J. 1988. Effects of organic matter decomposition level and cellulose amendment on the inoculum potential of *Rhizoctonia solani* in hardwood bark media. *Phytopathology*, 78: 836-840.
- 7) Craft, C. M. and Nelson, E. B. 1996. Microbial properties of composts that suppress damping-off and root rot of creeping bentgrass caused by *Pythium graminicola*. *Appl. Environ. Microbiol.*, 62: 1550-1557.
- 8) Gera, J. V. O. and Wilma, J. M. V. G. 1996. Possibilities for biological control of *Pythium* root rot in ornamental bulb culture with composted organic household waste. *In: The Science of Composting*. pp. 1374-1375. M. Bertoldi, P. Sequi, B. Lemmes and T. Papi, (Eds.). Blackie A & P: London.
- 9) Hadar, Y. and Gorodecki, B. 1991. Suppression of germination of sclerotia of *Sclerotium rolfsii* in compost. *Soil Biol. Biochem.*, 23: 303-306.
- 10) Hadar, Y. and Mandelbaum, R. 1986. Suppression of *Pythium aphanidermatum* damping-off in container media containing composted liquorice roots. *Crop Prot.*, 5: 88-92.
- 11) Hoitink, H. A. J. 1980. Composted bark, a lightweight medium with fungicidal properties. *Plant Dis.*, 64: 142-147.
- 12) Hoitink, H. A. J., Boehm, M. J. and Hadar, Y. 1993. Mechanisms of suppression of soilborne plant pathogens in compost-amended substrates. *In: Science and Engineering of Composting: Design, Environmental, Microbiological and Utilization Aspects*. pp. 601-622. H. A. J. Hoitink and H. M. Keener, (Eds.). Renaissance Publications: Worthington, OH.

- 13) Hoitink, H. A. J., Stone, A. G. and Grebus, M. E. 1996. Suppression of plant disease by composts. *In: The Science of Composting*. pp. 373-381. M. Bertoldi, P. Sequi, B. Lemmes and T. Papi, (Eds.). Blackie A & P: London.
- 14) Inbar, Y., Hadar, Y. and Chen, Y. 1993b. Recycling of cattle manure: the composting process and characterization of maturity. *J. Environ. Qual.*, 22: 857-863.
- 15) Kuter, G. A., Hoitink, H. A. J. and Chen, W. 1988. Effect of municipal sludge compost curing time on suppression of *Pythium* and *Rhizoctonia* diseases of ornamental plants. *Plant Dis.*, 72: 731-756.
- 16) Kwok, O. C. H., Fahy, P. C., Hoitink, H. A. J. and Kuter, G. A. 1987. Interactions between bacteria and *Trichoderma harzianum* in suppression of *Rhizoctonia* damping-off in bark compost media. *Phytopathology*, 77: 1206-1212.
- 17) Mandelbaum, R., Gorodecki, B. and Hadar, Y. 1985. The use of composts for production of disease suppressive container media. *Phytoparasitica*, 13: 185 (Abstract).
- 18) Mandelbaum, R. and Hadar, Y. 1990. Effect of available carbon source on microbial activity and suppression of *Pythium aphanidermatum* in composted and peat container media. *Phytopathology*, 80: 794-804.
- 19) Nelson, E. B., Kuter, G. A. and Hoitink, H. A. J. 1983. Effect of fungal antagonist and compost age on suppression of *Rhizoctonia solani* damping-off in container media amended with composted hardwood bark. *Phytopathology*, 73: 1457-1462.
- 20) Schmitthenner, A. F. 1979. *Pythium* species: isolation, biology and identification. *In: Advances in Turfgrass Pathology, Proceeding of Symposium on Turfgrass Disease*. Harcourt Brace Jovanovich, Inc. Duluth, MN. Pp. 33-36.

- 21) Shnürer, J. and Rosswall, T. 1982. Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter. *Appl. Environ. Microbiol.*, 43: 1256-1261.
- 22) Schöler, C., Biala, J., Gottschall, S., Ahlers, S. and Vogtman, H. 1989. Suppression of root rot on peas, beans and beetroots caused by *Pythium ultimum* and *Rhizoctonia solani* through the amendments of growing media with composted organic household waste. *Pytopathology*, 127: 227-238.
- 23) Serra-Wittling, C.; 1995. Valorisation de composts d'ordures ménagères en protection des cultures: Influence de l'apport de compost sur le développement de maladies d'origine tellurique et le comportement de pesticides dans un sol. Thèse présentée en vue de l'obtention du diplôme de docteur. l'INA PARIS-GRIGNON.
- 24) Volland, R. P. and Epstein, A. H. 1994. Development of suppressiveness to diseases caused by *Rhizoctonia solani* in soils amended with composted and noncomposted manure. *Plant Dis.*, 78: 461-466.