BARD

FINAL REPORT

PROJECT NO. IS-2949-97

Physiological Regulation of Root System Architecture: The Role of Ethylene and Phosphorus

A. Eshel, J.P. Lynch, K.M. Brown

2002

BARD Final Scientific Report

Date of Submission of the report: December 30th 2001 ✓

BARD Project Number: IS-2949-97R

Project Title: Physiological Regulation of Root System Architecture:
The Role of Ethylene and Phosphorus

Investigators

Institutions

Principal Investigator (PI):

Amram Eshel

Tel-Aviv University

Co-Principal Investigator (Co-PI):

Jonathan P. Lynch
Kathleen M. Brown

Pennsylvania State University
Pennsylvania State University

Collaborating Investigators:

Keywords *not* appearing in the title and in order of importance. Avoid abbreviations.
Beans, Tomatoes, Fractal Analysis, Aeroponics

Abbreviations commonly used in the report, in alphabetical order:

Budget: IS: \$ 113,000

US: \$ 172,000

Total: \$ 285,000

Signature
Principal Investigator

Signature
Authorizing Official, Principal Institution

BARD Final Scientific Report

Publication Summary (numbers)

	Joint IS/US authorship	US Authors only	Israeli Authors only	Total
Refereed (published, in press, accepted)	1	3		4
Submitted, in review, in preparation	2	3	1	6
Invited review papers		1		1
Book chapters				
Books				
Master theses				
Ph.D. theses				
Abstracts		8		8
Not refereed (proceedings, reports, etc.)				

Postdoctoral Training: List the names and social security/identity numbers of all postdocs who received more than 50% of their funding by the grant.

Rina Shabtai-Wiezmann - Israeli ID 05883558-8

Cooperation Summary (numbers)

	From US to Israel	From Israel to US	Together, elsewhere	Total
Short Visits & Meetings	1			1
Longer Visits (Sabbaticals)				

Description of Cooperation:

On May 1999 the US investigators visited the laboratory in Israel for a short time for discussion of the
results of the experiments and coordination of the workplan.

Patent Summary (numbers)

	Israeli inventor (s) only	US inventor (s) only	Joint IS/US inventors	Total
Submitted				
Issued (allowed)				
Licensed				

BARD Final Scientific Report

Abstract

Specific Objectives and Related Results:

1) *Determine the effect of phosphorus availability on ethylene production by roots. Test the hypothesis that phosphorus availability regulates ethylene production*

Clear differences were found between the two plants that were studied. In beans ethylene production is affected by P nutrition, tissue type, and stage of development. There are genotypic differences in the rate of ethylene production by various root types and in the differential in ethylene production when P treatments are compared. The acceleration in ethylene production with P deficiency increases with time. These findings support the hypothesis that ethylene production may be enhanced by phosphorus deficiency, and that the degree of enhancement varies with genotype. In tomatoes the low-P level did not enhance significantly ethylene production by the roots. Wildtype cultivars and ethylene insensitive mutants behaved similarly in that respect.

2) *Characterize the effects of phosphorus availability and ethylene on the architecture of whole root systems. Test the hypothesis that both ethylene and low phosphorus availability modify root architecture.*

In common bean, the basal roots give rise to a major fraction of the whole root system. Unlike other laterals these roots respond to gravitropic stimulation. Their growth angle determines the proportion of the root length in the shallow layers of the soil. A correlation between ethylene production and basal root angle was found in shallow rooted but not deep-rooted genotypes, indicating that acceleration of ethylene synthesis may account for the change in basal root angle in genotypes demonstrating a plastic response to P availability. Short-time gravitropic response of the tap roots of young bean seedlings was not affected by P level in the nutrient solution.

Low phosphorus specifically increases root hair length and root hair density in *Arabidopsis*

We tested 7 different mutants in ethylene perception and response and in each case, the response to low P was lower than that of the wild-type. The extent of reduction in P response varied among the mutants, but every mutant retained some responsiveness to changes in P concentration. The increase in root hair density was due to the increase in the number of trichoblast cell files under low P and was not mediated by ethylene. Low P did not increase the number of root hairs forming from atrichoblasts. This is in contrast to ethylene treatment, which increased the number of root hairs partly by causing root hairs to form on atrichoblasts.

3) *Assess the adaptive value of root architectural plasticity in response to phosphorus availability.*

A simulation study indicated that genetic variation for root architecture in common bean may be related to adaptation to diverse competitive environments. The fractal dimension of tomato root system was directly correlated with P level.

Conclusions & Implications

A. P level affects root architecture and anatomy of the common bean in an adaptive way causing shallowness of the root system and increase in root hair length and density. Future selection and breeding programs for P efficient genotypes should target these traits.

B. Overall exploration of the rooting volume can be assessed by the Fractal Dimension of the root system. This parameter should be used in the future for characterizing the effectiveness of plant root systems and their relative ability to utilize soil resources.

C. Plant responses to ethylene and low P are mediated by separate signal transduction pathways and ethylene-insensitive or over-sensitive mutants could not be expected to be more P efficient.

D. P concentration of 25 μ M, which is 1% regular fertilization level for greenhouse tomatoes, is sufficient for normal growth and yield. It is therefore proposed that P fertilization for greenhouse tomatoes can be reduced by at least one order of magnitude, resulting in reduction of costs, but more importantly reduction of environmental pollution by greenhouse effluents. A research proposal aimed at achieving this goal was submitted to the Israeli Vegetable Growers Board.

BARD Final Scientific Report

Achievements

This research focused on the similarities between the effects of ethylene and low P level on root system anatomy and architecture. In the common bean P nutrition, tissue type, and stage of development affected ethylene production. There were genotypic differences in the rate of ethylene production by various root types and in the differential in ethylene production when P treatments are compared. These findings support the hypothesis that ethylene production may be enhanced by phosphorus deficiency, and that the degree of enhancement varies with genotype.

On the other hand in tomatoes we did not find a consistent increase of ethylene production by low P treatment in wildtype or ethylene-insensitive mutants. Moreover, neither external ethylene application nor low P treatment altered the allometric root-shoot relationships or basic qualitative characteristics of tomato root architecture. There were similar findings in a concomitant study that has used ethylene-insensitive mutant of petunia. We therefore conclude that the two plant families differ in their response to P deficiency and ethylene.

Root Morphology & Architecture

Gravitropism

In common bean, typical lateral roots were insensitive to the gravitropic stimulus however, the basal roots, exhibited a positive gravitropic response that was not as strong as that of the taproots. Since the basal give rise to most of the roots which comprise the bean root system, the angle of their growth determines the proportion of the root length found in the shallow layers of the soil, where phosphorus is typically found at higher concentrations. Genetic variation in basal root angle has been correlated with field performance of common bean genotypes. We have shown that basal root angle is affected by manipulation of ethylene synthesis. Promoting ethylene synthesis results in shallower roots, while inhibiting it results in deeper roots. A correlation between ethylene production and basal root angle was found in shallow rooted but not deep-rooted genotypes, indicating that acceleration of ethylene synthesis may account for the change in basal root angle in genotypes demonstrating a plastic response to P availability.

Root Hairs

Low phosphorus specifically increases both root hair length and root hair density in *Arabidopsis*. Our initial data on ethylene involvement seemed to indicate a possible role for ethylene in the low P response. Although we were unable to reduce root hair density or length under low P with an ethylene synthesis inhibitor, we were able to do so with the ethylene synthesis inhibitor and with the ethylene action inhibitors. Likewise, the ethylene precursor ACC increased root hair length and density in high P plants to create a morphology similar to low P treatment.

BARD Final Scientific Report

If ethylene action was important for the increases in root hair length and density, we would expect those responses to be greatly decreased in mutants with impaired ethylene perception. We tested 7 different mutants in ethylene perception and response and in each case, the response to low P was reduced. We further dissected the root hair density response by investigating the changes in root growth that could produce increased root hair density. We found that roots grown with P deficiency had more and smaller cortical cells, and thus increased the number of potential root hair cells. The anatomy remained consistent with the P treatment and not ethylene manipulation. Likewise, the anatomy of ethylene response mutants was altered with low P treatment in the same way as wildtype plants. Therefore, the increase in root hair density is not mediated by ethylene. Another factor that affected root hair density under low P was the length of individual trichoblast cells. However, ethylene did not mediate this aspect of root hair density. Also, low P did not increase the number of root hairs forming from atrichoblasts. This was in contrast to ethylene treatment, which increased the number of root hairs partly by causing root hairs to form on atrichoblasts. This is further evidence that ethylene and low P act via separate mechanisms to increase root hair density.

Fractal Analysis

Root exploration of the rooting volume is a most important characteristic for determining plant adaptation to low P conditions because of the low mobility of P in soil. One of the objectives of this research program was to determine whether the mathematical parameter called Fractal Dimension could be used as a quantitative measure of this complex morphological characteristic. It was found that the indeed the analyzed plant root systems are amenable to fractal analysis. Their two-dimensional Fractal Dimension corresponded to overall root system size and its degree of exploration of the soil. We therefore conclude that this Fractal Dimension can indeed be used for quantitative comparison between different root systems, in terms of their effectiveness in utilization of immobile edaphic resources.

Root Competition

Competition for nutrients among neighboring roots occurs when their individual depletion volumes overlap, causing a reduction in nutrient uptake. By exploring different spatial niches, plants with contrasting root architecture may reduce the extent of competition among neighboring root systems.

A dynamic geometric model, *SimRoot*, was used to simulate common bean root growth and to compare the overlap of depletion volumes. The main objectives of this study were i) to evaluate the impact of root architecture on competition for phosphorus among neighboring plants, and ii) to compare the magnitude of competition among roots of the same plant vs. roots of neighboring plants. By varying the gravitropism of basal roots, we simulated three distinct root architectures corresponding with observed genetic variation for root architecture in this species. Combinations of roots having the same architecture resulted in more intense inter-plant competition. Genetic variation for root architecture in common bean may be related to adaptation to diverse competitive environments.

BARD Final Scientific Report

Carbon Utilization

Alteration of the relative investment of plant carbohydrate resources is another mechanism by which plants adapt to scarcity of edaphic resources. P efficient genotypes are expected to show higher plasticity in the allocation of carbohydrates to root structure and function. A common response to low phosphorus availability is increased relative biomass allocation to roots. The resulting increase in root:shoot ratio presumably enhances phosphorus acquisition but may also reduce growth rates by diverting carbon to the production of heterotrophic rather than photosynthetic tissues. To assess the importance of increased carbon allocation to roots for the adaptation of plants to low P availability, we constructed carbon budgets for four common bean genotypes with contrasting adaptation to low phosphorus availability in the field. Low phosphorus plants utilized a significantly larger fraction of their daytime net carbon assimilation for root respiration. Genotypes also had similar rates of P absorption per unit root weight and plant growth per unit of P absorbed. However, P-efficient genotypes allocated a larger fraction of their biomass to root growth, especially under low P conditions.

P Fertilization of Greenhouse Tomatoes

In the course of this research it was shown that fertilization of greenhouse tomatoes with 25 μM P, which is 1% the level of this nutrient used in common agricultural practice, will suffice for normal plant development and yield. It is therefore proposed that P fertilization for greenhouse tomatoes can be greatly reduced. Reduction in P input down to 10% of current usage is expected to be successful and result not only in reduction of costs, but more importantly reduction of environmental pollution by greenhouse effluents. A research proposal aimed at changing the common practice of P fertilization of greenhouse tomatoes by reducing input levels by at least one order of magnitude was submitted to the Israeli Vegetable Growers Board.

Published Articles

- Nielsen, KL, A Eshel, JP Lynch. 2001. The effect of phosphorus availability on the carbon economy of contrasting common bean (*Phaseolus vulgaris* L.) genotypes. *Journal of Experimental Botany*, 52:329-339.
- Liao, H, G Rubio, X Yan, A Cao, KM Brown, JP Lynch. 2001. Effect of phosphorus availability on basal root shallowness in common bean. *Plant and Soil* 232:69-79.
- Rubio, G, T Walk, Z Ge, X Yan, H Liao, JP Lynch. 2001. Root gravitropism and below-ground competition among neighboring plants: a modeling approach. *Annals of Botany* 88:929-940.
- Lynch, JP and KM Brown. 2001. Topsoil foraging – an architectural adaptation to low phosphorus availability. *Plant and Soil* 237:225-237.