BARD

77

FINAL REPORT

PROJECT NO. US-1160-86

Bovine Relaxin: A Placental Source and Effects on Prostaglandin and Steroid Metabolism

M.J. Fields, M. Shemesh, P.A. Fields

1991

100

100

100

100

100

A. Standard BARD Cover Page for Scientific Reports

Date: December 9, 1991

BARD
P.O. Box 6
Bet Dagan, ISRAEL

BARD Project No. US-1160-86

Title

Bovine Relaxin: A Placental Source and Effects on Prostaglandin and Steroid
Metabolism

Investigators' Name

Investigator's Institutions

(Principal listed first)

Michael J. Fields

University of Florida

Mordechai Shemesh

Kimron Veterinary Institute

Phillip A. Fields

University of South Alabama

Anna-Riitta Fuchs

Cornell Medical School

Michael Diskin

Ag Food Development Authority

Galway, Ireland

Project's starting date: January 1, 1988

Type of Report: 1st Annual ☐ 2nd Annual ☐ Final ☒

Signatures

Principal Investigator

Institution's Authorizing Official

B. Table of Contents

Page

A. Cover Page	1
B. Table of Contents	2
C. Abstract	4
D. Objectives	5
E. Description of Research Results	7
1. Isolation and Amino Acid Sequencing of High Molecular Weight Relaxins	7
- Extraction of Acidic High MW Relaxins	8
- Isolation Via PVDF - Relaxin-Immunoaffinity	9
- Cyanogen Bromide Cleavage	11
- Isolation of Prorelaxin Via 2-D Electrophoresis	12
- Transfer of Prorelaxin onto PVDF	14
- Internal Amino Acid Sequencing of Prorelaxin	17
- Conclusion	18
2. Extraction of Relaxin From Bovine Corpora Lutea and Placenta	20
- Relaxin Isolated From Bovine Corpora Lutea	20
- Attempts to Enhance the Recovery of Bovine Relaxin	26
- SDS Gel Separation of the A and B Chains of Relaxin	27
- Isoelectric Focusing Separation of the A and B Chains of Relaxin	32
- Two D-SDS Separation of A and B Chains of Relaxin	37
- Isolation of Relaxin From the Bovine Placenta	40
- Acid Acetone VS Tris Extraction of tissue	41
- Sephadex G5044	44
- SDS - Electrophoresis Gel Slices	45
- Parallelism of Bovine and Porcine Relaxin	46
- Conclusion	49
- Addendum	50
3. Effects of Relaxin on Steroidogenesis and Prostaglandin Synthesis	51
- Effect of Relaxin on Luteal Progesterone Synthesis	51
- Effect of Relaxin on Placental Steroid Metabolism	53
- Effect of Relaxin on Prostaglandin Synthesis	57
- Conclusion	60

4. Response of Nonpregnant and Periparturient Cows to Porcine Relaxin	62
- Evaluation of Different Routes of Administration of Relaxin to Cows	62
- Production of Antibodies to Porcine Relaxin in Cows	66
- Relaxin Induction of Parturition	69
- Relaxin Combined With Corticosteroids to Induce Parturition	72
- Conclusion	73
5. Oxytocin Receptors in Reproductive Tissues	74
- Oxytocin and Vasopressin Receptors in Early Pregnancy	74
- Oxytocin and Vasopressin Receptors in the Ovary	80
- Addendum	80
6. Large Luteal Cells and Their Secretory Granules	82
- Depletion of Secretory Granules with $\text{PGF}_{2\alpha}$	84
- Conclusion	91
7. Literature Cited	92
F. Description of Cooperation	97
G. Evaluation and Benefit to Agriculture	99
H. List of Publications	104

C. Abstract

Relaxin (RXN) is an enigma in the ruminant. The cow is responsive to porcine relaxin but little is known about the endogenous form of bovine relaxin. Although, the nucleotide and amino acid sequence of numerous species of RXN have been determined none of this information is available for the ruminants. This study evaluated the placenta as a source for isolation of sufficient quantities of RXN for chemical characterization. RXN was immunohistochemically localized in the syncytiotrophoblast and subsequently shown in isolates from the cotyledon that gave a linear dose response in the pRXN RIA. Attempts to increase the recoverable yields of bRXN were not successful. Although microtechniques for successfully separating the A and B chain of pRXN and determining their amino acid sequence were developed, insufficient quantities of bRXN preempted their application. These techniques were, however, successfully utilized to show that the high molecular weight porcine immunoreactive relaxin-like peptides were processed prohormone. We are encouraged that most recently we have shown with western blots a 17,000 Dalton immunoreactive relaxin from the bovine placenta that could be a form of the bovine prohormone.

RXN had a clear dose and time related effect on the production of P_4 or $PGF_{2\alpha}$ and PGE in either the caruncle or cotyledon in a few in vitro experiments but we could not repeat these observations. It was concluded that if RXN is involved in steroidogenesis or PG synthesis the action is indirect and mediated through some other compound. When pRXN was administered with or without corticosteroids to late pregnant cows the interval to parturition was decreased. The mid-pregnancy increase in oxytocin receptors (OT-Rc) in the endometrium and myometrium was unexpected. However, the sharp increase in oxytocin receptors (OT-RC) in the endometrium and fetal membranes at or preceding labor indicates a role for OT at this time that may complement the action of RXN. During the cycle but not pregnancy luteal secretory granules were shown to contain OT and neurophysin. In the course of these studies we observed two different populations of large luteal cells, the Alpha cell ($PGF_{2\alpha}$ insensitive) and the Beta cell sensitive to subluteolytic doses of $PGF_{2\alpha}$.

D. Objectives

The finding of placental relaxin in another ruminant, the ewe as well as placental and/or endometrial relaxin in the human, rabbit, cat, horse and guinea pig indicates there may be a similar source of relaxin in the cow, particularly given the extremely low luteal levels found in both the ewe and cow.

The specific objectives were to:

1. To identify the uterine (caruncle and intercaruncle endometrium) and fetal (cotyledon and non-cotyledonary chorion) tissues containing immunoreactive relaxin via the use of the homologous porcine relaxin radioimmunoassay.
2. Isolation of sufficient quantities of bovine relaxin that will lead to determination of the relaxin peptide amino acid sequence as well as building antibodies for development of an homologous bovine relaxin radioimmunoassay.
3. To identify the specific cell producing relaxin via immunohistochemical and immunocytochemical localization of bovine relaxin.
4. To determine the effects of relaxin on uterine-placental prostaglandin and steroid synthesis with dispersed cells of the cotyledon and caruncle.
5. To determine with dispersed uterine-placental cells whether relaxin synthesis in conjunction with progesterone is influenced by mediators of intracellular calcium and/or secretagogues that act via cyclic nucleotides. There is a paucity of information on factors regulating relaxin synthesis.
6. To determine the effect of various factors on the migration and exocytosis of secretory granules of binucleate cell. The content of secretory granules will be examined for relaxin via immunocytochemistry.

7. To clinically test relaxin under field conditions to determine if relaxin will reduce the incidence of retained placenta in heifers induced to calve with $\text{PGF}_{2\alpha}$ or glucocorticoids, a model shown to result in a high incidence of retained placenta.

The objectives of the research did not significantly change throughout the time frame of the grant. Our primary objective of evaluating the bovine placenta as a source of relaxin was met. However, without sufficient quantities of isolated bovine relaxin we were not in a position to attain the amino acid sequence of bovine relaxin nor build antibodies to this relaxin for development of an homologous bovine relaxin radioimmunoassay. In the course of these studies there was an opportunity to pursue two other areas of research that converge into the study of relaxin.

First, was the characterization of the oxytocin receptor in various reproductive tissues across a wide spectrum of the estrous cycle and pregnancy including parturition. Oxytocin and relaxin are both known for playing a critical role in the process of parturition. Characterizing the receptors of the uterus and placental tissues provides additional insight into the action of oxytocin. This information with that of relaxin contributes to a more informed understanding of the endocrine environment within which relaxin participates.

The second area introduced under the umbrella of this grant was that of luteal secretory granules. Even though the corpus luteum was shown to contain little immunoreactive relaxin and the focus of this grant was a placental source of relaxin, the only peptide shown to be packaged in luteal secretory granules across all species during pregnancy has been relaxin. Thus, study of the secretory granules was pursued as a source of relaxin in the cow. Specifically we were interested in the early formation of these granules.

In both cases the two introduced areas, oxytocin receptors and luteal secretory granules were very productive. These studies were complementary and did not distract or redirect monies from the primary objective of characterizing a placental source of relaxin in the cow.

E. Description of Research Results

1. Isolation and Amino Acid Sequencing of High Molecular Weight Relaxins.

Porcine relaxin, a hormone of pregnancy, consists of two polypeptide chains A and B linked by two inter-chain disulfide bonds (Schwabe and McDonald, 1977; Sherwood, 1988). It has a molecular weight of 6,000 and a basic pI of 10. Gast et al. (1980), and Gast (1982, 1983) showed that the pig corpus luteum contains mRNA that directs the synthesis of a 23,000 Da relaxin-immunoreactive protein and postulated that it is the primary translation product and that the membrane-dependent cleavage of a 3,000 Da signal peptide was the first step in relaxin biosynthesis (Fig. 1). Haley et al. (1982) determined the complete amino acid sequence of porcine preprorelaxin by cloning of relaxin cDNA (Fig. 2). Relaxin, like insulin, is initially

Figure 1. Schematic summary of the processing of porcine preprorelaxin to relaxin by the successive removal of the signal peptide (S) and the connecting peptide (C) by proteolytic digestion (Kemp and Niall, 1984; Sherwood, 1988).

synthesized as a single-chained prehormone with the overall structure: signal peptide/B chain/connecting peptide(C-peptide)/A chain as shown schematically in Figure 1. The function of C-peptide (104 amino acid residues) in relaxin is not known. It presumably helps in correct

Figure 2. Amino acid sequence of porcine preprorelaxin. Cleavage sites are indicated with vertical lines (Sherwood, 1988).

formation of disulfide bonds between A and B chains by directing proper folding of the precursor. Besides, its large size and the fact that its primary structure is highly conserved among species (more so than its A and B chains) led to the postulation that the C-peptide may contain peptide sequences with hormonal activities (Hudson et al., 1981; Kemp and Niall, 1984).

The recovery, purification, and amino acid sequencing techniques for high molecular weight, relaxin-like peptides isolated from porcine ovaries were developed for and applied to purification of bovine placental relaxin.

Extraction of Acidic, High Molecular Weight, Relaxin-Like Peptides from Porcine Ovarian Tissue. Porcine relaxin (6,000 Da, pI 10.) has routinely been purified in our laboratory from ovaries of the late pregnant sow utilizing the procedures previously described by Griss et al. (1967) and Fields et al. (1980). Briefly, with a Brinkman Polytron, frozen tissue was homogenized in an extraction solution of 3 parts 0.7 N HCl and 5 parts acetone (6 ml/g tissue) and incubated overnight at 4°C. The tissue homogenate was then centrifuged and the relaxin in the supernatant was precipitated with the addition of five volumes of cold acetone, collected by centrifugation, washed to remove HCl and solubilized with 0.2 M ammonium acetate, pH 5.0. This relaxin-containing crude extract was then fractionated by Sephadex G-50 column chromatography (Fig. 3) and the protein peak eluted at a volume corresponding to 6,000 Da was

Figure 3. Elution profile of pH 5 supernatant from acid-acetone crude extract of porcine ovarian tissue fractionated by Sephadex G-50. The column (5 X 90 cm) was pre-equilibrated and eluted by 0.01 M ammonium acetate, pH 5, at 60 ml/h. Fractions under peak IV, which contains 6,000 relaxin, were pooled and further purified by CM cellulose.

Fractions under peaks I and II were

pooled, heat treated, and passed through CM cellulose to remove the 6,000 Da relaxin. It's from this high Mr, acidic sample that the three high molecular weight, acidic, relaxin-immunoreactive peptides were later isolated by two-dimensional gel electrophoresis.

further purified by a cation-exchange chromatography on a pH 5.0 carboxymethyl cellulose (CM cellulose) column to obtain 6,000 Da relaxin of high purity. The proteins eluted at higher molecular weight region ($\geq 12,000$ Da) from the Sephadex G-50 fractionation were pooled and heated (5 min at 90 °C) to stop possible proteolysis. This sample was further purified and devoid of 6,000 Da relaxin by passing it through a pH 5.0 CM cellulose column. This time, only the flow through, i.e., unbound, acidic proteins, was collected. It was from this flow through that high molecular weight, acidic, relaxin-like proteins were isolated.

Isolation of Acidic, High Molecular Weight, Ovarian Peptides via PVDF-Relaxin-Immunoaffinity. Following a method described by Janssen Biotech (1984), a small piece (0.5 x 5 cm) of polyvinylidene difluoride membrane (PVDF; Immobilon-P, Millipore Inc., Bedford, MA) was coated with CM cellulose-purified porcine relaxin (lot# UF-16S) by passive blotting (spotting) technique to isolate rabbit antibody (polyclonal, lot# UF-1Q) against porcine relaxin. The isolated antibody was eluted from the membrane, by lowering the pH, and was used to coat yet a second sheet of PVDF membrane to isolate relaxin-immunoreactive peptides from the high

molecular weight, CM cellulose-flow through (acidic) sample prepared as mentioned above. The bound peptides were eluted from the PVDF membrane, again by lowering the pH, and separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) according to the method of Giulian et al.(1985) (pH 9.3; 20% T, 0.5% C_{bis}, 10% glycerol, 0.1% SDS) in a minigel system (SE 250, Hoefer Scientific Instruments, CA), and electroblotted (Western blot) onto a layer of PVDF membrane in Towbin's transfer buffer (Towbin et al.,1979) at 70 volts for 120 min. A low molecular weight calibration kit for SDS-gel (BDH Limited, Poole, England), consisting of horse heart myoglobin and it's five CNBr-cleaved fragments with molecular weights ranging from 2,512 to 16,949, was included in the SDS-gel as a molecular weight marker.

The membrane was then stained with 0.125% Coomassie blue in 10% acidic acid/50% ethanol and destained to visualize protein bands. Two major proteins bands, with apparent molecular weights (Mr) of 14,000 and 21,000, were detected (Fig. 4). To obtain sequence information about these peptides, the method of Matsudaira (1987) was employed. These protein bands were separately excised from the membrane, cut into 2 x 4 mm strips, and inserted directly into the reaction cartridge of an Applied Biosystems Model 470A gas-phase sequencer where the

Figure 4. Western blot analysis of a minigel SDS-PAGE separation of a) low molecular weight

standards (lane 2), b) porcine hemoglobin (lane 4), and c) high molecular weight, acidic, relaxin-like peptides isolated by PVDF-relaxin-immunoaffinity, Coomassie blue stained (lane 3) and immunostained (lane 1). The molecular weights of the standards in lane 2, from top down, are 16,950, 14,400, 10,600, 8,160, and 6,210, respectively. A sixth peptide (2,510 Da) was not visible in this gel system.

N-terminal amino acid sequences were determined by automated Edman degradation.

Unfortunately neither of these peptides isolated this way turned out to be relaxin-like, possibly due to nonspecific binding of proteins to the PVDF membrane. Amino acid sequencing data showed that while the 21,000 Da protein (65 pmole, as assessed by amino acid analysis) was N-terminally blocked the 14,000 Da protein band contained 105 pmole of another N-terminally blocked peptide and 15 pmole of porcine α -1 hemoglobin (Table 1).

Table 1. N-Terminal Amino Acid Sequence of an Acidic 14,000 Da Porcine Ovarian Peptide Isolated by PVDF-Relaxin-Immunoaffinity

	Cycle							
	1	5	10	15	20	25	30	35
PVDF-isolated	VLSAADKANVKAA UGKUGGQAGAUGAEALERMFLGFP							
Homology	VLSAADKANVKAA GKVGGQAGA GAEALERMFLGFP							
α -1 hemoglobin	VLSAADKANVKA AWGKVGGQAGAHGAEALERMFLGFP							

U=unidentified

Cyanogen Bromide Cleavage of 21,000 Da Peptide isolated by PVDF-Relaxin-

Immunoaffinity. Our initial interpretation about the N-terminally blocked 21,000 Da peptide was that this peptide represented the single chain prorelaxin that has a blocked amino-terminus with pyroglutamate. To obtain internal sequence information of this peptide it was necessary to subject it to enzymatic/chemical cleavage. Since prorelaxin contains methionine, *in situ* cyanogen bromide (CNBr) cleavage method of Scott et al. (1988) was used in an attempt to cleave the peptide into four fragments. Briefly, the stained band was excised from the PVDF membrane, cut into small strips (2 x 4 mm) and placed into a brown 1.5 ml microcentrifuge tube. To the tube were added 100 μ l 70% formic acid and 10 μ l 0.5 M CNBr (Aldrich Chemical, Milwaukee,

WD). The tube was immediately flushed with a stream of N₂, wrapped in aluminum foil and incubated at room temperature for 24 h with intermittent shaking. The membrane was subsequently separated from the reaction solution, washed with 100 µl water and the wash was added to the sample in the reaction solution. Peptides were extracted from the membrane with 100 µl 70% isopropyl alcohol/5% trifluoroacetic acid and added to the sample. Fifty microliters SDS-PAGE sample buffer (0.125 M Tris, 2% SDS, 20% glycerol), 2 µl bromophenol blue (0.5% w/v in 10% EtOH) and 1 ml water were added to the sample before the sample was lyophilized to dryness. The sample was lyophilized five times, with 100 µl water each time, to remove the acid completely. The resultant CNBr-cleaved peptide fragments were separated on a minigel SDS-PAGE and electroblotted (70 volts, 60 min) onto PVDF for sequencing. However, no cleavage of the 21,000 Da peptide by CNBr was detected on the PVDF membrane. Later through the use of an immunostaining technique (Janssen Biotech, 1984), in which the Western blots were first incubated with rabbit antiserum against porcine relaxin (1°Ab) followed by a gold conjugated goat anti-rabbit IgG antiserum (2°Ab, AuroProbe BLplus, Janssen Biotech) and silver-enhanced (IntenSE BL, Janssen Biotech) to identify the relaxin-like peptides, it was found that this 21,000 Da peptide was in fact not relaxin-immunoreactive at all (Fig. 4). Although the identities of these peptides were disappointing, importantly the technology for isolating and microsequencing a peptide was established in our laboratory in collaboration with the Protein Chemistry Core Facility.

Isolation of Prorelaxin via Two-Dimensional SDS Gel Electrophoresis. When it became apparent that not all peptides isolated by the PVDF-immunoaffinity method were immunoreactive, two-dimensional polyacrylamide gel electrophoresis (2-D PAGE), a modified method of O'Farrell (1975), was employed. The proteins were first separated on the basis of their net charges by isoelectric focusing (IEF) in a 4% T, 5.4% C_{bis} tube gel (2.5 x 130 mm) with 2% pH 3.5-10

Ampholine (Pharmacia LKB Biotechnology, Bromma, Sweden), then, in the second dimension, separated according to size in a standard (14 x 16 cm) slab SDS-PAGE gel (Giulian et al., 1985).

This technique has extraordinary resolving power and the protein species were separated in highly reproducible patterns. The proteins separated in 2-D gels were then electrophoretically transferred onto PVDF membranes, stained with Coomassie blue to reveal the patterns of protein separation. Some blots were immunostained to identify the relaxin-immunoreactive peptides. From the 6,000 Da relaxin-free, high molecular weight, CM cellulose-flow through of porcine ovarian extracts (see previous section for detail) a total of three acidic, high Mr, relaxin-like peptides were identified on the Western blots (Fig. 5). Unlike porcine relaxin, which has a basic

Figure 5. Western blot (PVDF) analysis of 2-D gel separation of the high molecular weight, CM

cellulose-flow through (acidic) sample from porcine ovarian extract.

4a). Coomassie blue stained blot showed the protein separation pattern obtained by 2-D gel electrophoresis.

4b). When a blot was immunologically stained, with rabbit antiserum against porcine relaxin, three intense spots (numbered 1-3) were revealed. Spot No. 3 migrated faster across the membrane hence it appeared hollow in the front side (the side contacted the gel) but solid on the back side of the membrane. The apparent molecular weights and pIs for immunoreactive peptides No. 1, 2, and 3 are 21,000, 14,000, and 14,000, and pH 6.4, 6.6, and 6.0, respectively.

pI of 10 and a molecular weight of 6,000, these relaxin-immunoreactive peptides were focused at pH 6.4, 6.6 and 6.0 with their apparent Mr estimated to be 21,000, 14,000 and 14,000, respectively. These peptides are believed to be prorelaxin and processed intermediates, respectively.

The amount of relaxin-like peptides found in a standard analytical 2-D gel were estimated to be in the low range of abundance, typically 100 ng, corresponding to about 5 pmole for a 20,000 Da peptide, from a 1 mg total protein load. This represents an upper limit of the sample load for a single standard size 2-D gel without severely compromising the resolution in either dimension.

Transfer of Prorelaxin onto PVDF. In order to increase the amount of protein separated by 2-D gels for sequence analysis, we applied multiples of the first dimension IEF gel to the slab SDS-gel. The segments of the IEF gels with pH range of interest (pH 5.5-6.8, known from preliminary analytical experiments), were cut out from each IEF gel and multiple (usually four) such gel segments were loaded across the top of a single second-dimension slab SDS-gel. Afterwards the proteins resolved in the gel were electroblotted onto PVDF membrane and revealed by Coomassie blue staining. A repeating pattern of protein separation was obtained. Multiple (six) copies of each spot were pooled, inserted directly into the reaction cartridge of the gas-phase sequencer for sequence analysis.

Results of amino acid sequencing (Table 2) of the three high molecular weight, acidic, porcine relaxin-related peptides, isolated by 2-D polyacrylamide gel electrophoresis, revealed that they all have an identical single amino acid sequence -- that of A-chain of porcine relaxin, with microheterogeneity at the N-termini. This for the first time confirms the processed forms of prorelaxin.

Table 2. N-terminal Amino Acid Sequence of Acidic, High Molecular Weight
Relaxin-Immunoreactive Peptides Isolated by 2-D Gels

	1	5	10	15	20
A-Chain of 6,000 Da Relaxin	FRMTLSEKCCQVGCIRKDIARLC				
21,000 Da Relaxin-Like Peptide	FRMTLSEK ??QVG ?I ?				
14,000 Da, pI 6.6, Relaxin-Like Peptide	FRMTLSEK ??QVG ?IRKDIARL ?				
14,000 Da, pI 6.0, Relaxin-Like Peptide	FRMTLS- - - - - - - - - -DIA ?L?				

? = Unidentified amino acid residues.

- - = Amino acid residues unassigned due to printer error causing data system shutdown.

The absence of extraneous contaminants demonstrated the high resolving power of 2-D polyacrylamide gel electrophoresis, coupled with Western Blotting, in obtaining peptide(s) pure enough for amino acid sequencing. More importantly, it confirms that these peptides truly are porcine relaxin related. Since only the amino acid sequence of A-chain was obtained, it suggests as expected that all of their B-chains are amino-terminally blocked as in the case of 6,000 Da porcine relaxin with pyroglutamate. In addition, apparently the post-translational modification that cleaves the peptide bond between the N-terminus of A-chain and the residue before it (i.e., the C-terminal amino acid residue of "C-peptide") occurred by the time they were isolated.

Since no sequence information was gained on the B-chain from any of these peptides, it is not possible to verify the presence (or absence) of all or a fragment of the C-peptide. Amino acid analysis would reveal the amino acid composition which would assist in verifying the molecular weights of these peptides. However, this approach is impractical due to the low sensitivity of amino acid analysis requiring significantly higher quantities of protein (2 - 5 µg or ~500 pmole for a 20,000 Da peptide). This compares to 10-20 pmole necessary for the amino acid sequence

determination.

The N-terminal of B-chain was not accessible to Edman degradation. Consequently, to verify the presence (or absence) of the C-peptide it became necessary to obtain an internal sequence of the B-chain by sequencing peptide fragments generated through cleavage of the polypeptide chain by chemical and/or enzymatic digestion. Due to the expected sample loss during each sample manipulation, it is necessary to start with at least 200 pmole of each peptide to ensure sufficient quantity to carry through *in situ* chemical/enzymatic cleavage and sequence analysis. However, the sequence data showed that the amount of peptides of interest are much lower than expected. As previously described, each sequenced sample (six copies of each spot) was isolated by 2-D gel from a total of 6 mg of sample and the sequencing data showed the presence of only 10 - 35 pmole of each purified peptide. This required an alternative approach since this method would require 120 spots of 21,000 Da relaxin-like peptide to obtain an internal amino acid sequence of B-chain.

In an effort to enrich relaxin-like peptides and to reduce the number of 2-D gel runs, experiments were performed with the following procedural modifications:

a.) The acidic, heat treated, high molecular weight sample, was purified one step further by isoelectric precipitating at pH 7.5, and only the supernatant was applied onto the first-dimensional IEF gels.

b.) IEF gel tubes of wider diameter (5 mm instead of 2.5 mm) were used which provides four times the cross sectional area, hence four times the sample loading limit.

c.) Shorter and wider IEF gels with pH gradient of narrower range (5 x 50 mm, pH 4 - 7 gels vs. the 2.5 x 135 mm, pH 4 - 8.8 gels) were used to prevent basic peptides from entering the gel thereby reducing their interference in the

separation of acidic peptides.

d.) Sample was included in the IEF acrylamide monomers gel solution during polymerization in an attempt to achieve increasing sample loading limit without compromising resolution of proteins.

None of these changes significantly improved the recovery of peptides of interest. The current approach is the accumulation of purified relaxin-like peptides utilizing the 2-D gel protocol where there is 1 mg sample per IEF gel followed by four relevant IEF gel pieces per second-dimensional SDS-gel.

Internal Amino Acid Sequencing of Prorelaxin. As mentioned earlier, because prorelaxin has a blocked amino terminus, it must be treated with protease or cleavage reagents to generate peptide fragments to obtain internal sequence information with which to verify the presence of the C-peptide in the high molecular weight, relaxin-like peptides isolated by 2-D gel. Due to the limited quantities, various *in situ* enzymatic/chemical cleavage methods for PVDF immobilized proteins (Kennedy et al., 1988; Walsh et al., 1988; Coli et al., 1989; Fimmel et al., 1989) have been studied. Chemical cleavage methods for PVDF bound proteins have been chosen because of the high residue-specificity of chemical reagents, the volatility of the solvents used in their reactions, and the typically larger fragments released from chemical cleavages than from a proteolytic digestion, which are better suited for separation by electrophoresis.

Since prorelaxin contains only four methionine (with two of them next to each other, i.e., Met-Met) one of the chemical reagents considered is cyanogen bromide (CNBr) to cleave the methionine-X peptide bonds, as described in the previous section. The other chemical reagent considered is BNPS-Skatole (Fontana and Gross, 1986) which cleaves specifically at the tryptophan-X peptide bonds. Prorelaxin contains only two tryptophan residues and both are located in B-chain proximal to the targeted C-peptide. The low number of this residue, the close

proximity of these residues to the C-peptide, and the availability of a tryptophan-specific reagent (BNPS-Skatole, Sigma Chemical, St. Louis, MO) make tryptophan one of the choices of site for cleavage. The Coomassie-stained spots of interest will be excised from the PVDF blot, pooled and cleaved *in situ* by BNPS-Skatole.

Briefly, the spots will be cut into small strips and placed into a 1.5 ml brown microcentrifuge tube. To the tube will be added 50 μ l water and 150 μ l skatole solution (1.3 mg/ml in glacial acetic acid) and it will be incubated for 2 h at 47 °C in the dark. The membrane will be separated from the incubation solution, washed five times with water and the peptides eluted from the membrane with 200 μ l 70% isopropanol/5% trifluoroacetic acid for 2 h. To the eluent will be added 100 μ l water, 50 μ l SDS-gel sample buffer (0.125 M Tris, 2% SDS, 20% Glycerol, pH 6.8), and 2 μ l bromophenol blue (0.5% w/v in 10% EtOH) and the sample lyophilized to remove isopropanol and trifluoroacetic acid. To recover peptides from the acetic acid incubation solution, two volumes water will be added to the sample to precipitate skatole which can be removed by centrifugation in a microcentrifuge (Eppendorf centrifuge 5415). The peptide-containing supernatant will be lyophilized and combined with the membrane wash and continued lyophilization with water (100 μ l each) for five times, or until solution is blue. Final volume will be adjusted to 100 μ l with water. The resulting chemical-cleavage peptide fragments will be separated by a 1-D slab SDS-gel, electroblotted onto a PVDF membrane and stained with Coomassie blue. The peptide-containing bands will subsequently be excised, inserted into the reaction cartridge of a gas-phase sequencer for amino acid sequence determination.

Conclusions. Three acidic, high molecular weight, relaxin-related peptides (porcine prorelaxin and intermediate peptides) have been isolated from sow ovarian extracts. They are focused at pH 6.4, 6.6 and 6.0 with estimated molecular weights of 21,000, 14,000, and 14,000, respectively. Two-dimensional polyacrylamide gel electrophoresis, in conjunction with Western

blotting of 2-D gel separated peptides onto PVDF membrane, have proved to be a powerful technique in microsequencing, especially when dealing with limited quantities of proteins.

We are in the process of determining the internal sequences of the extended B-Chain (the fragment of C-peptide connected to B-chain) to gain insight into the cleavage sites and mechanisms in the processing of porcine prorelaxin to relaxin. The progress of this study has been slowed by the limited amounts of prorelaxin present in ovarian tissues as well as the difficulty of isolating the prohormone without modification.

2. Extraction of Relaxin From Bovine Corpora Lutea and Placenta

Relaxin, a hormone of pregnancy, has been isolated from the ovary of the pig, rat, sand tiger shark, dogfish shark and skate (Sherwood, 1988). The amino acid sequence of these relaxins have been determined directly while the sequence of human relaxin has been determined via isolation of the gene. The relaxin molecule is characteristically basic, has a relative molecular mass (M_r) of 6,000, and consists of two covalently linked peptide chains. Amino acid sequence homology between all known sequences is only about 20%. However, the predominant feature of all relaxins is the insulin-like structure with six precisely placed cysteine residues that provide two interchain disulfide bonds and a single intrachain link.

Evidence for a bovine relaxin aside from extraction from tissue (Fields et al., 1980) and plasma (Wada and Yuhara, 1960, 1961; Castro-Hernandez, 1975) are a) its immunohistochemical localization in the large luteal cell (Fields et al., 1980); b) the relaxed pelvic ligaments and dilation of the cervix at the time of parturition (Perezgrovas and Anderson, 1982; Musah et al., 1986a,b), characteristics classically associated with relaxin in other species attributed to the remodeling of collagen (Sherwood, 1988); c) the detection of immunoactive relaxin in plasma of the cow approaching parturition (Anderson et al., 1982; Musah et al., 1987; Randel et al., 1988); and d) the induction of parturition with the administration of an exogenous relaxin (Perezgrovas and Anderson, 1982; Musah et al., 1986a,b).

Relaxin Isolated From Bovine Corpora Lutea. Relaxin was isolated from the bovine CL utilizing extraction and chromatographic procedures that have been successfully used for the isolation of relaxin from the other species. Corpora lutea were collected at the abattoir from late pregnant cows and immediately placed in liquid nitrogen and transported to the laboratory where they were stored at -70°C . Luteal tissue (1 kg) was extracted by modifications of the procedures described by Griss et al. (1967) and Fields et al. (1980). Briefly, frozen tissue was homogenized at 4°C at a ratio

of 1 g tissue to 6 ml extraction solution [0.7 N HCl/acetone (3:5, vol:vol)] with a Brinkman Polytron homogenizer and incubated with stirring at 4 °C for 24 h. Insoluble material was removed by centrifugation at 27,000 x g for 12 min at 4 °C and the proteins in the supernatant were precipitated with the addition of five volumes of cold acetone (4 °C). The relaxin containing acetone precipitate was collected by centrifugation at 17,000 x g for 20 min at 4 °C. The pellets were washed with acetone, and isoelectric precipitated with a total of 100 ml 0.2 M ammonium acetate (pH 5.0).

Size fractionation of the sample was performed on a Sephadex G-50 fine column (5 x 150 cm) at 4 °C. The column was pre-equilibrated and eluted with 10 mM ammonium acetate (pH 5.0) at a flow rate of 60 ml/h. Effluent were collected in 10 ml fractions and the elution profile was determined by measuring uv absorbance at 220, 280 and 415 nm.

The combined fractions 175 through 230 from Sephadex G50 chromatography was loaded onto a 4 g CM-cellulose column and eluted with a linear NaCl gradient (0-0.3 M NaCl in 10 mM ammonium acetate, pH 5.0) at the flow rate of 20 ml/h. Five ml fractions were collected and uv absorbed fractions associated with individual peaks were combined and desalted by dialysis (3M Spectrapor Membrane Tubing, Mr 3,500 cutoff) or by ultrafiltration (YM2 Filter, Mr 1,000 cutoff, Amicon Corp. Lexington, MA) against 1 mM ammonium acetate (pH 5.0). Samples were then lyophilized and stored at -20 °C.

The homologous porcine relaxin radioimmunoassay was used to measure bovine relaxin utilizing an antibody (UF-1M) developed against a CM-cellulose purified porcine relaxin (Fields and Fields, 1985). Since relaxin lacks tyrosine, relaxin was conjugated (McMurtry et al., 1978) with 3-(p-hydroxyphenyl) propionic acid N-hydroxy-succinimidyl ester (TAGIT; Calbiochem-Behring, La Jolla, CA), followed by iodination with [¹²⁵I]-Na (Amersham, Arlington Heights, IL).

The assays were carried out in 12 x 75 mm plastic tubes in 500 µl volumes consisting of 200 µl buffer (60 mM Tris, 0.15 M NaCl, 5 mM EDTA, 0.02% NaN₃, 0.2% bovine serum albumin, 0.2%

normal rabbit serum), 100 μ l of sample or standard, 100 μ l of [125 I]-relaxin and 100 μ l of relaxin antiserum. The antibody was used at a dilution which would bind 30-40% of the total 12,000 cpm added. After a 22 h incubation at 4°C, 100 μ l of goat anti-rabbit gamma-globulin (diluted 2.5x) was added to each tube. Twenty-two hours later, 1 ml of chilled buffer (minus the BSA and normal rabbit serum) was added and centrifuged at 4,000 rpm (4°C) for 30 min. The supernatant was discarded and the pellet counted in a Packard gamma-counter (model #5650) at 75% efficiency and a counting error of 2%.

Using an antibody to porcine relaxin, immunoreactive material was detected in the extracts of bovine corpora lutea from late pregnant cows (Fig. 6). Relaxin-like immunoactivity was detected in the high Mr region of the void volume and in the 6,000 Da region at the same relative position as porcine relaxin eluted from Sephadex G50. The estimated mass under this peak is about 40 μ g calculated as porcine relaxin equivalents. The presence of immunoactivity in the higher Mr region was expected since we determined in the pig the large Mr molecules with immunoactivity are incompletely processed pro-relaxin. The detected relaxin-like bioactivity appears to be associated

Figure 6. Size fractionation of bovine and porcine luteal extracts with Sephadex G-50.

with the immunoreactive material. The loss of bioactivity after reduction with dithiothreitol indicates that disulfide bonds are essential for bioactivity. Like relaxin of other species, it is also basically charged.

The 6,000 Da immunoreactive relaxin was further fractionated by cation exchange chromatography into three peaks (Fig. 7). The estimated mass in the peaks were 2, 5 and 10 μ g calculated as porcine relaxin equivalents. All four CM-cellulose isolates displaced [125 I]-labeled porcine relaxin in a dose response manner that was similar to porcine relaxin (Fig. 8). The highest cross-reactivity was consistently associated with CMC-2 at levels of 0.01 - 0.1% the immunoreactivity of porcine relaxin.

Although the major immunoreactive peak co-eluted with the major peptide component (CMC-2), we were able to demonstrate that the immunoreactivity could be separated from CMC-2 on a G-50 (Superfine) column (Fig. 9). Chemical characterization via SDS-gel electrophoresis, RP/HPLC and amino acid analysis of the CM-cellulose peptides confirmed that they did not contain two peptide

Figure 7. Ion exchange chromatography of Sephadex G-50 immunoreactive components.

Figure 8. Cross-reactivity of bovine luteal CM-cellulose fractions with porcine relaxin antiserum.

Figure 9. Separation of relaxin immunoreactivity from CMC-2 on a Sephadex G-50 superfine column.

chains with the interchain disulfide bonds. It is the conservation of two peptide chains with six cysteine residues that is characteristic of all species of relaxin.

Evidence that the detected immunoactivity is not the result of contamination by porcine relaxin is the difference in elution time of these two relaxins from a CM-cellulose column (Fig. 10). To guard against contamination, bovine relaxin is extracted and purified in a different laboratory with glassware not used in the extraction and purification of porcine relaxin.

Figure 10. Ion exchange chromatography of bovine and porcine relaxins.

Conclusion. A formidable obstacle to isolation and purification of bovine luteal relaxin is a single cow has a corpus luteum of the ovary that weighs only 5 g for extraction of 5-20 ng relaxin and only the corpora lutea of late pregnant cows contain relaxin. However, a quantitative determination of the true crossreactivity of this relaxin could not be ascertained since the major portion of the tested bovine preparation was a basically charged non-relaxin protein. In addition at these levels, bovine relaxin in purified form was found to become irreversibly bound to the containing vessel. Attempts to enhance the extractable yield of bovine relaxin from the corpus luteum at these low levels were unsuccessful.

Attempts to Enhance the Recovery of Bovine Relaxin. The recovery of low concentrations of relaxin raised the possibility of proteolytic degradation of the relaxin. In an attempt to prevent degradation during tissue storage tissue samples were frozen in liquid nitrogen immediately upon collection. Unexpectedly, short term or long term storage of tissues in liquid nitrogen resulted in a markedly decreased recovery of relaxin. A possible explanation is the fracture of lysosomal granules may have exposed the relaxin to greater proteolytic degradation upon tissue thawing. Recovery from tissues with high concentrations of relaxin (such as corpus luteum from day 60 pregnant sow) were not affected by freezing. This difference can be explained by the stoichiometric relationship between the relaxin and putative protease(s). When the efficiency of the extraction procedure was examined, no evidence of proteolysis in the extraction process was found since an average recovery of 104% was obtained when luteal tissue (2-3 g) was spiked with 39 μ g of porcine relaxin immediately prior to extraction with acid/acetone.

An alternative considered for increasing the recovery of relaxin is the prorelaxin form may predominate and is processed to the native relaxin under only the optimal physiological conditions. We have previously shown this to be the case for luteal oxytocin. When tissue extracts were passed through Sephadex G50, relaxin immunoactivity profile of the eluates indicated significant activity eluted at a volume close to the void volume (Fig. 6). A prorelaxin with a Mr of 17,000 to 20,000 as previously shown for rat, human and pig, would elute at this region of detected immunoactivity. Thus, the immunoactivity associated with the high Mr is a good candidate for bovine prorelaxin. In addition we have shown that limited tryptic digestion of the porcine high Mr immunoreactive relaxin resulted in its conversion to 6,000 Da relaxin. We have determined the isoelectric point of porcine prorelaxin is 5.4. This explains that when the crude extract (acetone precipitate) was isoelectric precipitated at pH 7.6 there was a ten times higher yield of porcine 6,000 Da relaxin than when isoelectric precipitation was conducted at pH 5.0. The latter is the standard PI used for purification

of 6,000 Da relaxin. However, in the case of bovine relaxin from corpora lutea there was no difference in the yield whether the isoelectric precipitation was carried out at pH 7.6 or 5.0.

Conclusion. In exploring a number of possibilities, we were unsuccessful in enhancing the recovery of bovine luteal relaxin over that seen with classical extraction techniques that have been successfully used to recover relaxin from tissues of other species. Relaxin in bovine tissues is at a low concentration in contrast to that found in the pig, rat and horse.

SDS Polyacrylamide Gel Electrophoresis Separation of the α and β Chains of Porcine Relaxin.

The decrease in recovery of relaxin from tissue in stored liquid nitrogen, the finding of relaxin in the placenta of sheep (Wathes et al., 1988) and the readily available kilogram quantities of fresh placenta from late pregnant cows lead us to the evaluation of the placenta as a source of relaxin in the cow. We believed the application of some of the recent advances in protein chemistry coupled with a source of relaxin from a greater tissue mass that there was a reasonable possibility of purifying and determining the amino acid sequence of bovine relaxin. The following new techniques were developed for isolating the relaxin and separating the α and β chain before attempting amino acid sequencing.

One-dimensional SDS polyacrylamide gel electrophoresis (SDS-PAGE) was carried out according to the method of Giulian et al. (1985) in a minigel system. The gels (pH 9.3; 20% acrylamide, 10% glycerol, 0.1% SDS) were stained with Coomassie blue. Standards included horse heart myoglobin (MW range 2,512 to 16,949; BDH Limited, Poole, England) and prestained markers (Diversified Biotech, Newton Centre, MA) consisting of horse heart myoglobin and trypsin inhibitor (MW range 2,512 to 20,400). Although Schagger and Von Jagow (1987) reported that these marker kits contain six and seven bands, respectively, the smallest band (2,512 Da) stained poorly with Coomassie blue and was usually not detected. Samples and standards were combined with equal volumes of sample buffer (0.125 M Tris, 2% SDS, 20% glycerol) for a final concentration of 0.5

mg/ml. Porcine relaxin was reduced by incubation in sample buffer at 100 °C containing 5% β -mercaptoethanol. One microgram of each sample or standard was loaded onto the gels for electrophoresis. In an attempt to improve chain separation several modification of the protocol were made.

An SDS-PAGE gel of both unreduced and porcine relaxin reduced with β -mercaptoethanol is shown in Figure 11. The Coomassie blue stained gel indicated a single band for the unreduced relaxin (lane c) and a single band for the reduced relaxin (lane d). The reduced relaxin migrated faster and the band was more diffuse. The 2,512 Da band reportedly contained in the horse heart myoglobin standards was not detected. Considering the reduction of relaxin was incomplete, the protocol for reduction of porcine relaxin was changed from an incubation of 90 s at 100 °C in 5% β -mercaptoethanol to an incubation of 180 s and sample buffer that contained 10% β -mercaptoethanol.

Figure 11. A one-dimensional SDS-polyacrylamide gel of unreduced relaxin versus relaxin reduced with β -mercaptoethanol that was stained with Coomassie blue. Horse heart myoglobin standards MW range 6,210 to 16,949 (lane a); myoglobin + trypsin inhibitor prestained standards MW range 6,210 to 20,400 (lane b); unreduced relaxin (UF-17S; lane c); reduced relaxin (UF-17S; lane d). Note that reduced relaxin migrated further into the gel than unreduced relaxin.

No improvements were detected in the separation of the α and β chains.

An SDS-PAGE gel of both unreduced and reduced porcine relaxin in which the percentage of acrylamide was adjusted from 20% total and 0.5% bis crosslinking to the following concentrations: 23.0% total:0.5% bis, 23.0 total:2.0% bis, 20.0% total:5.0% bis, 20.0% total:1.25% bis, in order to decrease the pore size is shown in Figure 12. Two bands were detected in the lanes containing reduced relaxin, one migrating at the same rate as unreduced relaxin and one migrating faster. The band migrating with unreduced relaxin is believed to be intact relaxin that was incompletely reduced and detected when electrophoresis immediately followed addition of β -mercaptoethanol. There was a time dependent loss in the appearance of this band after reduction with mercaptoethanol (Fig. 11). Separation of the peptides of the standards was improved but only a single band that migrated faster than unreduced relaxin could be detected for reduced relaxin. Additional alterations of the acrylamide concentration produced similar results.

Figure 12. A one-dimensional SDS-polyacrylamide gel of unreduced relaxin versus relaxin reduced with β -mercaptoethanol that was stained with Coomassie blue in a gel containing 23% total acrylamide instead of 20%. Horse heart myoglobin standards MW range 2,512 to 16,949 (lanes a and f); unreduced relaxin (UF-17S; lanes b and e); reduced relaxin (UF-17S; lanes c and d). A slight improvement in the separation of the standards was detected but the two chains in the reduced relaxin lane still migrated as one.

An SDS-PAGE gel of both unreduced and reduced porcine relaxin in which an attempt was made to prevent the chains from reannealing by the addition of 11 ml β -mercaptoethanol to the upper

buffer chamber is shown in Figure 13. The β -mercaptoethanol was added to the buffer since it was thought this reducing agent migrated away from porcine relaxin, allowing the reannealing of the disulfide bonds (Fig. 13). The addition of the β -mercaptoethanol had a negative effect not only on relaxin itself but also the standards as seen by the "smile" that extended across the lanes containing relaxin in addition to the lanes directly adjacent to relaxin. The occurrence of the "smile" was a result of proteins containing disulfide bonds rather than due to technical problems (Gurusinghe et al., 1986). It was reported that β -mercaptoethanol electrophoresed in lanes adjacent to proteins containing disulfide bonds resulted in convex and/or concave bands. Prerunning the gel for 20 min with sample buffer containing 5% β -mercaptoethanol and addition of 20 μ l β -mercaptoethanol to each sample well 45 min after samples entered the separating gel eliminated the "smile" effect. However, no improvement was detected in separation of the chains.

Figure 13. A one-dimensional SDS-polyacrylamide gel of unreduced relaxin versus relaxin reduced with β -mercaptoethanol that was stained with Coomassie blue in which 1 ml β -mercaptoethanol was added to the upper buffer chamber. Horse heart myoglobin standards MW range 6,210 to 16,949 (lanes a and e); myoglobin + trypsin inhibitor prestained standards MW range 6,210 to 20,400 (lane b); unreduced relaxin (UF-17S; lane c); reduced relaxin (UF-17S; lane d). The continued infusion of β -mercaptoethanol throughout electrophoresis caused the appearance of a "smile" which masked any potential improvement in separation of the chains.

An SDS-PAGE gel of porcine relaxin which exhibited multiple bands upon oxidation with performic acid is shown in Figure 14. The disulfide bonds of porcine relaxin were broken by oxidation with performic acid according to the method of Sanger (1949) instead of β -mercaptoethanol. Performic acid converts cystine to cysteic acid whereas β -mercaptoethanol causes formation of sulfhydryl groups which readily reoxidize in the absence of reducing agent. Comparison with unoxidized relaxin on the same gel indicated that the fastest migrating band of the oxidized sample was 6,000 Da and that the other bands were higher Mr. The absence of lower Mr bands (< 6,000 Da) indicated chain separation had not occurred. When performic acid oxidized relaxin was reduced with β -mercaptoethanol, two bands were formed, one corresponding to the 6,000 Da unreduced relaxin and the other corresponding to the low Mr band of reduced relaxin. Presence of a lower Mr

Figure 14. A one-dimensional SDS-polyacrylamide gel stained with Coomassie blue of unreduced versus performic acid oxidized relaxin. Unreduced relaxin (UF-17S; lane a); relaxin reduced with β -mercaptoethanol (UF-17S; lane b); relaxin reduced with dithiothreitol (UF-16S; lane c); oxidized relaxin (UF-17S; lane d); oxidized relaxin reduced with β -mercaptoethanol (UF-17S; lane e); horse heart myoglobin standard MW range 6,210 to 16,949 (lane f); myoglobin + trypsin inhibitor prestained standard MW range 6,210 to 20,400, (lane g). Oxidation of relaxin resulted in the formation of aggregates, despite the presence of SDS and did not improve chain separation.

band indicated breakage of the disulfide bonds but absence of two distinct lower Mr bands meant electrophoretic separation of the chains had not occurred. Lane c contains porcine relaxin treated with dithiothreitol. Dithiothreitol was originally used to reduce porcine relaxin but, as indicated, separation was minimal. β -mercaptoethanol was found to be more effective and was used in all subsequent studies on reduction of relaxin.

Since the antibody recognition site of reduced relaxin was still intact, either reduction prior to electrophoresis was incomplete or the sulfhydryl groups had reoxidized, causing the disulfide bands to reform. If reduction of relaxin were incomplete, the 6,000 Da relaxin should still be detected. This was confirmed when a 6,000 Da relaxin was detected after electrophoresis of freshly reduced samples. Electrophoresis of reduced samples stored for several days resulted in the appearance of only one band and it migrated faster than intact relaxin. This indicated complete reduction had taken place when incubation occurred over a longer period while in storage.

Since there was evidence indicating that complete reduction had taken place, it was thought that reoxidation of the chains was occurring either during or after electrophoresis. Lundblad and Noyes (1984) reported that at alkaline pH, most proteins will rapidly reoxidize to form the original disulfide bonds. Reoxidation of the relaxin chains was thus considered since the pH of the separating gels used in these experiments was approximately 9.3. However, reoxidation of the chains before electrophoresis would be unlikely because the sample buffer has a neutral pH 6.8.

Schagger and von Jagow (1987) developed a tricine-SDS gel electrophoresis system reported to improve separation of low Mr peptides. However, this system was also unable to resolve the A and B chains of porcine relaxin (Dr. R. Renegar, unpublished results). Like the method of Giulian et al. (1985), the pH 8.9 of the separating gel in the tricine-SDS system is basic, an environment that can lead to reannealing of the disulfide bonds.

One-Dimensional Isoelectric Focusing to Separate the A and B Chain of Relaxin. Since

previous experiments with SDS gels did not result in chain separation suitable for sequencing, separation based on differences in isoelectric points was investigated. One-dimensional denaturing isoelectric focusing was performed in the vertical slab minigel system according to the method of Robertson et al. (1987). The range of ampholytes (Pharmacia LKB Biotechnology, Bromma, Sweden) used was either pH 3.5 to 10.0 or pH 3.5 to 10.0 (10%) and pH 9 to 11 (90%). Approximately 1 to 3 μ g of each standard and 5 μ g of porcine relaxin were loaded and electrophoresed.

In later trials, some of the protein bands were excised from the gel. The gel slices were incubated overnight in sample buffer (0.125 M Tris, 2% SDS, 20% glycerol) and then placed in the wells of a one-dimensional polyacrylamide gel and electrophoresed as described earlier according to the original protocol of Giulian et al. (1985). Following electrophoresis, the gels were either stained with Coomassie blue or electroblotted for 30 min according to Towbin et al. (1979) onto polyvinylidene difluoride membranes (PVDF) (Immobilon-P, Millipore Inc., Bedford, MA) and visualized by immunostaining (Auroprobe BLplus GAR, Amersham Corporation, Arlington Heights, IL) and silver enhancement (IntenSE BL, Amersham Corporation).

The disulfide bonds of porcine relaxin were broken by incubation with β -mercaptoethanol and the resultant chains were separated by isoelectric focusing. Two bands were detected by Coomassie blue staining (Fig. 15) that migrated differently from those seen with unreduced relaxin. The change in pI of the bands caused by reduction with mercaptoethanol indicated that all disulfide bonds were broken and that separation of the chains may have occurred. If reduction had been incomplete, i.e., some of the disulfide bonds were still intact, the pI of the resultant peptides should have remained the same as that for unreduced relaxin since there would have been no additions or deletions of amino acids. Although the pH gradient of the gel was determined, the standards did not migrate according to their pI, but as the pI of the standards became more acidic or basic the calculated pI differed more from the known pI. This prevented determination of the pI of the two bands from the reduced relaxin

Figure 15. Slab isoelectric gel stained with Coomassie blue of unreduced relaxin versus relaxin reduced with β -mercaptoethanol. Hemoglobin (lane a); unreduced relaxin (UF-19S; lane b); reduced relaxin (UF-19S; lane c); chymotrypsinogen (lane d); trypsinogen (lane e); cytochrome C (lane f). The reduced relaxin migrated as two bands, one migrating near cytochrome C and the other near hemoglobin.

lane.

Numerous faint Coomassie blue stained bands were found in addition to a heavily stained band in the lane containing intact relaxin on the isoelectric focusing gel (Fig. 15) that have not been previously reported. These faint bands are thought to be microheterogeneous forms of relaxin but could be contaminating proteins. If the faint bands were contaminating proteins, it would be unlikely that they all contained disulfide bonds. Those proteins not containing disulfide bonds should have migrated in a similar fashion in both the reduced and unreduced relaxin lanes, yet neither of the two bands in the reduced relaxin lane correlated with any of the faint bands.

The two putative A and B protein bands were excised from the slab isoelectric gel, re-electrophoresed on a one-dimensional SDS polyacrylamide gel and electroblotted onto a PVDF membrane in an attempt to confirm the identity of the two bands as the α and β chains of relaxin. Neither of the two bands could be detected by immunostaining (Fig. 16), whereas unreduced relaxin electrophoresed in the same manner was detected by the antibody. If true chain separation had

Figure 16. A western blot of unreduced versus β -mercaptoethanol reduced relaxins immunostained and enhanced with silver staining. Unreduced relaxin electrophoresed only on the SDS gel (UF-17S; lane a); reduced relaxin electrophoresed only on the SDS gel (UF-17S; lane b); cytochrome C excised from the slab isoelectric gel and re-electrophoresed (lane c); unreduced relaxin excised from the slab isoelectric gel and re-electrophoresed (UF-19S; lane d); the two bands of reduced relaxin excised from the slab isoelectric gel and re-electrophoresed (UF-19S; lanes e and f). Neither of the two bands of reduced relaxin that were re-electrophoresed could be detected indicating true separation of the chains may have occurred. Separation of the chains should disrupt proper chain folding thereby interfering with subsequent immunodetection.

occurred, then the binding site for the antibody supposedly would have been destroyed, resulting in a lack of affinity by the antibody for the two chains. Detection of the individual chains by an antibody produced against intact relaxin has not been reported. While it is possible excision of the bands from the isoelectric gel was faulty, the experiment was repeated several times with similar results. If reduction were incomplete or if some reannealing of the chains had occurred, then the resultant bands should have been detected by immunostaining as is the case in one-dimensional SDS gel electrophoresis. It is also possible that the two bands for reduced relaxin were not detected by immunostaining because sample losses were too great. However, this seems unlikely since unreduced relaxin was always detectable when similar quantities were used.

Positive identification of the two bands detected in the slab isoelectric gel as the α and β

Figure 17. A one-dimensional SDS-polyacrylamide gel of unreduced relaxin versus relaxin reduced with β -mercaptoethanol that was stained with Coomassie blue. Horse heart myoglobin standard MW range 6,210 to 16,949 (lane a); myoglobin + trypsin inhibitor prestained standards MW range 6,210 to 20,400 (lane b); unreduced relaxin electrophoresed only on the SDS gel (UF-17S; lane c); reduced relaxin electrophoresed only on the SDS gel (UF-17S; lane d); unreduced relaxin excised from the slab isoelectric gel and re-electrophoresed (UF-19S; lane e); reduced relaxin excised from the slab isoelectric gel and re-electrophoresed (UF-19S; upper band, lane f, lower band, lane g); cytochrome C excised from the slab isoelectric gel and re-electrophoresed (lane h). The cause of the smearing was not determined.

chains of porcine relaxin could only be obtained by amino acid sequencing. Therefore, the gels were stained with Coomassie blue instead of being electroblotted and immunostained in order to determine where the two bands were migrating. Unfortunately, both bands were heavily smeared (Fig. 17). However, there appeared to be a wide separation between the two bands with one migrating near the dye front, whereas the other migrated closer to unreduced relaxin. Unreduced relaxin electrophoresed in a similar manner was also heavily smeared although a band corresponding to 6,000 Da could be distinguished within the smear. Cytochrome C treated in the same manner appeared to have little smearing. Presence of contaminating proteins was not likely because electrophoresis of the same relaxin preparation on one-dimensional SDS gels resulted in the appearance of only one band when stained with Coomassie blue.

A more likely possibility for the smearing was inadequate transfer of relaxin from the isoelectric gel slices into the SDS gel. Placement of the gel slices into the wells of the SDS gels was a difficult task. In the case of cytochrome C gel slices were smaller and easier to manipulate. The gel slices for relaxins tended to be bigger to ensure that excision of the protein band was complete

since they were located by their orientation to cytochrome C from previous gels. It was concluded that further work needed to be conducted to refine the transfer of the cut gel to the SDS slab gel to preempt the smearing.

Two-dimensional Polyacrylamide Gel Electrophoresis Separation of A and B Chain of Porcine Relaxin. The use of conventional two-dimensional PAGE, which uses a tube gel for isoelectric focusing, was investigated in an attempt to eliminate the smearing. The procedure was modified for non-equilibrium pH-gradient electrophoresis (NEPHGE) according to Pollard (1984) for the first dimension using the minigel system. A slight modification was made by changing the pH range from 5 to 7 to pH 9 to 11 in order to accommodate the basic nature of relaxin. One microgram of either unreduced relaxin or relaxin that had been reduced with β -mercaptoethanol was loaded into the tube gels and electrophoresed. The second dimension was run as described above for SDS-PAGE except the slab gels were overlayed with 1% agarose in sample buffer (0.125 M Tris, 2% SDS, 20% glycerol) to adhere the tube gel to the second dimension. Porcine relaxin was reduced by incubation in 5% β -mercaptoethanol at 100 °C for 90 s. Following electrophoresis, the gels were blotted according to Towbin et al. (1979) onto PVDF membranes and visualized by immunostaining and silver enhancement.

A Western blot of unreduced relaxin that has been immunostained with a polyclonal antibody against relaxin, followed by silver enhancement is shown in Figure 18. The relaxin band in lane a was electrophoresed in both dimensions, whereas the relaxin band in lane b was electrophoresed in the second dimension only. Electrophoresis of unreduced relaxin resulted in a band migrating towards the basic end of the isoelectric tube gel in the first dimension and migrating with unreduced relaxin applied only to the second dimension (Fig. 18). A Western blot of reduced relaxin that has been immunostained with a polyclonal antibody against relaxin, followed by silver enhancement is shown in Figure 19. The resultant lengthwise smear indicated that reduced relaxin was not being focused

Figure 18. Western blot of unreduced relaxin from two-dimensional PAGE onto a PVDF membrane that has been immunostained followed by silver enhancement. Unreduced relaxin electrophoresed in both dimensions (UF-19S; lane a); unreduced relaxin electrophoresed in the second dimension only (UF-17S; lane b). The unreduced relaxin in lane 1 migrated toward the basic end of the gel in the first dimension and migrated similar to the relaxin in lane 2 in the second dimension. The fainter band is believed to result from aggregation of 6,000 D relaxin.

Figure 19. Western blot of β -mercaptoethanol reduced relaxin (UF-19S) from two-dimensional PAGE onto a PVDF membrane that has been immunostained followed by silver enhancement. Smearing of the protein widthwise in the gel indicates that it was not focused in the first dimension. This smearing is believed to be a result of the reduction process since unreduced relaxin (Fig. 10) migrated as expected.

in the first dimension. This behavior of reduced relaxin was puzzling since it did focus when electrophoresed on slab isoelectric gels. Unreduced relaxin focused in both systems as expected. When the slab gel protocol was adapted for the first dimension tube gel, reduced relaxin still did not focus. Although slight separation of the A and B chains was detected in several instances, separation was not reproducible nor sufficiently adequate for purification of each chain. Numerous efforts to improve chain separation by slight changes in protocol were attempted but did not result in better resolution. However, reduced relaxin did migrate faster than unreduced relaxin and was readily stained by the relaxin antibody when electroblotted.

Conclusion. Failure to separate the chains before sequencing would cause detection of two amino acids with every cycle instead of one. Under these circumstances, it is usually impossible to identify which chain the amino acid belongs to. However, sequence analysis of other relaxins has revealed strict conservation of several residues, including the cysteines responsible for disulfide bonds (Sherwood, 1988). Based on this information, direct sequencing of the unseparated chains of bovine immunorelaxin should still provide evidence the protein in question is relaxin. For example, in all relaxins that have been sequenced, glycine is present at the tenth position of the A chain followed by a cysteine. Therefore, a glycine should be one of the two amino acids detected near the tenth cycle. Since cysteines appear as blanks when sequenced, the eleventh cycle should have only one amino acid. In addition, the B chain is longer than the A chain by at least three residues meaning the last cycles of the sequencer would detect only one amino acid.

A problem frequently encountered in sequence determination of relaxin is blockage by a pyroglutamyl residue on the N-terminus of either chain prevents sequence analysis of that chain. The problem was first identified with the B chain of porcine relaxin by Schwabe and McDonald (1977) and has since been encountered in the B chain of shark (Gowan et al., 1981) and whale relaxins (Schwabe et al., 1989), and with the A chain of rat relaxin (John et al., 1981). In the case of a

relaxin with unseparated chains, this can be advantageous since only one amino acid would be detected with each cycle, thereby producing a sequence for one of the two chains. Identification of the chain sequenced could be determined by comparing the number and location of cysteine (absence) and glycine residues with other relaxins.

Isolation of Relaxin from the Bovine Placenta. Recently, Wathes et al. (1988) reported isolation and purification of relaxin from the placenta of another ruminant, the ewe. This report was of great interest because previous reports had indicated very low concentrations of relaxin in sheep Renegar and Larkin (1985). Since placentomes, endometrium, allantochorion and whole ovaries all contained low concentrations of relaxin immunoreactivity, Wathes et al. (1988) concluded that, as the largest tissue component by weight, placentomes were the major source of relaxin-like peptides in the ewe. With low concentrations of relaxin-like peptides existing in the cow, it was thought a similar situation might be occurring in this species.

Preliminary studies were undertaken with an acid/acetone extraction for relaxin in tissues from maternal (caruncle and intercaruncle endometrium and corpora lutea) and fetal (cotyledon and non-cotyledonary chorion) origin the last half of pregnancy. The concentration of relaxin in the placenta (cotyledon) was about 1 ng/g tissue. A 4 kg placenta (cotyledon) from a single cow yields 2-4 µg relaxin in porcine relaxin equivalents. Immunohistochemistry confirmed the localization of relaxin in the cotyledon. The only other tissue that specifically stained for relaxin was the corpus luteum from the same animal. Staining in the cotyledon was confined to the syncytiotrophoblast.

Relaxin containing placental extracts were isoelectrically precipitated at pH 5.0. Bovine relaxin was shown to be basically charged and that isoelectric precipitation at this pH removed a majority of the non-relaxin proteins. The extracts were applied to a CM-cellulose column and eluted with a stepwise gradient of NaCl, i.e., 0.05, 0.15 and 0.3 M. Following lyophilization, each fraction was desalted using Spectrapor 3 dialysis tubing. Samples were re-lyophilized and applied to gel

electrophoresis. From the electrophoresis experiments it became apparent there was insufficient material available for visualizing relaxin on an electrophoretic gel. The next step chosen was to evaluate alternative methods of extraction in an attempt to increase the yield of relaxin from the placenta.

Acid Acetone Versus Tris Extraction of Tissue. From late gestation cows placentomes were immediately excised and the cotyledonary portion of the placentomes removed, cut into smaller pieces and divided into three portions. Each portion was extracted according to one of the following procedures: acid-acetone, Tris buffer at 10 ml/g tissue, Tris buffer at 5 ml/g tissue (Table 3). All tissues were extracted on the day of collection since freezing of tissues were shown to decrease the content of relaxin.

Table 3. Cotyledons Collected and Subsequent Partitioning into Three Extraction Methods					
			Cotyledons extracted (g)		
Cow No.	Crown rump length (cm)	Cotyledons Collected (g)	Tris buffer (5 ml/g)	Tris buffer (10 ml/g)	Acid-acetone
50	50	1500	250	250	1000
84	84	1772	-	500	1250
90	90	1500	375	250	500
100	100	190	40	50	100
6555	55	1050	200	100	750
5454	56	150	100	50	-
900	90	1200	200	100	900

Up to 500 g of cotyledonary tissue were ground with 144 ml 0.75 N HCl in a food processor (Type 700, Krups, Germany). The ground tissue was homogenized (Polytron, Brinkman Instruments Inc., Westbury, NY) in an acidic solution with a final concentration of 0.5 N HCl + 85% acetone at 2.5 ml/g of tissue overnight at 4 °C (Steward and Papkoff, 1986). The tissue homogenate was subsequently centrifuged at 5,800 x g at 4 °C for 2 min and the supernatant filtered through three layers of cheese cloth after which 5.67 volumes of ice cold acetone were added to precipitate the

relaxin. The precipitate was collected by centrifugation at 3,000 x g at 4°C for 10 min, washed several times with ice cold acetone to remove the HCl, allowed to air dry and then stored at -20°C. The yield of immunoreactive relaxin averaged 79 ng (range 30 to 150 ng) relaxin immunoreactivity/kg fresh tissue.

An unpublished procedure (R Renegar, personal communication) that involved extraction with a Tris buffer containing salt and triton X-100 (detergent) as extracting agents that increased the yield of relaxin immunoreactive from hamster placenta 10 fold was investigated. The Tris buffer extraction was used at 10 ml buffer/g fresh tissue as well as at 5 ml/g tissue in an attempt to decrease the volume of extract obtained.

On the day prior to tissue collection, 150 ml EDTA (8 g/200 ml H₂O; Boehringer Mannheim Biochemicals, Indianapolis, IN) and 30 ml pepstatin (0.8 mg/40 ml ethanol; Boehringer Mannheim Biochemicals) were added to 15 l Tris buffer (pH 9.0 at 4°C; 0.15 M NaCl, 0.01 M Tris, 0.1% Triton X-100). In addition, a stock solution of phenylmethanesulfonyl fluoride (PMSF; 1.32 g/132 ml 100% ethanol; Boehringer Mannheim Biochemicals) was prepared. On the day of tissue collection, each 250 g batch of cotyledons was combined with either 2.5 l Tris buffer (10 ml/g tissue) or 1.25 l Tris buffer (5 ml/g tissue) in a 4 l beaker and either 16.5 ml PMSF (10 ml/g) or 8.25 ml PMSF (5 ml/g) were added. The tissue was immediately homogenized (Polytron, Brinkman Instruments Inc., Westbury, NY) and subsequently centrifuged at 12,000 x g at 4°C for 70 min to remove the insoluble material. The supernatants were stored at -20°C. The extracts were later concentrated to a tenth of the original volume by either ultrafiltration (Amicon, Alloy Products Corp., Waukesha, WI) using a YM1 membrane (1,000 Mr cutoff) or lyophilization and stored in 10 ml aliquots at -20°C.

Extraction of bovine cotyledons by the 10 ml/g Tris buffer procedure yielded 514 ng relaxin immunoreactivity/kg tissue, a six fold increase over the 80 ng/kg immunoreactive relaxin obtained

Table 4. Comparison of Immunoreactive (IR)-Relaxin in Crude Extracts of Cotyledons from Late Pregnant Cows.				
		Equivalents IR-relaxin (ng/kg)		
Cow Number	Crown rump length (cm)	Tris buffer 5 ml/g	Tris buffer 10 ml/g	Acid-Acetone ^b
50 ^a	50	84	574	33
6655	55	165	560	45
5454	56	184	248	-
84	84	-	745	94
90	90	570	490	120
900	90	132	422	40
100	100	164	560	150
Mean $\pm$ SE	75 $\pm$ 8	217 $\pm$ 72 ^d	514 $\pm$ 58 ^c	80 $\pm$ 20 ^d

^aCow #50 is a combination of tissue from three different cows all of approximately the same stage of gestation as determined by crown-rump length.

^bValues were obtained from the pH 5 soluble fraction since the acid-acetone crude extract is not in soluble form.

^{c,d}Means with different superscripts were different ($P < .01$).

from acid-acetone extraction of a portion of the same cotyledonary tissue (Table 4). No significant differences in yield were detected when the 5 ml/g procedure was compared to the acid-acetone method. Lack of significance was probably due to the variation encountered in these data. The yields ranged from 84 to 570 ng/kg for the 5 ml/g Tris method, 248 to 745 ng/kg for the 10 ml/g Tris method, and 40 to 150 ng/kg for the acid-acetone method (Table 4). Wathes et al. (1988) found similar variability in immunoreactive relaxin concentrations of ewe placenta but could offer no explanation for this observation.

The purification of relaxin from other species has generally involved a combination of gel filtration and either ion-exchange chromatography or isoelectric focusing. Gel filtration was selected as an initial step over ion exchange chromatography for purification since the former eliminated the steps involving salt removal. Ultrafiltration and dialysis, commonly used procedures for desalting,

are associated with sample loss, especially with increasing purity of the desired peptide (Eckerskorn and Lottspeich, 1990).

Sephadex G50. The crude extract supernatants were applied onto a 1.5 x 50 cm Sephadex G-50 column which had been pre-equilibrated with 0.02 M ammonium acetate, pH 5.0 and eluted at a flow rate of 9.5 ml/h. The resultant fractions (1 ml) were monitored for protein by measuring absorbance at 278 nm and for relaxin immunoreactivity by using the porcine relaxin radioimmunoassay previously described. Although samples were loaded at maximum capacity (10%

Figure 20. Elution profile of 1 ml fractions IR-relaxin and UV-absorbance from a bovine cotyledonary extract spiked with 50 ng CM-cellulose purified porcine relaxin (UF-19S) loaded onto a 1.5 x 50 cm Sephadex G-50 column equilibrated with 0.02 M ammonium acetate, pH 5 compared to the elution profile of the standards: bacitracin, BSA, cytochrome C, hemoglobin.

Figure 21. Elution profile of 1 ml fractions IR-relaxin and UV-absorbance from a bovine cotyledonary extract loaded onto a 1.5 x 50 cm Sephadex G-50 column equilibrated with 0.02 M ammonium acetate, pH 5.0 compared to the elution profile of the standards: bacitracin, BSA, cytochrome C, hemoglobin.

of bed volume), levels of immunoreactive relaxin in the eluates were below the detection limit of the relaxin RIA (Fig. 20 and 21).

SDS - Electrophoresis Gel Slices. Despite low levels, attempts were made to determine the Mr of the relaxin using a combination of gel electrophoresis and the relaxin RIA. Crude extracts of placentomes were loaded at maximum capacity onto SDS polyacrylamide gels and electrophoresed. Instead of using conventional staining methods, the gels were sliced and each slice was assayed in the relaxin RIA. Unfortunately, the sensitivity of this system was so great that relaxin immunoreactivity was detected even in slices from lanes in which no samples were loaded (Fig. 22). Apparently, trace amounts of porcine relaxin remained on the electrophoresis equipment from the previous chain separation experiments. After washing the equipment several times, the "background"

Figure 22. Relaxin immunoreactivity (pg/slice), as measured in the relaxin RIA, eluted from 1 mm SDS polyacrylamide gel slices with TBS containing .2% BSA and .2% NRS. The control lane contained no samples whereas the other two lanes contained the pH 5 soluble portion of the acid-acetone crude extracts of cows #1359 and #9312. No significant peaks were detected above background.

Figure 23. Relaxin immunoreactivity (ng/slice), as measured in the relaxin RIA, eluted from 4 mm SDS polyacrylamide gel slices in the relaxin lane and 2 mm gel slices in the control lane with TBS containing .2% BSA and .2% NRS. The control lane contained no samples whereas the relaxin lane contained 250 ng CM-cellulose purified porcine relaxin (UF-17S). A significant peak was detected in the relaxin lane.

immunoactivity was minimized to low levels (0 to 15 pg) and the bovine extracts were electrophoresed (Fig. 23). Thereafter increase in bovine relaxin immunoreactivity could not be detected likely the results of the estimated 40 pg of immunoreactive relaxin loaded onto the gels spread across several gel slices. At such low levels, any peaks would have been indistinguishable from background.

Parallelism of Bovine and Porcine Relaxin. Dose-response curves generated from 12.5, 25, 50, 100 or 20 μ l placental and cotyledonary acid acetone extracts were compared to the porcine relaxin standard in the relaxin RIA. Sensitivity of the assay, determined as two standard deviations above the zero dose level, was 19.92 ± 3.47 pg/assay tube (mean $\pm$ SE). Intra- and interassay coefficients of variation, determined for sow plasma from day 112 of pregnancy having 21.57 ± 0.66 ng relaxin per ml plasma, were 7.03% and 10.81% for 100 μ l aliquots of ten fold diluted day 112 sow plasma, respectively.

All comparison between bovine relaxin (both placental and cotyledonary) and porcine relaxin showed heterogeneity of regression (Fig. 24-27). Non-parallelism to porcine relaxin has been detected in another species, the rat (Sherwood and Crnekovic, 1979) in which it was concluded that, although the structure of rat relaxin was similar to porcine relaxin, it was not identical. A puzzling feature of the parallelism studies was that the slopes of the dose-response curves for bovine extracts appeared to be steeper (Fig. 24-27) than the porcine relaxin standard curve. In typical cases of non-parallelism, the slope of the dose-response curves tends to be lower than the standard curve, as is the case with rat relaxin (Sherwood and Crnekovic, 1979). Curves with lower slopes indicate the antibody has less affinity for the antigen in question than for the antigen which it was produced against. Steeper curves in these studies indicated the antibody had greater affinity for bovine relaxin than for porcine relaxin. Unfortunately, greater affinity by the antibody for bovine relaxin would result in overestimation in the RIA meaning levels in the cow may be even lower than estimated.

Figure 24. Dose-response curves for the pH 5 soluble fraction of the placental crude extracts of cows #8131 (o), #9312 (□) and #110 (v)

Figure 25. Dose-response curves for the pH 5 soluble fraction of the placental crude extracts of cows #1359 (o) and #9318 (□)

Figure 26. Dose-response curves for the pH 5 soluble fraction of the placental crude extracts of cows #1359 (o), #9312 (v), #9318 (Δ) and #110 (□)

Figure 27. Dose-response curves for the pH 5 soluble fraction of the cotyledonary crude extracts of cows #84 (o) and #90 (v)

When samples were assayed within a single assay differences were not detected between the slopes of the dose-response curves for the cotyledonary extracts nor between the combined curve for cotyledons and the combined curve for the placentomes. In a few instances where non-parallelism was noted the use of crude extracts could have caused the non-parallelism between the individual extracts. Interference by other proteins seems especially evident since wide variations were presented

within each set of triplicates whereas measurements of the standard and the intra- and interassay control (sow plasma) were more uniform indicating the variation was not due to pipetting errors.

Recovery of relaxin from placentomal crude extracts spiked with 16, 31, 63, 125, 250, 500, 1,000 2,000 or 4,000 pg porcine relaxin ranged from 31 to 155% (Table 5, Fig. 28). The control curve represents recovery of porcine relaxin in the absence of placentomal crude extract. The control

Table 5. Percent Recovery of Porcine Relaxin from Crude Extracts of Placentomes.						
Relaxin (pg)		Relaxin recovery (%)				
added	recovered mean $\pm$ SE	#8131	#9312	#9318	#110	#1359
15.6	15 $\pm$ 4	118	72	-	31	155
31.2	25 $\pm$ 5	66	139	74	73	54
62.5	51 $\pm$ 5	78	83	56	84	107
125	101 $\pm$ 4	73	89	76	79	86
250	225 $\pm$ 8	94	84	82	91	98
500	450 $\pm$ 24	87	94	77	86	106
1000	904 $\pm$ 29	93	96	82	85	96
2000	1608 $\pm$ 66	80	87	69	79	87
4000	3184 $\pm$ 52	83	82	79	78	76
	Mean $\pm$ SE	86 $\pm$ 5	92 $\pm$ 6	74 $\pm$ 3	76 $\pm$ 6	96 $\pm$ 9

Figure 28. Recovery of porcine relaxin added to the pH 5 soluble fraction of crude extracts of placentomes. The control represents recovery of porcine relaxin in the absence of placentomal crude extract.

curve ranged from 19.53 to 5,000 pg. The mean recovery of porcine relaxin was $85 \pm 4\%$ in the presence of placentomal crude extracts and $100 \pm 0\%$ in the absence of crude extracts.

Conclusions. The higher concentrations of the relaxin-like peptide found in the bovine corpus luteum compared to that of the placenta is interpreted that there is a systemic role for luteal immunoreactive relaxin. Anderson et al. (1982) and Randel et al. (1988) have detected immunoreactive relaxin in the plasma of pregnant cows that peaks near parturition. Immunoreactive relaxin, of luteal origin, may thus be acting on the cervix and pelvic ligaments of the cow through transport in the blood.

Transport of immunoreactive relaxin in the blood could also account for the low concentrations found in placental tissues. Placental tissue is heavily vascularized and systemic immunoreactive relaxin may be binding to its receptor. This possibility is further sustained by the fact that the molarity of immunoreactive relaxin ($8.3 \times 10^{12} \text{ M}^{-1}$) calculated from the Tris buffer crude cotyledonary extracts was below the K_d (approximately 10^9 to 10^{10} M^{-1}) reported for the relaxin receptor in the rat and pig (Mercado-Simmen et al., 1980, 1982). Location of the mRNA for immunoreactive relaxin in bovine placental tissues would resolve whether there is local synthesis rather than sequestration from a luteal source. Local synthesis by the placenta may result in a paracrine action for relaxin in the cow similar to that proposed for the human.

Isolation and purification of the relaxin-like peptide from the cow, whether from the corpus luteum or the placenta, appears to be a difficult task using the techniques presently available. A possibility for finding the relaxin-like peptide in the cow exists in the isolation of the mRNA for relaxin as was done with two human relaxin genes (Hudson et al., 1983, 1984). The nucleotide sequences were used to deduce the amino acid sequence of human relaxin. Although mRNA levels for the relaxin-like peptide apparently are low in view of the low levels of native peptide, use of the polymerase chain reaction, a recently reported procedure, may enable amplification of the bovine relaxin mRNA which could lead to isolation and sequence information.

Addendum. Utilizing a 20 to 40% ammonium sulfate precipitation from extracts of placentoms day 246 to day 275 of pregnancy two bands of relaxin immunoactivity were detected in a western blot. An antibody to porcine relaxin was utilized after transferring the proteins to Nitrocellulose paper. The two immunoactive bands relative molecular weight are 60,000 and 17,000 has been confirmed in extracts from four cows of 14 Israeli Holsteins evaluated.

The 17,000 Dalton peptide is of an appropriate size for the prohormone as discussed in previous sections. The 60,000 Dalton proteins has been observed in other species but its significance has never been explained. A 6,000 Dalton immunoreactive relaxin was not observed nor expected since the tested material had previously been dialyzed with loss of all proteins less than 8,000 Daltons. These findings are very exciting for we are in a good position from the above reported work to pursue further purification and amino acid sequencing directly from a western blot.

3. Effects of Relaxin on Steroidogenesis and Prostaglandin Synthesis.

In addition to the characterization of bovine relaxin, we have evaluated the effect of relaxin on in vitro progesterone and prostaglandin synthesis. Besides a systemic role for placental relaxin there could be an important paracrine role in remodelling of connective tissue (Steinetz et al., 1983) via regulation of placental (Shemesh et al., 1984; Gross and Williams, 1988) and uterine (Williams and Tahir, 1982) prostaglandin synthesis that may impinge on the release of the placenta at parturition. In addition relaxin has been shown to cause a decline in systemic concentrations of progesterone in the late pregnant cow (Musah et al., 1986a).

Administration of a pharmacological dose of relaxin to late pregnant beef heifers resulted in a precipitous decrease in peripheral plasma concentrations of progesterone within 90 min of infusion, and a rise in the peripheral concentration of estrone and 17 β -estradiol. Musah et al. (1987a) concluded that the decrease in secretion of progesterone caused by exogenous relaxin made available additional precursors for the biosynthesis of estrogens in the placenta. When high levels (6000 U) of relaxin were administered to heifers on day 275 of gestation, a peak release of oxytocin occurred followed by a significant decrease in plasma concentrations of progesterone (Musah et al., 1987a).

Effect of Relaxin on Luteal Progesterone Synthesis. When dispersed luteal cells from this first trimester pregnant cow were incubated with relaxin no effect on production of progesterone was observed (Table 6). However, when relaxin was incubated with only the small cell there was an increase in progesterone synthesis. These results might be interpreted that relaxin releases an inhibitory factor from the large cell that inhibits the progesterone increase induced by relaxin when incubated with the small cell. These observations need confirming and tested for at other stages of pregnancy.

Table 6. Progesterone Production by Dispersed Cells of the Corpus Luteum and Cotyledon from the Cow Days 60-90 of Pregnancy.			
Treatment	Placenta X $\pm$ SE	Progesterone (ng/ml)	
		Small Luteal Cell X $\pm$ SE	85% Small + 15% Large Luteal Cell X $\pm$ SE
Control	2.45 $\pm$.40	4.83 $\pm$ 2.12	2.26 $\pm$ 3.7
1000 pg Relaxin	4.78 $\pm$.52	12.61 $\pm$ 3.51	2.48 $\pm$.43
100 pg Relaxin	3.99 $\pm$.36	20.64 $\pm$ 3.85	2.44 $\pm$.61
10 pg Relaxin	6.34 $\pm$.25	74.70 $\pm$ 8.91	2.51 $\pm$.72

As reviewed above relaxin administration to late pregnant cows induces a marked decline in peripheral plasma concentrations of progesterone. We were unable to confirm this observation when eight late pregnant cows were infused, via the cervix, with 1 mg of highly purified porcine relaxin (Fig. 29). A possible explanation for the lack of an effect by relaxin on peripheral progesterone is

Figure 29. Three cows were administered 1 mg porcine relaxin (T) intracervically an average of 10 days prior to calving. Relaxin had no apparent effect on the plasma concentrations of progesterone. C = cow administered placebo.

no relaxin was observed in the plasma following its administration via the cervix. The reason for this is unexplained but consistent with additional studies that have yielded a low to undetectable systemic concentration of relaxin when administered via the cervix (*vide infra*).

Effect of Relaxin on Placental Steroid Metabolism. The effect of relaxin on the uterus is varied, including increased collagen remodelling, water inhibition, protein and glycogen synthesis (Bryant-Greenwood, 1982). Other effects of relaxin include modulation of $\text{PGF}_{2\alpha}$ and oxytocin induced uterine contractility (Porter, 1979). The relaxin induced precipitous decline in plasma progesterone in the late pregnant cow (Musah et al., 1986a) may be mediated through the release of uterine $\text{PGF}_{2\alpha}$. Relaxin reportedly modulates prostaglandin metabolism through inhibition of basal and oxytocin stimulated prostacyclin formation in the myometrium of the pregnant rat (Williams and Tahir, 1982). When myometrial tissues were incubated with arachidonic acid no inhibitory effect was observed, indicating an inhibition of the enzyme phospholipase A_2 . Effects on other prostaglandins, i.e., $\text{PGF}_{2\alpha}$ and PGE_2 were not studied.

Our unexpected preliminary findings indicate that dispersed cells of the cotyledon from day 60 to 90 of pregnancy responds to the addition of relaxin with increased progesterone production (Table 6). In further investigation, dispersed cells of the caruncle (500,000 cells/ml) were incubated for 24 h in the absence or presence of relaxin (10 ng/ml) or methyl isobutyl xanthine (MIX), a phosphodiesterase inhibitor (0.3 mM). Tissues were harvested and incubated as previously described Shemesh et al. (1983; 1984). Briefly, tissues were enzymatically dispersed with collagenase and dispersed cells incubated in medium 199 supplemented with 5% calf serum at 37°C in an atmosphere of 5% CO_2 - 95% air. MIX alone increased progesterone production over control levels (Table 7). Relaxin and MIX together produced an additive effect that resulted in a significant increase over control levels ($P < 0.05$). This additive effect did not appear to be dependent on the stage of pregnancy tested. The increase due to relaxin alone was not statistically significant.

Table 7. Effect of Relaxin and MIX on Progesterone Production by Cells of the Maternal Caruncle.				
Crown rump length (cm)	Relaxin	MIX	Relaxin + MIX	No.
15-28	122 ± 10	144 ± 23	160 ± 6	8
35	92	281	274	1
45	90	112	124	1
term	103	97	305	1
all	114 ± 10	135 ± 18	181 ± 20	11

MIX vs MIX + Relaxin $P < 0.05$

MIX vs Control $P < 0.05$

Relaxin vs Control NS $P < 0.1$

Control Progesterone 5.44 ± 1.8 ng/500,000 cells/24 h

Table 8. Eliminated

Table 9. Comparison of Fetal Cotyledon vs Maternal Canuncle Progesterone Production (15-45 cm Fetal Crown-Rump Length)				
Control	Fetal 2.84 ± 1.08	Maternal 1.05 ± 0.34	ng/500,000 cells	No.
Expressed as Percent of Control				
Relaxin	117 ± 5	112 ± 11		6
MIX	140 ± 10	151 ± 45		6

Relaxin alone did not significantly increase progesterone levels in both the cotyledon and caruncle above that seen in controls (Table 9). Time course experiments for maternal caruncle (Table 10) and fetal cotyledon (Table 11) showed that the increased synthesis of progesterone with time of incubation over 24 h did not differ between relaxin treated and non-relaxin treated controls.

Table 10. Time Course Experiment for Maternal Caruncle from Second Trimester. Effect of Relaxin on Progesterone and Prostaglandin F _{2α} Production.			
Experiment #1			
(ng/500,000 cells)			
<u>Progesterone</u>			
	2h	6h	24h
Control	0.34	0.68	2.54
Relaxin 10 ng	0.33	1.05	2.20
<u>PGF_{2α}</u>			
Control	1.19	2.21	3.12
Relaxin 10 ng	0.94	1.88	3.50
Experiment #2			
<u>Progesterone</u>			
Control	0.14		0.54
Relaxin 1 ng	0.12		0.63
MIX	0.34		0.40
<u>PGF_{2α}</u>			
Control	2.15		3.67
Relaxin	1.09		5.92
MIX	0.65		2.50

Table 11. Time Course Experiment for Fetal Cotyledons From Second Trimester. Effect of Relaxin on Progesterone and Prostaglandin F _{2α} Production.		
Progesterone		
	<u>2h</u>	<u>24h</u>
Control	0.94	2.15
Relaxin 1 ng	1.63	2.54
MIX	2.86	5.20
PGF _{2α} (ng/500,000 cells)		
Control	1.02	2.24
Relaxin 1 ng	3.16	4.09
MIX	0.67	1.49

Relaxin was not observed to have an effect on estrogen production by second trimester placental cells. In the near term maternal caruncle, estrogen was not detectable in long term incubations (Table 12). In one of the five short term incubations, relaxin did increase estrogen production by the maternal caruncle cells. In addition relaxin increased estrogen production by

Table 12. Net Production of 17β-Estradiol by Mid-Semester Maternal Caruncle Cells (pg/500,000 cells/24h).		
Added compound	Estradiol (pg)	No.
Control	ND*	11
MIX	541 ± 58	11
Relaxin	ND	11
8Br-cAMP	ND	8

*ND = (50pg/500,000 cells/24 h

the placental in the periparturient cow. Cyclic AMP also stimulated estrogen production (Table 13). In both cases stimulation of estrogen production was only observed in the cow shortly before parturition.

Table 13. Net Production 17 β -Estradiol by Term Maternal Caruncle. (33 h prior to parturition, 284 days)	
Added compounds	Estradiol (pg/500,000 cells/2 h)
Control	252
MIX	593
Relaxin 1 ng	800
Relaxin 5 ng	1004
Relaxin 10 ng	1258
cAMP	1441
cAMP	975
cAMP + cGMP	519

Conclusion. Time course studies on progesterone secretion showed a viable incubation of caruncular and cotylendnary tissues. Although, in several cases relaxin alone increased progesterone synthesis the increase was not statistically significant. When relaxin and MIX were added there resulted an increase in progesterone synthesis that was greater than the addition of MIX alone. This indicates no direct effect of relaxin on progesterone synthesis but possibly an indirect effect mediated through other compound(s). The finding of an increase in relaxin and cAMP stimulated estradiol synthesis only in the one cow tested shortly before parturition, is indicative of the rapid changes which occur in steroidogenesis in the cow shortly before parturition.

Effect of Relaxin on Prostaglandin Synthesis. Prostaglandin synthesis by dispersed cells of the cotyledon (500,000 cells/ml) was evaluated following incubation for 24 h in the absence or presence of relaxin (10 ng/ml) or MIX (0.3 mM). Relaxin ($P < .10$) and MIX ($P < .01$) resulted in a depression of prostaglandin synthesis (Table 14 and 17). Relaxin and MIX together had no significant effect above that seen with MIX alone. This was true for fetuses smaller than 30 cm crown rump length. However, in a larger fetus relaxin and MIX together did cause a further decrease in prostaglandin production and at term relaxin alone caused a substantial decrease in prostaglandin production. However, in another series of experiments near parturition relaxin had no effect on PGF_{2 α} production by the maternal caruncle (Table 16). However, MIX did have a suppressive effect similar

to that seen earlier in gestation. Very pronounced was the suppressive effect of the cyclic nucleotides, cAMP and cGMP, on both synthesis of $\text{PGF}_{2\alpha}$ and PGE (Table 18).

Table 14. Effect of MIX and Relaxin on Prostaglandin Production by the Maternal Caruncle.				
		% of Control		
Crown rump length (cm)	Relaxin	MIX	Relaxin + MIX	No.
15-28	117 ± 13	50 ± 7	63 ± 9	8
35	92	41	28	1
45	102	74	62	1
term	52	68	62	1
term 3 ng/ml Relaxin	61	68	62	1
10 ng/ml Relaxin	18	-	-	1
30 ng/ml Relaxin	35	-	-	

MIX vs Control $P < 0.01$

Relaxin vs Control NS $P < 0.1$

Relaxin + MIX vs MIX alone NS

Control prostaglandin 2.17 ± 78 ng/500,000 cells/24h

Table 15. Eliminated

Table 16. Comparison of PGF _{2α} to PGE Production by Maternal Caruncle Taken From Cows Close to Parturition.			
(ng/500,000 cells/24h)			
	PGF _{2α}	PGE	
Control	1.31 ± 0.04	12.43 ± 4.7	No.
Expressed as Percent of Control			
MIX	82 ± 21	88 ± 7	5
Relaxin 1 ng	105 ± 8	101 ± 8	5
Relaxin 5 ng	101 ± 8	115 ± 17	5
Relaxin 10 ng	102 ± 14	99 ± 14	5

Table 17. Dose Response for the Effect of Relaxin on PGF _{2α} Production in Maternal Caruncle of Second Trimester	
ng/500,000 cells/24 h	
Control	2.17
Relaxin 3 ng	1.32
Relaxin 10 ng	0.39
Relaxin 50 ng	0.75

Table 18. Effect of cAMP, cGMP and cAMP+cGMP on Total Prostaglandin Production by the Caruncle in Two Cows Near Parturition (0.5 mg/ml).						
Treatment	Incubation time	PGF _{2α} ng/500,000 cells	PGE ng/500,000 cells	% control F _{2α} E (net synthesis)		Age days
Experiment 1#						
Control	2h	0.03 ± 0.05	0.40 ± 0.14		100	250
cAMP		0.31 ± 0.02	0.28 ± 0.06	-	44	
cGMP		0.25 ± 0.04	0.30 ± 0.07	-	51	
cAMP+cGMP		0.25 ± 0.01	0.18 ± 0.03*	-	1	
Experiment #2						
Control	2h	0.72 ± 0.08	10.57 ± 0.61	100	100	284
cAMP		0.52 ± 0.03	5.99 ± 0.18*	51	1**	
cGMP		0.69 ± 0.05	7.17 ± 0.30	92	1	
cAMP+cGMP		0.51 ± 0.09	3.82 ± 0.15*	72	1	
Experiment #1						
Control	24h	0.95 ± 0.08		100		260
cAMP		0.44 ± 0.05	-	20	-	
cGMP		0.79 ± 0.05	-	60	-	
cAMP+cGMP		0.28 ± 0.02	-	1	-	
Experiment #2						
Control	24h	2.83 ± 0.09	24.66 ± 2.49	100	100	284
cAMP		2.16 ± 0.14*	8.69 ± 0.71*	73	6	
cGMP		3.11 ± 0.31	12.99 ± 0.92*	111	31	
cAMP+cGMP		1.91 ± 0.13*	4.37 ± 0.39*	63	1	

*Significantly different from control P<0.05; Mean ± SE of five replicates

**Values less than time zero incubation recorded as 1%.

Conclusion. It was difficult to demonstrate that relaxin had a statistically significant effect on the production of progesterone or prostaglandin in either maternal caruncle or fetal cotyledon. On the other hand, in a few experiments there was a clear dose and time related increase or suppression of progesterone and synthesis but we could not repeat the observations. Therefore if relaxin is involved in steroidogenesis or prostaglandin synthesis the action is indirect and is mediate through some other compound.

It is of interest to note that MIX stimulated progesterone production and inhibited $\text{PGF}_{2\alpha}$ production. Since cAMP and cGMP also suppressed PGE and $\text{PGF}_{2\alpha}$ production, it is probable that the effect of MIX on prostaglandin production is the result of its inhibition of phosphodiesterase rather than increasing intercellular calcium. This assumption is supported by the lack of an effect of calcium ionophore on PG synthesis (data not included).

The finding of an increase in cAMP stimulated estradiol synthesis only in the one cow tested shortly before parturition, is indicative of the rapid changes which occur in steroidogenesis and PG synthesis in the cow shortly before parturition.

The possibility that relaxin has a paracrine effect in that it inhibits prostaglandin synthesis during gestation and increases prostaglandin synthesis once parturition is initiated can still not be ruled out and there is some evidence that this is indeed the case in the human placenta (Bernal et al., 1987).

4. Response of Nonpregnant and Periparturient Cows to Porcine Relaxin.

Previous studies have shown that exogenously administered relaxin to the cow results in an induced parturition accompanied by a significant increase in pelvic area, cervical dilation, no incidence of retained placenta and no mortality of calves up to 1 week post-partum (Anderson et al., 1982; Perezgrovas and Anderson, 1982; Musah et al., 1986a,b). Induction of parturition with glucocorticoids or prostaglandins is accompanied with the problem of a high incidence of retained placenta. Musah et al. (1987b; 1988) showed that when relaxin was administered in combination with dexamethasone or cloprostenol to induce parturition the problem of retained placenta was eliminated.

Evaluation of Different Routes of Administration of Relaxin to Cows. To complement our in vitro studies and the above reports several studies were conducted in which relaxin was administered to the nonpregnant and late pregnant cow (Paccamonti, 1988; Paccamonti et al., 1989; 1991). In an attempt to determine the optimal carriers for extending the circulating concentrations of relaxin and thus biological potency, five nonpregnant Angus cows were submitted to five treatments with relaxin administered 72 h post-PGF_{2α}. The experiment was replicated five times with each cow receiving any treatment only once. Treatments with relaxin for Trial 1 were: a) 1 ml of 5% beeswax in sesame oil i.m. (control), b) 1 mg porcine relaxin in 1 ml K-Y gel intracervically (i.cx.), c) 1 mg relaxin in 1 ml 5% beeswax in sesame oil s.c., d) 1 mg relaxin in 1 ml 5% beeswax in sesame oil i.m., or 3) 1 mg relaxin in 1 ml saline i.v. There was a 2 week interval between treatments (Paccamonti, 1988; Paccamonti et al., 1989; 1991).

Much variation existed between individual animals in Trial 1, believed in part to reflect variations in absorption and delivery within the carriers used. The time at which the peak concentration of relaxin occurred was earlier for the i.cx. (2.7 h) than for the s.c. (10.8 h; $P < 0.01$) or for the i.m. in oil (8.3 h; $P < 0.03$) treatments (Fig. 30). However, there was no difference detected between the time of peak concentration of relaxin resulting from i.m. and s.c. routes of administration

EXPERIMENT 1

1 mg Porcine Relaxin

Figure 30. Regression curves of plasma concentrations of relaxin in cows administered 1 mg relaxin in 1) K-Y gel intracervically, 2) 5% beeswax in sesame oil intramuscularly, 3) 5% beeswax in sesame oil subcutaneously.

($P>0.10$). The maximum concentration of relaxin after s.c. administration (1.8 ng/ml) was greater than after i.c.x. (0.4 ng/ml; $P<0.05$) or i.m. (1.0 ng/ml; $P<0.10$) administration. No difference could be detected between the i.c.x. and i.m. routes ($P>0.10$) in maximum concentration of relaxin. The area under the curve was also lower for i.c.x. (5.5) than for s.c. (30.5; $P<0.05$) or i.m. (30.0; $P<0.05$) treatments but did not differ between the s.c. and i.m. ($P>0.10$). Orthogonal contrasts of treatments showed that the regression curve of relaxin treated cows differed from the vehicle treated controls, i.c.x. treatment differed from the i.m. and s.c., and that i.m. differed from the s.c. ($P<0.001$).

The failure of i.m. relaxin in beeswax/oil to attain high plasma concentrations of relaxin led to Trial 2 in which there was a comparison of an i.m. injection of relaxin in beeswax/oil versus a saline carrier. In addition, the i.v. treatment was repeated from Trial 1. Trial 2 was repeated with

five nonpregnant Hereford cows utilizing the randomized block design of Trial 1. The entire trial was repeated three times such that each cow received each treatment once and no cow received any treatment twice. The administered relaxin in beeswax/oil in Trial 2 (Fig. 31) had a much sharper initial rise than the same treatment in Trial 1 (Fig. 30). Plasma concentrations of relaxin returned to baseline by 24 h post-injection. However, cows injected with relaxin in beeswax/oil i.m. varied considerably in their response. The peak level seen with the i.m. route using saline is over four times greater than seen with the beeswax/oil carrier, reaching 12.6 ng/ml by 1 h post-injection and returning to baseline by 24 h (Fig. 31). Although the peak concentration of relaxin after injection in beeswax/oil was lower than after injection in saline ($P < 0.005$), the time of the peak concentration of relaxin did not differ ($P > 0.1$) between treatments. The area under the curve was less for the relaxin administered i.m. in beeswax/oil (26.0) than for either the i.m. in saline (71.7) or i.v. in saline (75.5)

Figure 31. Regression curves of plasma concentrations of relaxin in cows administered 1 mg of relaxin intramuscularly in a) 5% beeswax in sesame oil or b) saline.

($P < 0.001$) treatments. There was no difference between the i.m. in saline and i.v. routes ($P > 0.1$). Neither cow nor period (replication) significantly affected the concentration of relaxin over time ($P > 0.10$) in either trial. The concentration of progesterone at the time of relaxin administration also did not affect the level of relaxin in either Trial 1 or 2 as determined by MANOVA ($P > 0.10$). The half-life of relaxin was calculated as $t_{1/2} = [(ln 2)/b]$ and determined from the i.v. injections to be 16.56 min (Fig. 32).

Figure 32. Regression curve of plasma concentrations of relaxin in cows administered 1 mg relaxin intravenously in 1 ml saline.

Conclusion. The use of beeswax/oil as a carrier resulted in variation in plasma concentrations of relaxin among cows. This variation reflects the apparent difficulty in evenly distributing the relaxin throughout the carrier that is coupled with differences in absorption of relaxin from the injection site. The use of saline as an alternative carrier resulted in greater homogeneity of plasma

concentrations of relaxin between cows. The 5% beeswax in sesame oil carrier was chosen because earlier reports showed the potentiation or prolongation of relaxin resulting from its slow release and thereby increasing the biological activity of the hormone. However, in cows the rate of release was so slow that the maximum concentrations attained were low in relation to those seen after administration of relaxin in saline. In addition, the area under the curve for i.m. administered relaxin in beeswax/oil is only one third the area for i.m. in saline. Although potentiation is assessed by biological response and is not solely dependent on blood concentrations, it is unlikely that the beeswax/oil carrier would result in potentiation of relaxin in the cow in view of these observations.

Production of Antibodies to Porcine Relaxin Following Administration of Porcine Relaxin to Cows. In Trial 1 and 2 numerous samples gave assay values of relaxin which differed greatly from the expected (Paccamonti, 1988; Paccamonti et al., 1989; 1991). Initially this was considered a technician error at the time of sample collection or processing in the laboratory. These aberrant results were characterized by high concentrations of relaxin in the pretreatment plasma samples. This is a time (nonpregnancy) when there should be no relaxin. In Trial 1 period 3, one cow (336) had apparent pretreatment concentrations of 11-15 ng/ml relaxin. In period 4, this same animal, treated i.c.v. had apparent pretreatment values of 27-29 ng/ml and post-treatment values that ranged from 19-32 ng/ml over the 48 h sampling period. Concentrations of relaxin of this magnitude were seen only within the first 30 min after i.v. administration. In period 5, cow 336, a control receiving only the beeswax in oil carrier, had an apparent pretreatment concentration of relaxin of 1 ng/ml which decreased over the next 24 h. At 36 h the apparent concentration of relaxin rose to 31 ng/ml. Cow 839 also had high pretreatment concentrations in period 4 and period 5 (37-40 ng/ml).

Once an animal began to show elevated pretreatment concentrations of relaxin (>1.0 ng/ml), pretreatment concentrations remained elevated in subsequent periods. Another trend observed with the assays is that one animal (336) had high pretreatment levels in period 3, two animals in period

4 (335 and 839), and four in period 5 (245, 336, 449 and 839). There is a pattern that with increased exposure to relaxin, more animals were showing apparent high plasma concentrations before receiving relaxin (Fig. 33). This pattern of high pretreatment relaxin in nonpregnant cows indicated these values were a function of the assay and in this case the in vivo production of endogenous antibodies to porcine relaxin.

Period 1 2 3 4 5						Period 1 2 3			
Cow						Cow			
245	i.cx.	i.v.	control	i.m. beeswax	(s.c. beeswax)	270	i.m. beeswax	i.v.	(i.m. saline)
336	s.c. beeswax	i.m. beeswax	(i.v.)	(i.cx.)	(control)	300	i.m. saline	i.v.	i.m. beeswax
449	control	i.cx.	i.m. beeswax	s.c. beeswax	(i.v.)	388	i.v.	i.m. saline	i.m. beeswax
564	i.v.	s.c. beeswax	i.cx.	control	i.m. beeswax	3103	i.v.	i.m. saline	i.m. beeswax
839	i.m. beeswax	control	s.c. beeswax	(i.v.)	(i.cx.)	4224	i.m. saline	i.m. beeswax	(i.v.)
Experiment 1						Experiment 2			

Figure 33. Animal assignments for Experiment 1 and 2. Circles represent pretreatment samples precipitating >15% [125 I] relaxin at a 1:10 dilution of plasma. i.cx.=intracervical; s.c.=subcutaneous; i.v.=intravenous; i.m.=intramuscular.

Ammonium sulfate precipitation of the immunoglobulin bound [125 I] relaxin confirmed the presence of antibodies to porcine relaxin. Plasma samples precipitating 15% or more of the [125 I] relaxin at a 1:10 dilution of plasma were considered to contain antibodies to porcine relaxin at a concentration high enough to interfere with the porcine relaxin RIA (Fig. 34). Once a cow developed an antibody response, she continued to show the response in subsequent periods. In addition, with repeated exposures, additional cows demonstrated the antibody response. A review of

Figure 34. Percentage of [^{125}I] relaxin precipitated by ammonium sulfate from plasma collected prior to relaxin administration.

the pattern of antibody response showed that animals received an i.m. injection of relaxin in 5% beeswax/oil before development of measurable antibody production. In addition plasma from cows in Trial 1 which precipitated the highest counts of [^{125}I] relaxin also received a s.c. injection of relaxin in 5% beeswax in oil. In Trial 2, cows were only treated with porcine relaxin three times. One cow in each of the first two periods received relaxin in beeswax/oil i.m. Both cows developed an antibody response by the third period. Data from periods in which cows showed significant antibody production were not used in the statistical analyses for evaluation of treatments (vide supra).

One of the unexplained observations in Trial 1 and 2 was the low circulating concentrations of relaxin following its intracervical administration. The cows used in the present study were estrus induced with $\text{PGF}_{2\alpha}$ in an attempt to control the variability of the stage of the estrous cycle. At

estrus, copious amounts of mucus are produced which flow in the direction of the vulva and external environment that may result in the removal of material placed in the cervix. During pregnancy there is a thick, mucoid cervical plug providing a barrier between the external environment and the developing fetus. This could serve to trap substances deposited in the cervix and may afford more effective absorption. In addition estrous synchronized cows would be under a shorter estrogenic stimulation than seen with pregnant cows. With this in mind late pregnant cows were administered relaxin i.c.v. and frequent blood samples taken for 24 h following treatment. No significant concentrations of relaxin were detected in the plasma of these cows post-treatment. Subsequent studies indicated the site of deposition, i.e., os-cervical, mid-cervical or uterus, did not influence the absorption of relaxin into systemic circulation.

Conclusion. It appears that the beeswax in sesame oil carrier, rather than potentiating or prolonging the effects as seen in mice and guinea pigs, respectively, acts as an adjuvant, in effect immunizing the cows against relaxin. The clinical significance of these findings should not be overestimated since seldom would cows receive repeated injections of porcine relaxin particularly at such short intervals. In light of the findings in this report, it is also highly unlikely that a 5% beeswax in sesame oil carrier would be used. Considering these two factors, production of antibodies to porcine relaxin would not be expected under field conditions. Nonetheless a preparation of bovine relaxin would be an ideal solution to the problem. The failure to detect relaxin in systemic circulation after cervical administration remains unexplained.

Relaxin Induction of Parturition. One of the objectives of this grant was to evaluate the effect of exogenous administration of porcine relaxin on parturition in the dairy and crossbred beef cow. Relaxin was deposited i.c.v. in a K-Y gel carrier or injected i.m. in either beeswax/oil or saline carrier (Paccamonti, 1988; Paccamonti et al., 1989). Animals receiving relaxin i.c.v. or i.m. had gestation lengths 6.6 and 1.0 d shorter than controls ($P < 0.05$; Table 19). Animals treated with relaxin.

One possible explanation for the prolonged interval to parturition may be the day of treatment since animals in three experiments were treated 7.3 ± 0.4 day (mean $\pm$ SE) before their calculated due date (10.2 days before actual due date). There may be an "endocrine window" immediately prior to parturition in which exogenous relaxin would more effectively induce parturition. To test this hypothesis an expanded field study was conducted to evaluate both doubling of the dose of relaxin and day of treatment. Crossbred cows were treated twice with an i.m. injection of relaxin (am and pm) at 8 or 4 day before the expected day of parturition. Although there was a 2.0 day difference in gestation length between animals treated with relaxin 8 day before their due date and controls treated with saline, this difference was not statistically significant ($P=0.16$; Table 21).

Table 21. Effect of Day of Relaxin Treatment on Gestation Length.			
Treatment	No. animals	Gestation length (d)	Difference in gestation length from controls
Control	12	287.5	0.0
Relaxin 4 day before due	11	287.9	-0.2
Relaxin 8 day before due	14	285.5	2.0

Conclusion. Intracervical deposition of relaxin may be more effective since the cervix is a target tissue of relaxin. Uptake could be direct, thereby avoiding the problems of absorption into the systemic circulation, redistribution, and metabolism of the hormone before reaching the target tissue. The results indicate intracervical administration of relaxin during a specified "endocrine window" may be the most effective approach for effectively inducing parturition or shortening gestation. This "endocrine window" would be a period of time in which relaxin would not initiate a series of events leading to parturition until the proper endocrine setting was present (Paccamonti, 1988; Paccamonti et al., 1989).

Relaxin Combined with Corticosteroids to Induce Parturition. If relaxin is only effective in inducing parturition when given shortly before naturally occurring parturition, its usefulness as an agent for induction of parturition is limited. However, reports of the use of relaxin in conjunction with other agents, such as prostaglandins or corticosteroids, offer an avenue of other beneficial clinical uses for relaxin. Relaxin may be useful in narrowing the time span in which parturition is expected as seen in the pig when combined with prostaglandin. Relaxin may also serve to reduce the incidence of retained placenta and decrease the interval from treatment to parturition when used with corticosteroids or prostaglandins.

Two experiments were conducted in which relaxin was used to induce parturition in conjunction with glucocorticoids to alleviate dystocia and the prevention of retained placenta. In collaboration with Dr. Michael Diskin, Galway, Ireland and induced parturition regimen was tested that was based on an initial administration of a long-acting glucocorticosteroid (30 mg Betsopart, Glaxo) followed 5 to 6 days later with a short-acting preparation (40 mg Betsolan Soluable, Pitman-Moore). Porcine relaxin (1.5 mg) was administered intramuscularly at the time of the second treatment with corticosteroid. This model (without relaxin) will induce calving up to one week before the expected calving date resulting in a 95% calving rate, considerably reduced incidence of calving difficulty, good calf survival rates and a 25% incidence of retained fetal membranes (RFM). The use of relaxin in this model is an ideal complement to the induction regimen.

Continental Friesian cross heifers pregnant to a Limousin sire with a known history of a high incidence of calving difficulty were treated on day 274 with Betsopart and on day 279 with Betsolan and porcine relaxin. Control animals received saline. The results are reported in Table 21. In a second trial heifers were treated on day 272 with Betsopart and on day 279 Betsolan and porcine relaxin. There were no saline controls in this trial. The results are reported in Table 22.

Table 22. Experiment 2- Induction of Parturition in Continental Friesian Cross Heifers with a Combination of Long and Short Acting Glucocorticoids and Relaxin

	Corticoid Induced Only	Corticoid Induced + Relaxin
Heifers	10	10
No calving	9	8
Gestation length (days)	279	279
Mean birth weight (kg)	38.0	38.7
Mean Calving Score ($\pm$ SE)	3.0 ± 0.4^a	1.9 ± 0.4^b
No Cows with RFM (%)	2 (22)	0 (0)

Induction Regimen: Day 272 30 mg Long-acting glucocorticoid. (Betsopart, Glaxo)
Day 277 40 mg short-acting glucocorticoid (Betsolan Soluable, Pitman-Moore)
Day 272 1.5 mg Porcine relaxin.

Control animals received saline on treatment days.

Calving Score: Score 1-unassisted birth to Score 6-Cesarean Section.

Calves were weighed at birth, were allowed to suckle their dam and weighed again at Day 50.

Values with different superscripts differ ($P < 0.05$).

Conclusion. In both trials the ease of calving (calving score) was significantly improved in heifers receiving relaxin. The area of the pelvis was significantly increased in heifers that received relaxin in Experiment 1. Pelvic area was not monitored in Experiment 2. In both experiments there was a trend towards a reduction in the incidence of retained fetal membranes in animals that received relaxin. Since the incidence of retained fetal membranes was not as high as previously experienced, relaxin was not adequately tested as to its efficacy to alleviate this problem in the corticosteroid induced heifer. The prematurity of parturition had no effect on calf survival or subsequent performance of the calf. These results are encouraging, particularly, since our earlier work with intramuscular injections of relaxin were ineffective in the induction of parturition.

5. Oxytocin Receptors in Reproductive Tissues.

Musah et al. (1989) investigated the role of relaxin in luteolysis by determining the effects of porcine relaxin on the systemic plasma concentrations of both oxytocin and progesterone in late pregnant heifers. When high levels (6000 U) of relaxin were administered to heifers on day 275 of gestation, a peak release of oxytocin occurred. Following the peak release of oxytocin a significant decrease in plasma concentrations of progesterone was observed, confirming previous studies (Musah et al., 1987).

Musah et al. (1990) reported with cultured bovine luteal cells that exogenous relaxin suppressed the release of oxytocin into the culture medium in a dose-dependent manner. Furthermore, the suppression of oxytocin release by relaxin was greater when cells were obtained from corpora lutea of more advanced stages of gestation. On the other hand, addition of cloprostenol (an analogue of prostaglandin $F_{2\alpha}$) induced the release of oxytocin into the culture medium. However, the induction of oxytocin released by cloprostenol was suppressed when relaxin was added. Co-culture of relaxin and estrogen together resulted in an enhancement of the relaxin-induced suppression of oxytocin release by bovine luteal cells. The effects described above for relaxin indicate an interrelationship between relaxin and oxytocin. Based on this our studies were expanded to evaluate the oxytocin receptor in the cow.

Oxytocin and Vasopressin Receptors in the Bovine Uterus During the Estrous Cycle and Early Pregnancy. In collaboration with Dr. Anna Riitta Fuchs (Cornell Medical School) we have investigated the binding of oxytocin and vasopressin to the uterus of the cow on days 0, 7, 10, 14, 17 and 21 of the estrous cycle and early pregnancy (Fuchs et al., 1989, 1990a-d). The uterus was removed immediately after slaughter, and the removed endometrium was frozen. Serosa was removed from the myometrium before freezing. Briefly the tissues were homogenized in a 50 mM Tris buffer containing PMSF and EDTA which dissociates all endogenous ligands from the oxytocin (OT)

binding sites. A pellet precipitating between 1000 xg and 16000 xg was used for the receptor assay with a buffer containing 5 mM and 0.1% BSA, pH 1.8, protein concentration 200 to 300 µg/tube. The labeled ligand was [3 -H] OT and [3 H] AVP from New England Nuclear. All tissues were assayed in duplicate using four concentrations of labeled ligand and 1 µM unlabeled ligand to determine nonspecific binding. The binding data were analyzed with the LIGAND program. Values are expressed as femtomoles ligand bound per mg membrane protein. Saturation analyses were carried out with increasing concentrations of labelled ligands in the range from 0.10 to 11.5 nM.

Myometrium as well as endometrium possessed high affinity, saturable binding sites for both labelled OT and AVP. Analysis of the data indicated that for OT a single class of binding sites was present in both tissues and at all stages (Fig. 35). The binding affinity (K_D) was the same on days

Figure 35 Scatchard analysis of [3 H]-oxytocin (OT) binding in bovine endometrium of Day 0 of the estrous cycle.

examined, K_d = 0.9 ± 0.1 nM (X + SE). For AVP one class of binding sites was found in the myometrium (Fig. 36), K_d = 1.3 ± 0.2 nM (X ± SE), but in the endometrium of estrous cows (day 0), the two site model fit the data significantly better (Fig. 37). The high affinity site was absent on day 10, when the receptor concentration was much lower. The K_d of the lower affinity endometrial AVP binding sites was 6.7 ± 0.9 nM (± SE). The K_d of the high affinity endometrial AVP binding

Figure 36. Scatchard analysis of [3 H]-arginine vasopressin (AVP) binding in bovine myometrium on Day 0 of the estrous cycle.

Figure 37. Scatchard analysis of [3 H]-arginine vasopressin (AVP) binding in bovine endometrium on Day 0 of the estrous cycle. The two site model gave significantly better fit than one site model, $P=0.014$.

site (0.5 ± 0.08 nM, $X \pm SE$) was significantly greater than the K_d of the OT binding sites, which in turn was higher than the K_d of the lower affinity AVP binding site. There was no significant differences between the affinities of OT binding sites and AVP binding sites in the myometrium. (Fuchs et al., 1989, 1990a-c).

The concentration of [3 H] OT binding sites was on the average 50% higher than [3 H] AVP across the cycle in both tissues. Endometrial receptor levels varied significantly during the cycle, it was lowest on days 7, 10 and 14, rose significantly on day 17, and peaked on day 21 (Fig. 38). Myometrial receptor levels decreased from levels at estrus on days 7, 10 and 14, but the changes were not significant (Fig. 39). The ratio between endometrial and myometrial receptor concentrations

FIGURE 38
OXYTOCIN RECEPTORS IN BOVINE ENDOMETRIUM
DURING ESTROUS CYCLE AND EARLY PREGNANCY

FIGURE 39
OXYTOCIN RECEPTORS IN BOVINE MYOMETRIUM
DURING ESTROUS CYCLE AND EARLY PREGNANCY

changed from about 10 at estrus to less than 1 in the luteal phase. In early pregnancy, the receptor levels did not differ from nonpregnant levels on days 7 and 14, but on day 17 the endometrial receptor concentrations were significantly lower, and on day 21 those in both tissues were significantly lower. The endometrial OT and AVP receptor concentrations were inversely correlated with plasma concentrations of progesterone ($P = 0.005$) with no correlation to plasma estradiol,

whereas the concentrations of myometrial receptor showed no correlation to plasma progesterone but an inverse correlation with plasma estradiol ($P = 0.004$; Fuchs et al., 1989; 1990a-c). The difference between nonpregnant and early pregnant concentrations of endometrial OT receptor on day 17 and 21 after mating raised the question whether the embryo and its secretions influence endometrial OT receptor concentrations by a local paracrine route or by a systemic route, mediated by luteal progesterone secretion (Fig. 38). To investigate this question we measured OT receptor concentrations in both the ipsi- and contralateral uterine horns of pregnant cows on days 17 and 19 after mating (Fuchs et al., 1991b).

There was no difference in the endometrial OT receptor levels between the ipsi- and contralateral uterine horns of pregnant cows on either day 17 or 19, nor was any difference observed between the ipsi- and contralateral myometrial OT receptor concentrations on those days (Fig. 40).

FIGURE 40
OT RECEPTORS IN IPSI- AND CONTRA-LATERAL UTERINE HORNS OF PREGNANT COWS ON DAYS 17 AND 19 AFTER MATING.

FIGURE 41
PROGESTERONE IN BOVINE PLASMA DURING ESTROUS CYCLE AND EARLY PREGNANCY.

The cow with a degenerating embryo had significantly higher endometrial OT receptor concentrations than the other day 19 pregnant cows, and showed no difference between ipsi- and contralateral levels. The affinity of endometrial OT receptor was similar in all cases $K_d = 0.85 \pm 0.24$ nM all values are mean $\pm$ SD. B max for all ipsilateral endometria was 42.8 ± 35 fmol/mg and for contralateral endometria 27.7 ± 13 fmol/mg. They were significantly lower than the levels in nonpregnant cows (day 17: 345 ± 547 fmol/mg and day 19: 227 ± 190 fmol/mg (Fuchs et al., 1991b).

Plasma progesterone which began to fall on day 17 in nonpregnant cows remained at midluteal levels in pregnant cows (Fig. 41) and showed a strong negative correlation with endometrial OT receptor concentration ($r = -0.85$) while the correlation with OT receptor concentration ($r = -0.15$) was not significant. The suppression of endometrial OT receptor concentration may be due to systemic progesterone since endometrial OT receptors are more responsive to small changes in plasma progesterone than myometrial OT receptors. However, since conceptus secreted proteins are transferred to the contralateral horn during this period (Short et al., 1991) the suppression of OT receptor could also result from a local effect.

Conclusion. The presence of two classes of binding sites for AVP but only one for OT suggests that at least in the endometrium there are two separate binding sites for AVP and OT. The higher affinity AVP binding sites clearly is distinct from the OT receptor, whereas the lower affinity binding sites for AVP in the endometrium may or may not reflect binding of [3 H] AVP to the OT receptor with somewhat lower affinity than OT. Further work is required to distinguish between these possibilities. We conclude that the endometrial OT and AVP receptor concentrations are more tightly controlled by progesterone than in the case of the myometrium and that the bovine conceptus suppresses endometrial OT and AVP receptor concentrations in the preattachment stage.

No local effect of the embryo on the endometrial OT receptor concentrations could be demonstrated. The strong negative correlation of endometrial but not myometrial OT receptor

concentrations to plasma concentrations of progesterone suggests a hormonal rather than local paracrine regulation of the uterine OT receptor concentrations, by the bovine embryo. Endometrial OT receptor protein turnover appears to be more rapid and more sensitive to variations in circulating progesterone levels than myometrial OT receptor turnover.

Oxytocin and Vasopressin Receptors in the Ovary. Bovine ovarian stroma and follicles bound tritiated oxytocin and tritiated arginine vasopressin with similar affinity, whereas bovine corpora lutea bound tritiated oxytocin only (Fuchs et al. 1990d; 1991a). Competition for the binding of tritiated oxytocin by various agonist and antagonists was suggestive of receptor function. The number of oxytocin binding sites varied cyclically in all tissues. In bovine ovarian stroma and corpora lutea the concentrations were lowest on day 14 and highest on days 17, 19, and 21 after ovulation, with a striking peak in the luteal concentration on day 19. In large follicles the concentration of oxytocin binding sites was highest on the day of estrus and lowest on day 7. The number of arginine vasopressin binding sites was ~50% lower than oxytocin binding sites and the cyclic variations were not significant. Because bovine ovaries produce oxytocin and arginine vasopressin the results suggest a paracrine or autocrine role for these neuropeptides in luteolysis and ovulation (Fuch et al., 1990d; 1991a).

Addendum. Concentrations of oxytocin receptors in tissues of the uterus and fetal membranes were examined throughout pregnancy. The changes in bovine uterine concentrations of oxytocin receptor that take place during pregnancy and parturition are different from those reported in the rat and rabbit. In these species the myometrial oxytocin receptors remain very low throughout most of pregnancy and rise shortly before parturition.

The rise in bovine myometrial concentrations of oxytocin receptors during mid-pregnancy parallels the reported changes in uterine responsiveness to oxytocin (see figure below). The resulting contractions have been associated with periods of lowered fetal oxygenation that may in turn stimulate

the fetal-adrenal axis to secrete ACTH, a contributing factor in fetal adrenal maturation.

Endometrial concentrations of oxytocin receptors differed from the myometrial receptors with higher concentrations and a peak reached just before the onset of labor. Since oxytocin stimulates the release of $\text{PGF}_{2\alpha}$ from the nonpregnant bovine endometrium there may be a similar cascade operating during pregnancy. The failure of myometrial and endometrial concentrations of oxytocin receptor to follow a parallel course suggest a somewhat differential regulation of oxytocin receptor formation in the two uterine tissues.

In addition concentrations of oxytocin receptors were established throughout pregnancy in the amnion and chorio-allantoic membranes. Concentrations of receptors were low throughout most of gestation rising abruptly at time of labor. There were no differences in the affinity of the receptors between these membranes and the endometrium. These findings coupled with those for relaxin indicate both relaxin and oxytocin play an integral role in bovine parturition.

6. Large Luteal Cells and Their Secretory Granules.

Large luteal cells undergo progressive changes during the estrous cycle that conforms to similar changes observed during pregnancy (Field et al., 1985, 1991). On day 3 of the cycle, large cells resembled those on day 45 of pregnancy (i.e., contained little or no cytoplasmic lipid, no swollen or vacuolated mitochondria, no mitochondrial inclusions, and only a small numbers of secretory granules (Fields et al., 1992). Large luteal cells of corpora lutea taken at mid-cycle days 7, 11, and 14, like those of the second trimester of pregnancy, contained the largest number of secretory granules (Fig. 42) and some cytoplasmic lipid. On day 14 the large cells were observed to have certain morphological characteristics indicative of cells observed in the pregnant cow corpora lutea that are in transition to a regressive state (Fields et al., 1985), i.e., a decrease in secretory granules, cells with

Figure 42. Distribution of large luteal cell secretory granules throughout the estrous cycle of the cow. Percentage of large cells containing secretory granules(.....) was determined by scanning 100 large cells with the transmission electron microscope. The mean number of granules per large cell (.....) was determined from photomicrographs of 10 randomly selected large luteal cells with nucleus and cell membrane showing.

vacuolated mitochondria, and swollen mitochondria with dense inclusions. Involutive is a term we proposed (Fields et al., 1987; 1989) to define the corpus luteum where large luteal cells are in transition to senescence. The corpora lutea still have a large luteal weight and maintain high plasma concentrations of progesterone. Like in third trimester pregnant cows (Fields et al., 1985), these regressive characteristics were more pronounced during the last third of the cycle (days 17 and 19).

The immunostaining of oxytocin and neurophysin in a well defined region within the cytoplasm of large luteal cells reflects the discrete clustering of secretory granules. We reported for the first time that both oxytocin and neurophysin are co-localized in the same secretory granule of the large luteal cell (Fields et al., 1992). This was expected since the nucleotide sequence for oxytocin and neurophysin is contained within a single mRNA (Ivell and Richter, 1984; Ivell et al., 1985) localized in only the large luteal cell (Fehr et al., 1987). No oxytocin or neurophysin immunostaining or clustering of secretory granules were observed in small luteal cells.

The following paradigm is proposed. The early appearance and disappearance of the oxytocin mRNA indicates early formation of the oxytocin prohormone in the region of the rough endoplasmic reticulum. The delay in packaging of the prohormone into secretory granules in the Golgi complex may be the result of a delay in synthesis and thus availability of enzymes necessary for inclusion in the granules for processing of the oxytocin prohormone. The delay in processing of the oxytocin prohormone until day 7 reduces the hormones potency early in the cycle and a time when functional uterine oxytocin receptors are found in significant concentrations (Meyer et al., 1988; Soloff and Fields, 1989; Fuchs et al., 1990a-c and Jenner et al., 1991). Earlier studies have shown that oxytocin administered early in the cycle (days 1 to 7) will induce premature luteolysis. The high level of uterine oxytocin receptors found at estrus declines until reaching a nadir on day 7. This decline in receptor numbers reduces the possibility of mid-cycle oxytocin (days 7-14) initiating premature regression of the corpus luteum. If pregnancy occurs, uterine oxytocin receptors remain low (Fuchs

et al., 1990c; 1991b; Jenner et al., 1991) and luteal maintenance and pregnancy are sustained. If pregnancy does not occur, uterine oxytocin receptors increase significantly by day 17 and peak by day 21.

PGF_{2α} will deplete the large luteal cell of secretory granules (Fields et al., 1987; 1989) and induce an ovarian release of oxytocin (Schallenberger et al., 1984). In the nonpregnant animal, it is the timely release of oxytocin from the corpus luteum, its subsequent binding to uterine oxytocin receptors, and the release of uterine PGF_{2α} followed by a repetition of this sequence of endocrine events that leads to luteal regression in the ewe (Fint and Sheldrick, 1983; McCracken et al., 1984); Fields et al. (1991) showed that the days in which oxytocin is most rapidly depleted from the corpus luteum coincides with the nadir of uterine oxytocin receptors and the release of small nonluteolytic pulses of PGF_{2α} (Parkinson and Lamming, 1990) but prior to the luteolytic pulses of PGF_{2α} (Betteridge et al., 1984). Although a role for the mid-cycle secretion of oxytocin has not been determined, it may play a role in upregulating its own receptor as has been shown for other hormones. Luteal oxytocin that is released late in the cycle drives the luteolytic pulses of PGF_{2α}.

Oxytocin and neurophysin are major products of luteal secretory granules during the estrous cycle but not of secretory granules during the post-implantation period of pregnancy (>day 45). The number of granules and luteal oxytocin peak mid-cycle and then decrease precipitously. In pregnant cows, luteal granules rise from nearly undetectable numbers on day 45 to peak on days 180-210 (Fields et al., 1985). There are at least two populations of secretory granules, one of the cycle which contains oxytocin and one of the post-implantation period of which the contents are unknown.

Depletion of Secretory Granules with PGF_{2α} Early in the Cycle. The secretory granule population of pregnancy after day 40 has been characterized (Fields et al., 1985). Few granules were reported around day 40. Prior to day 40 the granule population has not been studied in the pregnant cow. In collaboration with Dr. Phillip Fields, University of South Alabama, a study is underway to

determine the effect of early pregnancy on the secretory granule population from day 11 to day 40. This is a transition period going from one that contains oxytocin to one with a population of secretory granules that do not contain oxytocin. We believe the latter contains relaxin but have been unable to immunocytochemically localize relaxin in these granules.

The major luteal secretory granule work conducted during this grant was to develop a model for depletion of secretory granules via a sub-luteolytic dose of $\text{PGF}_{2\alpha}$ during the early stages of the estrous cycle of the cow. The objective was to use this model to determine if a new cluster of granules can be generated in the granule depleted cell. Although, this is not a model of pregnancy, in the large luteal cell of early pregnancy there is depletion of the oxytocin containing granules prior to formation of granules devoid of oxytocin. It was also our objective to determine if there is regeneration of granules and do they contain oxytocin.

Cows were injected via the jugular vein with either saline or 25 μg $\text{PGF}_{2\alpha}$ at 0, 2 and 4 h on day 6 and 7 (Fig. 43). A preliminary study indicated this injection regimen had no effect on cycle length. However, 50 μg shortened the cycle. Corpora lutea ($n=3/\text{day}$) were harvested on day 6 (Pre- $\text{PGF}_{2\alpha}$), or day 7 (2 h post- $\text{PGF}_{2\alpha}$) or day 12 (recovery phase). $\text{PGF}_{2\alpha}$ caused a release of oxytocin into circulation and a decline in secretory granules on day 7 that were not replenished by day 12 (Fig. 44). Irrespective of treatment, all secretory granules contained oxytocin and neurophysin. It appears that early in the cycle a single cluster of secretory granules containing oxytocin and neurophysin are formed. Once secretory granules are depleted, the cell is incapable of generating a second population of secretory granules (Fields et al., 1991).

This model revealed two populations of large luteal cells which will be referred to as Alpha and Beta (Fig. 45). Beta large luteal cells (40-46 μm diameter) (Fig. 46) contained numerous mitochondria and a more apparent smooth endoplasmic reticulum than small Alpha large luteal cells (32-39 μm) (Fig. 47). In addition, Beta large luteal cells were more sensitive to $\text{PGF}_{2\alpha}$ as evidence

Treatment	No. Cows	Day Treatment	Day OVX	Objective
No Treatment	2	None	6	Baseline SG
Saline	2	6,7	7	Baseline SG
Saline	2	6,7	12	Increase SG
PGF _{2α}	3	6,7	7	Depletion SG
PGF _{2α}	3	6,7	12	Regeneration SG

Figure 43. Experimental design evaluating the effectiveness of a non-luteolytic dose of PGF_{2α} to deplete secretory granules and the ability of the large luteal cell to regenerate a new population of granules.

Figure 44. PGF_{2α} depletion of secretory granules in the Beta but not the Alpha large luteal cell.

Figure 45. Beta (left) and Alpha (right) large lutea cells from a saline treated cow. Note the abundance of smooth endoplasmic reticulum in the Alpha cell.

Figure 46. Beta cell from a cow treated with saline. Note the secretory granules at the plasma membrane adjacent to the capillary vessel and the abundance of smooth endoplasmic reticulum.

Figure 47. Alpha large luteal cell from a cow treated with saline.

Figure 48. Alpha (left) and Beta (right) large luteal cells taken from a cow treated with $\text{PGF}_{2\alpha}$. The Alpha cell has secretory granules and is otherwise normal whereas the Beta cell has been depleted of granules. In addition $\text{PGF}_{2\alpha}$ caused a swelling and washout effect of the Beta cell mitochondria but not in the case of the Alpha cell.

by their depletion of secretory granules and swelling of smooth endoplasmic reticulum and mitochondria (Fig. 48). These results indicate the corpora lutea contains at least two types of large luteal cells. Beta large luteal cells which are sensitive to sub-luteolytic doses of $\text{PGF}_{2\alpha}$ undergo regression first. This is followed by regression of Alpha large luteal cells when $\text{PGF}_{2\alpha}$ is elevated. This is evident by the fact that all large luteal cells are regressing by day 19.

Conclusion. In the nonpregnant cow there are two types of large luteal cells, Alpha and Beta. Secretory granules of both cell types contain oxytocin and neurophysin. $\text{PGF}_{2\alpha}$ decreased the number of granules in the Beta cell but had no effect on the granules of the Alpha cell. $\text{PGF}_{2\alpha}$ apparently inhibits formation of secretory granules in both the Alpha and Beta cell from day 7 to 12 of the cycle. The $\text{PGF}_{2\alpha}$ induced exocytosis of secretory granules from Beta cells on day 6 did not influence the length of the estrous cycle. We previously had shown (Fields et al., 1992) that the major depletion of oxytocin containing secretory granules occurred by day 14. This was prior to the reported rise in uterine oxytocin receptors between day 14 and 17 and the pulsing of uterine $\text{PGF}_{2\alpha}$ on days 16 and 17.

Our interpretation is there may exist a differential release of secretory granules from the Alpha and Beta cell throughout the cycle and that the Alpha cell release of granules may be in response to $\text{PGF}_{2\alpha}$ spikes that occurs late in the cycle. It is this latter release of oxytocin that leads to luteal regression. The $\text{PGF}_{2\alpha}$ sensitive Beta cell may provide for the early (day 11 to 14) depletion of granules (Fields et al., 1991). The role of this early secreted oxytocin is not understood (Fields et al., 1992).

7. Literature Cited

1. Bryant-Greenwood, G.D. Relaxin as a new hormone. *Endocr. Rev.*; 1982; 3: 62-90.
2. Choli, T.; Kapp, U.; Wittmann-Liebold, B. Blotting of proteins onto immobilon membranes: In situ characterization and comparison with high performance liquid chromatography. *J. Chromatogr.*; 1989; 476: 59-72.
3. Eckerskorn, C.; Lottspeich, F. Combination of two-dimensional gel electrophoresis with microsequencing and amino acid composition analysis: Improvement of speed and sensitivity in protein characterization. *Electrophoresis*; 1990; 11: 554-561.
4. Fehr, S.; Ivell, R.; Koll, R.; Schams, D.; Fields, M.; Richter, D. Expression of the oxytocin gene in the large cells of the bovine corpus luteum. *FEBS Lett.*; 1987; 210: 45-50.
5. Fields, M.J.; Barros, C.M.; Chang, S-M; Stallings-Mann, M.; Fields, P.A. In vivo effect of PGF2a on bovine large luteal cell secretory granules and their content, oxytocin and neurophysin. *Biol. Reprod.*; 1991; 44(Suppl 1): 100 (Abstr.).
6. Fields, M.J.; Dubois, W.; Fields, P.A. Dynamic features of luteal secretory granules: Ultrastructural changes during the course of pregnancy in the cow. *Endocrinology*; 1985; 117: 1675-1682.
7. Fields, M.J.; Dubois, W.; Ball, B.A.; Drost, M.; Fields, P.A. The effect of prostaglandin F2a on mitochondrial electron dense inclusions and secretory granules of the bovine large cell during late pregnancy. *Adv. Exp. Med. Biol.*; 1987; 219: 677-681.
8. Fields, M.J.; Dubois, W.; Brackett, K.H.; Faulkner, R.F.; Ball, B.A.; Martin, J.M.; Drost, M.; Fields, P.A. In-vivo effect of prostaglandin F2a treatment on secretory granules in the corpus luteum of the late pregnant cow. *J. Reprod. Fertil.*; 1989; Suppl. 37: 215-223.
9. Fields, M.J.; Fields, P.A.; Castro-Hernandez, A.; Larkin, L.H. Evidence for relaxin in corpora lutea of late pregnant cows. *Endocrinology*; 1980; 107: 869-876.
10. Fields, M.J.; Watkins, W.B.; Barros, C.M.; Fields, P.A. Characterization of large luteal cells and their secretory granules throughout the estrous cycle of the cow. *Biol. Reprod.*; 1992; (In Press).
11. Fields, P.A.; Fields, M.J. Ultrastructural localization of relaxin in the corpus luteum of the nonpregnant, pseudopregnant, and pregnant pig. *Biol. Reprod.*; 1985; 32: 1169-1179.
12. Fimmel, S.; Choi, T.; Dencher, N.A.; Eueldt, G.; Wittmann-Liebold, B. Topography of surface-exposed amino acids in the membrane protein bacteriorhodopsin determined by proteolysis and micro-sequencing. *Biochem. Biophys. Acta.*; 1989; 978: 231-240.
13. Fontana, A.; Gross, E. *Practical Protein Chemistry*. Darbre, A.; 1986: 94-95.
14. Fuchs, A.R.; Behrens, O.; Helmer, H.; Liu, H.C.; Barros, C.M.; Fields, M.J. Endometrial and myometrial OT and AVP receptors during the estrous cycle and early pregnancy. *Ann. Meet. Soc. Gynecol. Invest.*; 1990b: 127 (Abstr.).

15. Fuchs, A.R.; Behrens, O.; Helmer, H.; Ivanisevic, M.; Grifo, J.; Barros, C.M.; Fields, M.J. Ovaries and corpora lutea have receptors for oxytocin (OT) and vasopressin (AVP). 37th Ann. Meet. Soc. Gynecol. Invest.; 1990d; 300 (Abstr.).
16. Fuchs, A.R.; Behrens, O.; Helmer, H.; Fields, M.J.; Barros, C.M. Oxytocin and vasopressin receptors in cow uterus. Uterus. Proc. Inter. Conf. Neurohyp., Copenhagen; 1989 (Abstr.).
17. Fuchs, A.R.; Behrens, O.; Helmer, H.; Liu, H.; Barros, C.M.; Fields, M.J. Oxytocin and vasopressin receptors in bovine endometrium and myometrium during the estrous cycle and early pregnancy. Endocrinology; 1990c; 127: 629-636.
18. Fuchs, A.R.; Liu, H.C.; Nevarro, D.; Chang, S-M.; Stallings-Mann, M.; Fields, M.J. Preimplantation bovine conceptus suppresses endometrial oxytocin receptors in both IPSI and contralateral horn. Biol. Reprod.; 1991b; 44(Suppl. 1): 159 (Abstr.).
19. Fuchs, A.R.; Behrens, O.; Helmer, H.; Barros, C.M.; Fields, M.J. Oxytocin and vasopressin receptors in cow uterus. Proc. 4th Int. Conf. Neurohypophysis. New Aspects of Morphology, Function and Regulation, Oxford University Press; 1990a: 142-144.
20. Fuchs, A.R.; Behrens, O.; Helmer, H.; Vangsted, A.; Ivanisevic, M.; Grifo, J.; Barros, C.; Fields, M.J. Oxytocin and vasopressin binding sites in human and bovine ovaries. Am. J. Obstet. Gynecol.; 1991a; 163: 1961-1967.
21. Gast, M.J. Characterization of preprorelaxin by tryptic digestion and inhibition of its conversion to relaxin by amino acid analogs. J. Biol. Chem.; 1983; 258: 9001-9004.
22. Gast, M.J. Studies on luteal generation and processing of the high molecular weight relaxin precursor. Ann. N.Y. Acad. Sci.; 1982; 380: 111-125.
23. Gast, M.J.; Mercado-Simmen, R.; Niall, H.; Boime, I. Cell-free synthesis of a high molecular weight relaxin-related protein. Ann. N.Y. Acad. Sci.; 1980; 343: 148-154.
24. Giulian, G.G.; Shanahan, M.F.; Graham, J.M.; Moss, R.L. Resolution of low molecular weight polypeptides in a non-urea sodium dodecyl sulfate slab polyacrylamide gel system. Fed. Proc.; 1985; 44: 686.
25. Gurusinge, A.; Ryan, P.A.; Davis, P.F. Ready identification of disulfide-bonded polypeptides by one-dimensional sodium dodecyl sulfate-polyacrylamide slab gel electrophoresis. Electrophoresis; 1986; 7: 96-98.
26. Haley, J.; Hudson, P.; Scanlon, D.; John, M.; Cronk, M.; Shine, J.; Tregear, G.; Niall, H. Porcine relaxin: Molecular cloning and cDNA structure. DNA; 1982; 1: 155-162.
27. Hudson, P.; Haley, J.; Cronk, M.; Shine, J.; Niall, H. Molecular cloning and characterization of cDNA sequences coding for rat relaxin. Nature; 1981; 291: 127-131.
28. Ivell, R.; Brackett, K.H.; Fields, M.J.; Richter, D. Ovulation triggers oxytocin gene expression in the bovine ovary. FEBS Lett.; 1985; 190: 263-267.
29. Ivell, R.; Richter, D. The gene for the hypothalamic peptide hormone oxytocin is highly expressed in the bovine corpus luteum: Biosynthesis, structure and sequence analysis. EMBO J.; 1984; 3: 2351-2354.

30. Jassen Biotech. Guidelines for the use of auro probe BL plus and inten SE BL reagents for immunogold silver staining in dot and blot immune overlay assays.; 1984.
31. Jenner, L.J.; Parkinson, T.J.; Lamming, G.E. Uterine oxytocin receptors in cyclic and pregnant cows. *J. Reprod. Fertil.*; 1991; 91: 49-58.
32. Kemp, B.E.; Niall, H.D. Relaxin. *Vitam. Horm.*; 1984; 41: 79-115.
33. Kennedy, T.E.; Gawinowicz, M.A.; Barzilai, A.; Kandel, E.R.; Sweatt, J.D. Sequencing of proteins from two-dimensional gels by using in situ digestion and transfer of peptides to polyvinylidene difluoride membranes application to proteins associated with sensitization in Aplysia. *Proc. Natl. Acad. Sci.*; 1988; 85: 7008-7012.
34. Lundbald, R.L.; Noyes, C.M. The modification of cystine-cleavage of disulfide bonds. In: *CRC Chemical Reagents for Protein Modification*. Boca Raton, FL: CRC Press; 1984: 95-98.
35. Matsudaira, P. Sequence from picomole quantities of proteins electrophoretically onto polyvinylidene difluoride membranes. *J. Bio. Chem.*; 1987; 262: 10035-10038.
36. Meyer, H.H.D.; Mittermeier, Th.; Schams, D. Dynamics of oxytocin, estrogen and progesterin receptors in the bovine endometrium during the estrous cycle. *Acta Endocrinol.*; 1988; 118: 96-104.
37. Musah, A.I.; Schwabe, C.; Anderson, L.L. Acute decrease in progesterone and increase in estrogen secretion caused by relaxin during late pregnancy in beef heifers. *Endocrinology*; 1987a; 120: 317-324.
38. Musah, A.I.; Schwabe, C.; Willham, R.L.; Anderson, L.L. Dystocia, pelvic and cervical dilation in beef heifers after induction of parturition with relaxin combined with cloprostenol or dexamethasone. *Anim. Reprod. Sci.*; 1988; 16: 237-248.
39. Musah, A.I.; Schwabe, C.; Willham, R.L.; Anderson, L.L. Induction of parturition, progesterone secretion, and delivery of placenta in beef heifers given relaxin with cloprostenol or dexamethasone. *Biol. Reprod.*; 1987b; 37: 797-803.
40. Musah, A.I.; Schwabe, C.; Willham, R.L.; Anderson, L.L. Pelvic development as affected by relaxin in three genetically selected frame sizes of beef heifers. *Biol. Reprod.*; 1986b; 34: 363-369.
41. Musah, A.I.; Schwabe, C.; Willham, R.L.; Anderson, L.L. Relaxin on induction of parturition in beef heifers. *Endocrinology*; 1986a; 118: 1476-1482.
42. Musah, A.I.; Schwabe, C.; Anderson, L.L. Relaxin regulates oxytocin secretion in late-pregnant beef heifers. *Proc. Soc. Exp. Biol. Med.*; 1989; 191: 124-129.
43. Musah, A.H.I.; Schwabe, C.; Anderson, L.L. Relaxin and prostaglandin on oxytocin secretion from bovine luteal cells during different stages of gestation. *Acta Endocrinol.*; 1990; 122: 396-402.
44. O'Farrell. High resolution two-dimensional electrophoresis of proteins. *J. Bio. Chem.*; 1975; 250: 4007-4021.
45. Paccamonti, D. Clinical applications of relaxin in cattle. Thesis, University of Florida, Gainesville; 1988: 1-211.

46. Paccamonti, D.; Barros, C.; Drost, M.; Messine, O.; Fields, M.J. Response to nonpregnant and periparturient cows to porcine relaxin. *Biol. Reprod.*; 1989; 40(Suppl. 1): 70 (Abstr.).
47. Paccamonti, D.L.; Chang, S.T.; Dubois, W.; Barros, C.M.; Drost, M.; Wilcox, C.J.; Fields, M.J. Circulating concentrations of porcine relaxin in cows: Evaluation of vehicles and route of administration. *Theriogenology*; 1991; 35: 1131-1146.
48. Perezgrovas, R.; Anderson, L.L. Effect of porcine relaxin on cervical dilatation, pelvic area and parturition in beef heifers. *Biol. Reprod.*; 1982; 26: 765-776.
49. Pollard, J.W. Two-dimensional polyacrylamide gel electrophoresis of proteins. In: Walker JM, *Methods in Molecular Biology. 1. Proteins*. Clifton, NJ: Humana Press; 1984: 81-96.
50. Porter, D.G. The myometrium and the relaxin enigma. *Anim. Reprod. Sci.*; 1979; 2: 77-96.
51. Randel, R.D.; Musah, A.I.; Anderson, L.L. Temporal relationship between peripheral plasma relaxin, estradiol-17 β and progesterone in Brahman cows near parturition. *J. Anim. Sci.*; 1988; 66 (Suppl 1): 64 (Abstr.).
52. Renegar, R.H.; Larkin, L.H. Relaxin concentrations in endometrial, placental and ovarian tissues and in sera from ewes during middle and late pregnancy. *Biol. Reprod.*; 1985; 32: 840-847.
53. Robertson, E.F.; Dannell, H.K.; Malloy, P.J.; Reeves, H.C. Rapid isoelectric focusing in a vertical polyacrylamide minigel system. *Anal. Biochem.*; 1987; 167: 1290-294.
54. Sanger, F. Fractionation of oxidized insulin. *Biochem. J.*; 1949; 44: 126-128.
55. Schagger, H.; von Jagow, G. Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal. Biochem.*; 1987; 166: 368-379.
56. Schallenberger, E.; Schams, D.; Bullermann, B.; Walters, D.L. Pulsatile secretion of gonadotrophins, ovarian steroids and ovarian oxytocin during prostaglandin-induced regression of the corpus luteum in the cow. *J. Reprod. Fertil.*; 1984; 71: 493-501.
57. Schwabe, C.; McDonald, J.K. Demonstration of a pyroglutamyl residue at the n terminus of the B-chain of porcine relaxin. *Biochem. Biophys. Res. Commun.*; 1977; 74: 1501-1504.
58. Scott, M.G.; Crimmins, D.L.; McCourt, D.W.; Tarrand, J.J.; Eyerman, M.C.; Nahm, M.H. A simple in situ cyanogen bromide cleavage method to obtain internal amino acid sequence of proteins electroblotted to polyvinylidene difluoride membranes. *Biochem. Biophys. Res. Commun.*; 1988; 155: 1353.
59. Shemesh, M.; Hansel, W.; Strauss III, J.F. Bovine placentomes contain factors which decrease progesterone secretion. *Biol. Reprod.*; 1983; 29: 856-862.
60. Shemesh, M.; Hansel, W.; Strauss III, J.F. Calcium-dependent, cyclic nucleotide-independent steroidogenesis in the bovine placenta. *Proc. Natl. Acad. Sci. U.S.A.*; 1984; 81: 6403-6407.
61. Shemesh, M.; Hansel, W.; Strauss III, J.F. Modulation of bovine placental prostaglandin synthesis by an endogenous inhibitor. *Endocrinology*; 1984; 115: 1401-1405.

62. Sherwood, O.D. Relaxin. The Physiology of Reproduction. Knobil, E.; Neill, J.; Raven Press, New York, New York; 1988: 585-670.
63. Sherwood, O.D.; Ornekovic, V.E. Development of a homologous radioimmunoassay for rat relaxin. Endocrinology; 1979; 104: 893-897.
64. Short Jr., E C; Geisert, R D; Helmer, S D; Zavy, M T; Fulton, R W. Expression of antiviral activity and induction of 2',5'-oligoadenylate synthase by conceptus secretory proteins enriched in bovine trophoblast protein-1. Biol. Reprod.; 1991; 44: 261-268.
65. Soloff, M.S.; Fields, M.J. Changes in uterine oxytocin receptor concentrations throughout the estrous cycle of the cow. Biol. Reprod.; 1989; 40: 283-287.
66. Stallings-Mann, M.L. A Relaxin-Like Peptide in the Bovine Placenta. Thesis, University of Florida; 1991: 1-191.
67. Steinetz, B.G.; O'Byrne, E.M.; Butler, M.C.; Hickman, L.B. Hormonal regulation of the connective tissue of the symphysis pubis. Biology of Relaxin and Its Role in the Human. Bigazzi, M.; Greenwood, F.C.; Gasparri, F.; Excerpta Medica, Amsterdam, The Netherlands; 1983: 71-92.
68. Stewart, D.R.; Papkoff, H. Purification and characterization of equine relaxin. Endocrinology; 1986; 119: 1093-1099.
69. Towbin, H.; Staehelin, T.; Gordon, J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. Proc. Natl. Acad. Sci. U.S.A.; 1979; 76: 4350-4355.
70. Wada, H.; Yuhara, M. Concentration of relaxin in the blood serum of pregnant cow and cow with ovarian cyst. Proc. Silver Jubilee, Kyoto University; 1961: 61-66.
71. Wada, H.; Yuhara, M. Extraction of relaxin from sow ovary and cow tissues. Jpn. J. Zootech. Sci.; 1960; 31: 237-242.
72. Walsh, M.; McDongal, J.; Witmann-Liebold, B. Extended n-terminal sequence of proteins of archaebacterial ribosomes blotted from two-dimensional gels onto glass fiber and poly (vinylidene difluoride) membrane. Biochemistry; 1988; 27: 6867-6876.
73. Wathes, D.C.; Rees, J.M.; Porter, D.G. Identification of relaxin in the placenta of the ewe. J. Reprod. Fertil.; 1988; 84: 247-257.

F. Description of Cooperation

There has been an excellent cooperative spirit between the laboratories of Dr. Michael J. Fields at the University of Florida and Dr. Mordechai Shemesh at the Kimron Veterinary Institute. The first year of the grant Dr. Michael J. Fields participated in the BARD sponsored workshop entitled "Maternal Recognition of Pregnancy" held March 20-24 in Jerusalem in which Dr. Mordechai Shemesh was the organizer of the conference. The first visit provided an opportunity for reviewing the objectives and planned experiments of this grant. In addition a visit with Dr. Larry Shores at the Kimron Veterinary Institute led to a collaborative effort between scientist at Florida and Israel to evaluate a major Florida livestock forage, white dutch clover, for phytoestrogens. The preliminary results were positive and may provide valuable information as to the profound positive effect this clover has on the rebreeding of lactating cattle.

During the course of this grant data from experiments were exchanged, protocols modified and results reevaluated in view of additional experiments. Purified porcine relaxin, antibodies to porcine relaxin, tissue samples, plasma and extracts were exchanged. Both investigators participated in the 21st through 24th annual meeting of the Society for the Study of Reproduction in the United States. It was at these meetings that there was an extensive exchange of ideas and data and planning of experiments was accomplished.

In the fall of 1990 Dr. Shemesh joined us at the University of Florida for a series of studies on the in vitro effect of relaxin on uterine and placental steroidogenesis and prostaglandin synthesis in the pregnant cow. His visit was an informative one and the experiments conducted were successful. Prior to that visit Dr. Larry Shore from the Kimron Veterinary Institute made two trips to Gainesville in which information and experimental protocols were exchanged.

In addition a very successful collaborative research effort was maintained between the laboratory of Dr. Michael Fields and Dr. Phillip Fields at the University of South Alabama (secretory

granules) and Dr. Anna-Riitta Fuchs at the Cornell Medical School (oxytocin receptors). A collaborative study was also conducted with Dr. Michael Diskin, Agricultural and Food Development Authority at Galway, Ireland evaluating induction of parturition with relaxin when administered with glucocorticoids. These latter collaborative efforts involved numerous visits and exchange of antibodies, porcine relaxin and tissues for study between the respective laboratories.

G. Evaluation and Benefit to Agriculture

One of the more pressing reasons for understanding relaxin, the hormone of parturition, is the cattle industry's introduction of management systems that have led to an increase in problems at the time of calving. These systems have involved the use of bulls that have a propensity for siring calves with a higher birth weight. Furthermore, the practice of breeding heifers to calve earlier at a time when their skeletal frames are still developing, has led to an increase in the frequency of dystocia. In some locations these changes in breeding programs has resulted in a 5 to 15% loss in potential calf crop due to dystocia and early post-natal death. A 5% death loss for 5.1 million heifers in the United States is a loss of 250,000 calves. At \$400 per calf, the loss would total \$100 million. In the dairy industry the problem is an additional one of retained placenta that reportedly affects 10-14% of dairy cows for an estimated loss of \$15 million.

However, relaxin is an enigma in the ruminant. The cow is responsive to porcine relaxin but little is known about the endogenous form of bovine relaxin. Although, the nucleotide and amino acid sequence of numerous species of relaxin have been determined none of this information is available for the ruminants. The application of the most recent tools of molecular biology have not been successful to date.

Relaxin was immunohistochemically localized in the syncytiotrophoblast and subsequently shown in isolates from the cotyledon. Isolates from both the corpus luteum and placenta gave a linear dose response in the homologous porcine relaxin radioimmunoassay. This relaxin was shown to differ from porcine relaxin when eluted from a CM-cellulose column. Attempts to increase the recoverable yields of bovine relaxin were not successful. Although microtechniques for successfully separating the A and B chain of porcine relaxin and determining their amino acid sequence were developed, insufficient quantities of bovine relaxin preempted their application. These techniques were, however, successfully utilized to show that the high molecular weight porcine immunoreactive relaxin-like

peptides were in fact processed prohormone. We are encouraged that most recently we have shown with western blots a 17,000 Dalton immunoreactive relaxin from the bovine placenta that could be a form of the bovine prohormone.

These findings indicate there may be two sources of relaxin in the cow, i.e., the corpus luteum and the placenta. There may be a systemic role for luteal relaxin and a paracrine role for placental relaxin. Endocrinology is replete with many redundant systems. Relaxin in the cow produced by both the placenta and corpus luteum could be another example of one. Luteal relaxin would be expected to have a systemic role since it would be expected to be released from secretory granules to circulate in the vascular system. Removal of the corpus luteum during pregnancy results in maintenance of pregnancy in many instances, however, the pregnancies are characterized by dystocia, retained fetal membranes and in many instances death of the calf. The placental production of relaxin would allow the birth of live calves. This is in contrast to the pig in which there is high circulating relaxin of luteal origin, no evidence for placental relaxin and an obligatory need for the corpus luteum (relaxin) if live piglets are to be born.

We showed in addition to an immunoreactive placental relaxin that the cow responds to exogenous administration of porcine relaxin. Relaxin was shown to have potential to induce parturition and reduce the incidence of retained placenta when administered along with glucocorticoids to the cow approaching the end of a normal gestation. In an attempt to elucidate the cellular mechanisms by which this action occurs we evaluated the effect of relaxin on steroid metabolism and prostaglandin by the uterine caruncle and fetal membranes. There was not demonstrate a consistent direct effect but we can not rule out an indirect effect via other factors since relaxin did potentiate the effect of incubating MIX with caruncular and cotyledonary tissues. It is through understanding the underlying mechanisms of action of relaxin that more effective systems can be developed to possibly reduce dystocia and retained placenta in the cow.

One of the reported roles of relaxin is a lutelytic effect when utilized to induce parturition. In addition relaxin has been shown to induce the release of both hypothalamic (in vivo) and luteal (in vitro) release of oxytocin indicating an interrelationship between relaxin and oxytocin. Based on this the oxytocin receptor was characterized throughout the reproductive cycle of the cow. Endometrial oxytocin receptor levels varied significantly during the cycle being lowest on days 7, 10 and 14, rose significantly on day 17 and peaked on day 21. In the pregnant animal the oxytocin receptors were also low on day 7, 10 and 14 but on day 17 and 21 the endometrial receptor concentrations were significantly lower than seen in the nonpregnant cow. We conclude that contributing to the sustainment of pregnancy is the conceptus suppression of uterine oxytocin receptors. This study contributed to the fundamental understanding of the maternal recognition paradigm of pregnancy. This type of basic information will eventually be utilized for greater effective control of the estrous cycle and maintenance of pregnancy. It is at this early preimplantation stage of pregnancy that up to 25% of the pregnancies are lost.

The role of oxytocin during parturition and particularly during pregnancy is not understood. Our approach was one of characterizing the oxytocin receptor throughout pregnancy in an attempt to define the target tissues of oxytocin and stage of pregnancy for time of biological action of this hormone. The surprising finding was the mid-pregnancy rise in oxytocin receptors in both the myometrium and endometrium. However, only the receptors of the endometrium peaked just prior to parturition. In addition the concentration of the endometrial receptors were much higher than for the myometrium. Since oxytocin stimulates the release of $\text{PGF}_{2\alpha}$ from the nonpregnant bovine endometrium there may be a similar cascade operating during pregnancy. The failure of myometrial and endometrial concentrations of oxytocin receptor to follow a parallel course suggest a somewhat differential regulation of oxytocin receptor formation in the two uterine tissues.

In addition concentrations of oxytocin receptors were established throughout pregnancy in the amnion and chorio-allantoic membranes. Concentrations of receptors were low throughout most of gestation rising abruptly at time of labor. There were no differences in the affinity of the receptors between these membranes and the endometrium. These findings coupled with those for relaxin indicate both relaxin and oxytocin play an integral role in bovine parturition.

As mentioned above relaxin was also shown to be present in the corpus luteum. Peptide hormones are stored in secretory granules for release at a critical time during the reproductive cycle. We have shown that one of the factors initiating this release is prostaglandin $F_{2\alpha}$. During the estrous cycle these granules were shown to contain oxytocin and neurophysin. The granules are formed early in the estrous cycle as a single cluster. Once the granules are depleted, via exogenous $PGF_{2\alpha}$ the cell is incapable of generating a second population of secretory granules.

A model of $PGF_{2\alpha}$ depletion of secretory granules revealed two populations of large luteal cells which will be referred to as Alpha and Beta. Beta large luteal cells (40-46 μm diameter) contained numerous mitochondria and a more apparent smooth endoplasmic reticulum than small Alpha large luteal cells (32-39 μm). In addition, Beta large luteal cells were more sensitive to the $PGF_{2\alpha}$ as evidence.

Our interpretation is there may exist a differential release of secretory granules from the Alpha and Beta cell throughout the cycle and that the Alpha cell release of granules may be in response to $PGF_{2\alpha}$ spikes that occurs late in the cycle. It is this latter release of oxytocin that leads to luteal regression. The $PGF_{2\alpha}$ sensitive Beta cell may provide for the early (day 11 to 14) depletion of granules. The role of the early secreted oxytocin is not understood but we believe it plays an important role in the reproductive cycle.

During early pregnancy prior to day 40 there is no available information on secretory granules. During this period there is a transition of a corpus luteum going from one believed to contain

oxytocin granules to one devoid of oxytocin. The content of granules of pregnancy after day 40 is unknown. Since relaxin is the only hormone identified in luteal secretory granules of pregnancy of other species we believe these granules also contain relaxin in the cow.

In conclusion relaxin and oxytocin are important hormones of pregnancy. The findings of this series of studies contributes to the basic understanding of the presence of these hormones, their mechanism of action and their function of in the cow. This information will be built on leading to an improvement of management and thus reduction in losses that occur during the periparturient period in the cow.

H. List of Publications

1. Fields, M.J.; Barros, C.M.; Chang, S.-M.; Stallings-Mann, M.; Fields, P.A. In vivo effect of PGF_{2a} on bovine large luteal cell secretory granules and their content, oxytocin and neurophysin. *Biol. Reprod.*; 1991; 44(Suppl 1): 100 (Abstr.).
2. Fields, M.J.; Watkins, W.B.; Barros, C.M.; Fields, P.A. Characterization of large luteal cells and their secretory granules throughout the estrous cycle of the cow. *Biol. Reprod.*; 1992; (In Press).
3. Fuchs, A.R.; Behrens, D.; Helmer, H.; Liu, H.C.; Barros, C.M.; Fields, M.J. Endometrial and myometrial OT and AVP receptors during the estrous cycle and early pregnancy. *Ann. Meet. Soc. Gynecol. Invest.*; 1990b; 127 (Abstr.).
4. Fuchs, A.R.; Behrens, D.; Helmer, H.; Invanisevic, M.; Grifo, J.; Barros, C.M.; Fields, M.J. Ovaries and corpora lutea have receptors for oxytocin (OT) and vasopressin (AVP). 37th Ann. Meet. Soc. Gynecol. Invest.; 1990d; 300 (Abstr.).
5. Fuchs, A.R.; Behrens, D.; Helmer, H.; Fields, M.J.; Barros, C.M. Oxytocin and vasopressin receptors in cow uterus. *Uterus. Proc. Inter. Conf. Neurohyp.*, Copenhagen; 1989 (Abstr.).
6. Fuchs, A.R.; Behrens, D.; Helmer, H.; Liu, H.; Barros, C.M.; Fields, M.J. Oxytocin and vasopressin receptors in bovine endometrium and myometrium during the estrous cycle and early pregnancy. *Endocrinology*; 1990c; 127: 629-636.

7. Fuchs, A.R.; Liu, H.C.; Nevarro, D.; Chang, S-M.; Stallings-Mann, M.; Fields, M.J. Preimplantation bovine conceptus suppresses endometrial oxytocin receptors in both IPSI and contralateral horn. Biol. Reprod.; 1991b; 44(Suppl. 1): 159 (Abstr.).
8. Fuchs, A-R.; Behrens, D.; Helmer, H.; Barros, C.M.; Fields, M.J. Oxytocin and vasopressin receptors in cow uterus. Proc. 4th Int. Conf. Neurohypophysis. New Aspects of Morphology, Function and Regulation, Oxford University Press; 1990a: 142-144.
9. Fuchs, A-R.; Behrens, D.; Helmer, H.; Vangsted, A.; Ivanisevic, M.; Grifo, J.; Barros, C.; Fields, M.J. Oxytocin and vasopressin binding sites in human and bovine ovaries. Am. J. Obstet. Gynecol; 1991a; 163: 1961-1967.
10. Paccamonti, D.; Barros, C.; Drost, M.; Messine, D.; Fields, M.J. Response to nonpregnant and periparturient cows to porcine relaxin. Biol. Reprod.; 1989; 40(Suppl. 1): 70 (Abstr.).
11. Paccamonti, D.L.; Chang, S.T.; Dubois, W.; Barros, C.M.; Drost, M.; Wilcox, C.J.; Fields, M.J. Circulating concentrations of porcine relaxin in cows: Evaluation of vehicles and route of administration. Theriogenology; 1991; 35: 1131-1146.
12. Stallings-Mann, M.L. A Relaxin-Like Peptide in the Bovine Placenta. Thesis, University of Florida; 1991: 1-191.