Winter chickpea; towards a new winter pulse for the semiarid Pacific Northwest and wider adaptation in the Mediterranean basin

Abbo, S. The Hebrew University of Jerusalem

Zhang, H. Texas A&M Research Foundation TX

Coyne, C. USDA, Agricultural Research Service WA

Shtienberg, D. Agricultural Research Organization

Sherman, A. Agricultural Research Organization

Vandemark, G.J. USDA, Agricultural Research Service WA

Project award year: 2011

Three year research project

Abstract

Original objectives: [a] Screen an array of chickpea and wild annual *Cicer* germplasm for winter survival. [b] Genetic analysis of winter hardiness in domesticated x wild chickpea crosses. [c] Genetic analysis of vernalization response in domesticated x wild chickpea crosses. [d] Digital expression analysis of a core selection of breeding and germplasm lines of chickpea that differ in winter hardiness and vernalization. [e] Identification of the genes involved in the chickpea winter hardiness and vernalization and construction of gene network controlling these traits. [f] Assessing the phenotypic and genetic correlations between winter hardiness, vernalization response and Ascochyta blight response in chickpea. The complexity of the vernalization response and the inefficiency of our selection experiments (below) required quitting the work on ascochyta response in the framework of this project.

Background to the subject: Since its introduction to the Palouse region of WA and Idaho, and the northern Great Plains, chickpea has been a spring rotation legume due to lack of winter hardiness. The short growing season of spring chickpea limits its grain yield and leaves relatively little stubble residue for combating soil erosion. In Israel, chilling temperatures limit pod setting in early springs and narrow the effective reproductive time window of the crop. Winter hardiness and vernalization response of chickpea alleles were lost due to a series of evolutionary bottlenecks; however, such alleles are prevalent in its wild progenitor's genepool.

Major conclusions, solutions, achievements: It appears that both vernalization response and winter hardiness are polygenic traits in the wild-domesticated chickpea genepool. The main conclusion from the fieldwork in Israel is that selection of domesticated winter hardy and vernalization responsive types should be conducted in late flowering and late maturity backgrounds to minimize interference by daylength and temperature response alleles (see our *Plant Breeding* paper on the subject). The main conclusion from the US winter-hardiness studies is that excellent lines have been identified for germplasm release and continued genetic study. Several of the lines have good seed size and growth habit that will be useful for introgressing winter-hardiness into current chickpea cultivars to develop releases for autumn sowing.

We sequenced the transcriptomes and profiled the expression of genes in 87 samples. Differential expression analysis identified a total of 2,452 differentially expressed genes (DEGs) between vernalized plants and control plants, of which 287 were shared between two or more *Cicer* species studied. We cloned 498 genes controlling vernalization, named *CVRN* genes. Each of the *CVRN* genes contributes to flowering date advance (FDA) by 3.85% - 10.71%, but 413 (83%) other genes had negative effects on FDA, while only 83 (17%) had positive effects on FDA, when the plant is exposed to cold temperature. The cloned *CVRN* genes provide new toolkits and knowledge to develop chickpea cultivars that are suitable for autumn-sowing.

Scientific & agricultural implications: Unlike the winter cereals (barley, wheat) or pea, in which a single allelic change may induce a switch from winter to spring habit, we were unable to find any evidence for such major gene action in chickpea. In agricultural terms this means that an alternative strategy must be employed in order to isolate late flowering – ascochyta resistant (winter types) domesticated forms to enable autumn sowing of chickpea in the US Great Plains. An environment was identified in U.S. (eastern Washington) where autumn-sown chickpea production is possible using the levels of winter-hardiness discovered once back-crossed into advanced cultivated material with acceptable agronomic traits. The cloned *CVRN* genes and identified gene networks significantly advance our understanding of molecular mechanisms underlying plant vernalization in general, and chickpea in particular, and provide a new toolkit for switching chickpea from a spring-sowing to autumn-sowing crop.

Summary Sheet
PubType IS only Joint US only
Reviewed 0 2 2

Description & details of cooperation

The field experiments conducted in Israel and in Pullman WA served for both phenotypic assessments of the target traits (flowering vernalization response and winter hardiness, respectively) as well as for RNA extraction for the expression analyses and gene tagging by the Texas A&M group. Therefore, the molecular work done in Texas A&M was the hub of this project and we look forward to the completion of the RNA-seq work which is likely to provide new and fresh insight into the genetic control of these two important adaptive traits in chickpea.

The field experiments in Israel and in Pullman WA provided the raw material for the detailed gene sequence and expression analyses conducted in Texas A&M (Zhang lab). The last shipment of RNA from the last field nursery of winter 2015 is due in December 2015. We look forward to the completion of the RNA-seq work which is likely to provide new and fresh insight into the genetic control of these two important adaptive traits in chickpea on top of the already identified *CVER* genes and networks. Shahal Abbo visited both in Texas A&M University and in Pullman WA (in March and July 2013, respectively) and presented seminars on his work. In the meetings we discussed in details the future of this work as well as other options for collaborations on agronomic and genetic improvement of cool season food legumes (chickpea, pea). As a result of the successful work and accumulated data we are now beginning to formulate a future grant application for BARD that builds on the present work.

Major achievements

From our field work it became appears that both vernalization response and winter hardiness are traits controlled by many genes in the wild-domesticated chickpea genepool. The main conclusion from the fieldwork in Israel is that selection of domesticated winter hardy and vernalization responsive types should be conducted in late flowering and late maturity backgrounds to minimize interference by daylength and temperature response alleles (see our ***Plant Breeding*** paper on the subject). The main conclusion from the US winter-hardiness studies is that excellent lines have been identified for germplasm release and continued genetic study. Several of the lines have good seed size and growth habit that will be useful for introgressing winter-hardiness into current chickpea cultivars to develop releases for autumn sowing.

We have profiled the expression of genes in 87 samples. This analysis identified a total of 2,452 differentially expressed genes (DEGs) between vernalized plants and control plants, of which 287 were shared between two or more *Cicer* species studied. We cloned 498 genes controlling vernalization. The cloned *CVRN* genes provide new toolkits and knowledge to develop chickpea cultivars that are suitable for autumn-sowing.

Unlike the winter cereals (barley, wheat) or pea, in which a single genetic change may induce a switch from winter to spring habit, we were unable to find any evidence for such effects in chickpea. In agricultural terms this means that an alternative strategy must be employed in order to isolate late flowering – ascochyta resistant (winter types) domesticated forms to enable autumn sowing of chickpea in the US Great Plains.

An environment was identified in U.S. (eastern Washington) where autumn-sown chickpea production is possible using the levels of winter-hardiness discovered once back-crossed into advanced cultivated material with acceptable agronomic traits. The cloned *CVRN* genes and identified gene networks provide a new toolkit for switching chickpea from a spring-sowing to autumn-sowing crop.

Publications for Project IS-4427-11C

Status	Type	Authors	Title	Journal	Volume: Pages	Year	Country
Published	Reviewed	Zhang M, Yang Zhang, CF Scheuring, C-C Wu, JJ Dong & H-B Zhang	Preparation of megabase-sized DNA from a variety of organisms using the nuclei method for advanced genomics research	<i>Nature Protocols</i>	7 : 467- 478	2012	US only
Published	Reviewed	Zhang H-B, CF Scheuring, M Zhang, Y Zhang, C-C Wu, JJ Dong & Y Li van Oss R, Abbo S, Eshed R, Sherman A, Coyne CJ, Vandemark GJ, Zhang H-B, & Peleg Z	Construction of BIBAC and BAC libraries from a variety of organisms for advanced genomics research	<i>Nature Protocols</i>	7 : 479- 499	2012	US only
Published	Reviewed	Pinhasi-van Oss R, Sherman A, Zhang H-B, Vandemark G, C. Coyne C & Abbo S	Genetic relationship in Cicer sp. expose evidence for gene flow between the cultigen and its wild progenitor	<i>PLOS One</i>	10 : e01397- 89	2015	Joint
Accepted	Reviewed		Vernalization response of domesticated x wild chickpea progeny is subject to strong genotype by environment interaction	<i>Plant Breeding</i>	:	2015	Joint

Achievements

Genetic analysis of vernalization response (Rehovot & Bet-Dagan, Israel)

As stated in our original grant application, apart from a preliminary report suggesting that vernalization response in chickpea is a polygenic trait (Abbo et al. 2002), prior to the initiation of this project, no information on this trait was available to chickpea breeders and agronomists who are interested in introducing chickpea into high latitude growing regions as a winter crop. Our major achievements after four experimental growing seasons in Rehovot, Israel are as follows.

1] Confirmation of past reports concerning the polygenic nature of flowering vernalization response in chickpea. This conclusion has been also confirmed by cloning 498 *CVRN* genes involved in the process of vernalization in this research (see below). This trait is subject to strong genotype by environment interaction. This is in sharp contrast to the well-studied systems of the winter cereals. This part of the project is included in a forthcoming paper in the journal *Plant breeding* (Pinhasi-van Oss et al.).

2] Identification of several QTL affecting flowering time following vernalization in chickpea using a set of SNP markers [Hiremath et al. (2012) *Plant Biotech J* doi: 10.1111/j.1467-7652.2012.00710.x]. See table 1 below.

Table 1. Linkage groups (LGs) and marker locations of the QTL affecting flowering time following vernalization treatment in hybrid progeny population of chickpea derived from the cross cv Mekomit x Cr934. Percentage expected variation (PEV).

Vernalization response loci in F ₂			
	Nearest marker	PEV	LOD score
LG2	CKAM1428	0.079	4.3
LG3	CKAM0628	0.251	9.6
LG4	CKAM1455	0.079	4.9
LG7	TOG896873	0.082	4.9
Vernalization response loci in F ₃			
	Nearest marker	PEV	LOD
LG3	CKAM0628	0.290	6.1
LG4	CKAM1501	0.210	6.3

3] Our field work was conducted with two distinct genetic backgrounds (late vs. early flowering). The pattern of advance to flowering as reported in our *Plant Breeding* paper suggests that future work towards fine mapping, tight marker tagging and isolation of the vernalization response genes should be conducted in a late flowering background. This is imperative in order to minimize the co-segregation of day-length response and temperature response loci affecting flowering time. This is a very important message to chickpea breeders and agronomists because early flowering is an important trait in all chickpea growing environments and is subject to manipulation in many chickpea breeding programs worldwide.

Genetic analysis of winterhardiness (Pullman WA, USA)

Two populations of *C. arietinum* × *C. reticulatum* were tested in the field for winter-hardiness in one U.S. location in 2011-2012. One population (ICC 4958 × PI 498777; 130 lines) segregated for winter damage and was selected for further testing in eight environments at three locations using a randomized complete block design with three replications. Differential damage was recorded at four environments (years) from one location, with the line data exhibiting a bi-modal trend, strongest in 2012-2013 (Fig. 1), tending to more winter damage (death) in two of the

others years 2013-2014 and 2014-2015. This indicates a major gene/region is conferring the winter-hardiness from the *C. reticulatum* donor parent. Year to year correlations were positive and highly significant (Table 2).

Figure 1. Frequency histogram of the winter damage 2012-13 on the *C. arietinum* $\times$ *C. reticulatum* population (1= 50% leaf damage, 5=all plants dead).

Table 2. Correlation of winter damage in *C. arietinum* $\times$ *C. reticulatum* population. Spearman's correlation coefficient above the diagonal, Pearson's correlation coefficient below the diagonal.

Year	2011-2012	2012-2013	2013-2014	2014-2015
2011-2012		0.75***	0.62***	0.62***
2012-2013	0.74***		0.64***	0.60***
2013-2014	0.62***	0.58***		0.85***
2014-2015	0.64***	0.57***	0.92***	

*** significant at $P < 0.0001$

Winter-hardy and winter-susceptible lines from the first two years of field data were selected for study under controlled freeze conditions using a growth chamber. Lacking a published protocol for chickpea, a controlled freeze procedure was developed based on one used in wheat. The developed controlled freeze procedure was successful in confirming winter-hardiness and winter-sensitivity of inbred lines selected from the *C. arietinum* by *C. reticulatum* cross, plus the parents.

Field screening of autumn-sown chickpea was successful in determining winter-hardiness in a segregating population of recombinant inbred lines from a *C. arietinum* $\times$ *C. reticulatum* cross (manuscript in preparation). Superior winter-hardy introgression lines have been identified from the population. Release of winter-hardy germplasm is in progress for use in aggressive back-crossing programs based on the best winter-hardy lines discovered from four years of field study and under controlled freeze tests.

Identification of genes networks controlling vernalization response and winter hardiness (Texas A&M, USA)

We sequenced and profiled the transcriptomes of 87 samples, with 20 from the winter hardiness experiment (WSU, USA) and 67 from the vernalization experiment (Israel). We obtained 12 – 17 million 100 PE (paired end) clean reads for each sample. Together, we generated a total of >1,300 million 100 PE clean reads, with >98% of the clean reads having a quality of $\geq Q20$. Analysis of the sequences resulted in transcript unigenes from 38,731 – 62,118 for each sample, with an N50 size ranging from 1,346 bp to 1,752 bp. We also measured the expression profile of every gene expressed in the 87 samples.

Furthermore, we analyzed the sequence and expression data of the genes in two aspects. The first was differential expression analysis between vernalized plants and controls. From this analysis we identified a total of 2,452 differentially expressed genes (DEGs) between vernalized plants and control plants, of which 287 were shared between two or more *Cicer* species studied (Fig. 1). The second was cloning of the genes controlling chickpea vernalization using a novel genome-wide high-throughput gene and QTL cloning system (filed for a patent). From this analysis we cloned 498 genes controlling chickpea vernalization, named *CVRN* for *Cicer* vernalization genes. The *CVRN* genes each contribute to flowering date advance by 3.85% - 10.71%, but 413 (83%) of them had negative effects on FDA, while only 83 (17%) had positive effects on FDA, when the host plant is exposed to cold temperature. Systems analysis of the 498 cloned *CVRN* genes showed that vernalization is controlled by at least 498 genes and it is the consequence of balancing between the genes having positive and negative effects. We also found that genes controlling vernalization encode enzymes or proteins that function in a variety of biochemical reactions and multiple metabolic pathways. The *CVRN* genes are correlated in function, forming a single interaction network (Fig. 2). The structure of the network, including content and number of genes (nodes), gene action direction, and number and composition of interactions (edges), varied dramatically between vernalized and control plants and among lines with different responses to cold temperature, which suggests this work identified molecular determinants of chickpea vernalization. These results provide not only novel insights into the molecular mechanisms of plant vernalization and flowering, but also strategies and toolkits for enhanced manipulation of plant vernalization, such as switching the modern spring chickpea back to the autumn-sowing chickpea.

Figure 1 Venn diagram of the differentially expressed genes (DEGs) between cold-treated vernalized plants and their non-cold treated controls among four *Cicer* species.

Figure 2 The 498 *CVRN* genes controlling vernalization in *Cicer* and their co-expression interaction network. The gene network consists of 498 *CVRN* gene nodes, 16,921 interaction edges and 12 functional clusters.