8

356
87

77

BARD

FINAL REPORT

PROJECT NO. US-1362-87

**Hormonal and Developmental Control of Root
Initiation**

J.D. Cohen, E. Epstein

(9)

הספריה סמוכזית
למדעי החקלאות
בית דגן

1993

630.72
BAR/COH
2nd copy

Standard BARD Cover Page for Scientific Reports

Date: January 4, 1993

BARD
P.O. BOX 6
Bet Dagan, ISRAEL

BARD project No. US-1362-87C

Title

Hormonal and Developmental Control of Root InitiationInvestigators' Names
(Principal listed first)J.D. CohenE. Epstein

Investigators' Institutions

USDA-ARS BeltsvilleARO, The Volcani CenterProject's starting date: March 1, 1989Type of Report: 1st Annual 2nd Annual Final X

Signature
Principal Investigator

DB

Signature
Institution's Authorizing Official

630.72
BAR/COH

: 581.498 : 581.134 : 581.192.7 :
581.192.96 : 631.53

b. Table of Contents	1
c. Abstract	2
d. Original research proposal objectives as related to the research achieved.	3
e. Description of the research activities and results	3
Introduction:	3
Beltsville Agricultural Research Center	3
Results:	
Measurements of IAA and IBA in the same plant sample	3
Carrot embryogenesis	5
<i>Petunia</i> Organogenesis	7
Auxin biosynthesis	7
The Volcani Center	8
Mung bean studies	8
Petunia cell suspension cultures	8
Citrus leaf	9
Sweet cherry	9
Protea	9
Endo-(1,4)- β -glucanase gene expression during rooting	9-a
f. Description of Cooperation and Conclusions:	10
g. Evaluation of the research achievements with respect to the aims of the original research proposal.	10
h. Publications:	11
Appendix	

c. Abstract

The major accomplishments from this project were: 1) the development of a number of systems of easy-to-root and difficult-to-root cultivars for use for comparative biochemical studies of rooting; 2) the development of mass spectral techniques for the analysis of tryptophan, indolebutyric acid and indoleacetic acid in plant tissues; 3) use of this technology to study the changes and interrelationships between both auxins and tryptophan during rooting and micropropagation; and 4) demonstration that indoleacetic acid is produced by a new pathway not involving tryptophan as a direct precursor and that this pathway is under developmental control. These studies have changed the basic understanding of the dynamics and relationships of auxins during critical steps in plant development related to propagation technology. The results of these studies will be of interest to those involved in plant propagation, plant tissue culture and scientists involved in understanding the process of plant development.

d. Original research proposal objectives as related to the research achieved.

Our grant had four main goals: 1) continue the work on IAA and IBA metabolism; 2) begin to study the biology and biochemistry of organogenesis; 3) begin to do tissue localization studies on auxin metabolism during rooting; 4) work toward obtaining model systems using ER and DR cultivars or mutants to study organogenesis. Our finding of IBA as a natural product formed from IAA has added a new dimension to this work, represented in the progress from the Israeli group on this aspect. The Beltsville group focused its attention to the problem of organogenesis and embryogenesis as controlled by endogenous hormonal systems and on the route of IAA biosynthesis.

e. Description of the research activities and results:

Introduction:

Research funded under BARD US-1362-87C concerns the study of the hormonal and molecular signals involved in the induction of developmental events such as root formation, organogenesis, and embryogenesis that play a critical role in propagation of horticultural species. Several experimental systems were employed in work carried out in the US and in Israel. The US team worked primarily with carrot cell cultures, *Lemna* cultures, petunia explants and a mutant line of maize. In Israel, Dr. Epstein primarily studies citrus, maize and bean. Each of these experimental system has unique advantages that lend themselves to the studies undertaken. These aspects were discussed in more detail in the first annual report.

Beltsville Agricultural Research Center

Research funded by BARD was aimed at identification of the hormonal and molecular signals involved in the induction of organogenesis or embryogenesis in plant tissue systems. Several experimental systems are being studied in detail: the process of embryogenesis was studied in carrot cell cultures, the process of organogenesis was studied in petunia and the process of hormonal regulation was studied in maize. Each of these experimental systems has unique properties that lends itself to the investigations conducted. In particular, in carrot it is the removal of a single hormonal signal, the auxin in the medium, that initiated embryogenesis. In petunia, it is the addition of cytokinin that stimulates the process of organogenesis and the addition of auxin changes the pattern of such development. In maize, auxin biosynthesis was studied by utilization of a unique mutant, orange pericarp, that is a double recessive mutant blocked at the step of tryptophan synthase B, and is thus a complete tryptophan auxotroph.

Results:

Measurement of IAA and IBA in the same plant sample

The role of auxins as the primary group of hormones involved in adventitious root formation has been well established (Davis *et al.*, 1988). Although IAA has been considered the main endogenous auxin responsible for adventitious root formation, both research workers and commercial propagators have preferentially used IBA to induce rooting because of its greater efficacy. Until recently, IBA was considered to be only a synthetic auxin analog. A few early reports suggested - and more recent work has demonstrated - that IBA occurs naturally. For example, IBA was shown to occur in carrot in normal roots and in "hairy roots" resulting from transformation by *Agrobacterium rhizogenes* (Epstein *et al.*, 1991), as well as in maize (Epstein *et al.*, 1989) and in genetic tumors of *Nicotiana* inter species crosses (Chen, 1987). The role of auxin in root induction by *A. rhizogenes* has not been studied in detail and IAA levels before and after transformation have been measured directly only in carrot (Epstein *et al.*, 1991; Hwang *et al.*, 1986). The finding that IBA also occurs in transformed carrot tissue has prompted

us to study the relationship of IAA and IBA in root induction by *A. rhizogenes* in this system.

Synthesis of ^{13}C -indole-[ring 2]-3-butyric acid: The synthesis of ^{13}C -indole-[ring 2]-3-butyric acid followed the same general procedure described by Cohen and Schulze (1981) for the synthesis of the ^{14}C -labeled compound. Twelve mg of indole-2- ^{13}C (Cambridge Isotope Laboratories, CLM-1863) were placed in a 1 ml size (4 ml capacity) freeze-drying tear bulb (A.H. Thomas, 5136-610) fitted with a condenser collar. Freshly broken NaOH pellets (0.75 g) and 1.5 ml γ -butyrolactone (Sigma) were added to the bulb. The tear bulb was placed in a heating mantle, brought to a temperature of 220°C at the rate of $2^{\circ}\text{C}/\text{min}$ and then refluxed for an additional 23 hr period. The reaction was stopped by the addition of water. As the reaction mixture cooled it hardened to a white solid substance. The reaction mixture was dissolved slowly by adding several 2-5 ml aliquots of water, warming, and shaking the mixture on a vortex mixer, resulting in a 50 ml final volume.

The aqueous solution was next extracted two times with an equal volume of chloroform, adjusted to pH 2.5 with 6 N HCl, and extracted with diethyl ether three times. The ether extracts were combined and, along with the chloroform and aqueous fractions, were checked by TLC for complete extraction into the ether phase. The ether fraction was then dried over anhydrous sodium sulfate for 1 hr, filtered, and evaporated to near dryness in a Buchi rotovaporator. The residue was dissolved in 2 ml 50% 2-propanol (aqueous, v/v) and applied to a 2.5 x 48 cm column of Sephadex LH-20. The IBA was eluted with 50% 2-propanol and 6 ml fractions collected. IBA eluted at 276-294 ml as determined by TLC. The fractions containing IBA were combined and evaporated to dryness and 2 ml 2-propanol was added immediately. The yield was approximately 30% and the identity of the product was confirmed by GC-MS (see details below).

Determination of IBA and IAA levels in transformed carrot: Between 2-3 g of root tissue from carrots obtained from a California market were cut into small pieces and homogenized with imidazole buffer (35% 0.2 M imidazole, pH 7, 65% 2-propanol; 4 ml/g tissue) using a Polytron. The extract was transferred to a 16 ml centrifuge tube and the following standards were added: 100,000 dpm ^3H -IBA (a gift from Dr. Ephraim Epstein, Volcani Center, Israel), 100,000 dpm ^3H -IAA, 50 ng each of $^{13}\text{C}_6$ -IAA and ^{13}C -[ring 2]-IBA. The extract was allowed to equilibrate for 1 hr at 4°C . The sample was then centrifuged at $2000 \times g$ for 5 min and washed two times with additional extraction buffer. Extracts were combined and reduced to one third of the original volume, making sure that all propanol had been removed.

The extract was transferred quantitatively to a separatory funnel or Mixxor apparatus (Lida) and was partitioned three times against 1/3 volume diethyl ether. The ether phase was removed and discarded. The aqueous phase was then brought to pH 2.5 and partitioned two times against ether, saving the solvent phase. The sample was dried over 1/3 volume anhydrous sodium sulfate for 1 hr at 4°C and then filtered. The ether portions were combined, evaporated to dryness *in vacuo* and the residue dissolved in 60 μl methanol. Great care was taken to ensure that the residue was fully dissolved in the methanol. Water (60 μl) was used to rinse the flask, thus bringing the final concentration of methanol to 50%.

HPLC Analysis: Samples were further purified by reverse-phase HPLC using a Waters instrument (Waters 6000A pumps with 680 controller) fitted with a C_{18} , 3 μm Microsorb (Rainin) column, 4.6 mm ID x 100 mm, and an Upchurch guard column packed with Whatman Co: Pell ODS. The sample was purified by chromatography using a linear gradient of acetonitrile/1% aqueous acetic acid starting at 15% and going to 50% acetonitrile over 25 min. Under these conditions the retention volume of IAA was 7.8 ml and that of IBA was 13.5 ml. Samples containing the radioactive peaks were combined, keeping the IAA and IBA peaks separate. Each sample was evaporated *in vacuo* to dryness and 100 μl methanol was added immediately. The samples were methylated using ethereal diazomethane, dried under nitrogen, and dissolved in 20 μl ethyl acetate.

Analysis by GC-MS: GC-MS was performed using a Hewlett Packard 5890 GC equipped with a

15 m 0.21 mm i.d. J & W Scientific DB-1701 fused silica column, using helium as the carrier gas at a flow rate of 1 ml/min. IAA and IBA were analyzed separately. Mass spectrometry was performed on a Hewlett Packard 5971A mass selective detector. Selected ion monitoring of ions at m/z 130, 136, 189, and 195 for IAA and its $^{13}\text{C}_6$ -standard and m/z 130, 131, 217 and 218 for IBA and its $^{13}\text{C}_1$ -standard with a dwell time of 50 ms for each ion was used.

Carrot embryogenesis

Somatic embryogenesis involves the orderly progression of undifferentiated callus into organized embryonic stages (globular, heart and torpedo), resulting in mature embryos possessing the potential to become mature plants. Because there exists considerable evidence that both exogenous and endogenous auxins regulate embryo development, we are now investigating the concentration and metabolism of auxins during carrot somatic embryogenesis using stable isotope labeling techniques. The development of somatic embryos in culture closely parallels the development of zygotic embryos within immature seeds. However, carrot somatic embryogenesis offers two experimental advantages for hormone analysis: 1) somatic embryos can be isolated in large numbers sufficient for analytical methods and free of any non-embryogenic tissues, and 2) the somatic embryos can be separated by differential filtration on the basis of size into distinct fractions of each embryonic stage. The calli used for somatic embryogenesis are initiated and maintained in complete MS medium (Murashige and Skoog, 1962) with exogenous 2,4-dichlorophenoxyacetic acid (2,4-D). The removal of 2,4-D from the medium (accomplished by washing with medium devoid of 2,4-D) permits expression of the embryogenic potential to develop. The initiation of globular embryos is blocked by auxin analogs such as 2,4,6-trichlorophenoxyacetic acid and the transition from globular to torpedo embryos is disrupted by auxin transport inhibitors such as N-(1-naphthyl)phthalamic acid and 2,3,5-triiodobenzoic acid. Therefore, in order to further understand the process whereby the removal of the exogenous auxin-like regulator 2,4-D permits embryo development to proceed, it was important to determine the changes that occur in the levels of the endogenous IAA. It might be expected that the presence of the exogenous growth regulator would suppress endogenous auxin production, yet this is not what we discovered. The question we were still left with was: What is the relationship between 2,4-D and IAA metabolism in the induction of carrot somatic embryo formation? This question was addressed in a second series of experiments.

Both 2,4-D and IAA were quantitatively measured in calli and developing embryos at different stages using GC-MS-SIM after initial purification. IAA and 2,4-D were extracted from the carrot tissue using 65% 2-propanol/0.2 M imidazole buffer pH 7. Both [$^{13}\text{C}_6$] IAA and [$^{13}\text{C}_6$] 2,4-D (Cambridge Isotope Laboratories) were quantitatively added during extraction to serve as internal standards. After removing the cell debris by centrifugation and evaporating the organic phase *in vacuo*, the samples were divided into three fractions for analysis following selective base hydrolysis: 1) free IAA and free 2,4-D, 2) free plus ester conjugate IAA and free plus ester conjugate 2,4-D and 3) total IAA (free plus ester plus amide conjugates) and total 2,4-D. Fraction 1 was directly purified on a Fisher Prep Sep amino disposable column and by HPLC on a 4.6 X 125 mm 5l C_{18} (Partisil ODS-3, Whatman) column. The IAA was eluted within 15 min using 20% acetonitrile/1% acetic acid, and was followed by elution of the 2,4-D with 40% acetonitrile/1% acetic acid. The free plus ester conjugate fractions (Fraction 2) and the total fractions (Fraction 3) were purified in the same manner after either mild hydrolysis (1 hr, 1 N NaOH, 25 C) or harsh hydrolysis (3 hr, 7 N NaOH, 100 C) respectively, and desalting using a Fisher Prep Sep C_{18} disposable column. Individual samples of the IAA and 2,4-D fractions were methylated with ethereal diazomethane and analyzed by GC-MS-SIM as previously described by us. For 2,4-D analysis, ions at m/z 175 and 181 (unlabeled and labeled ions) and 234 and 240 (unlabeled and labeled molecular ions) were monitored.

The experiments show that in the presence of 2,4-D there are high levels of total IAA (ca. 600 ng/g f.w.), 80-90% of which is amide conjugate with the remainder consisting of equal amounts of free and esterified forms. Once the 2,4-D is removed, the amide conjugate level dropped dramatically as globular embryos formed, and was followed by a slow decline in the levels of all forms of IAA as embryogenesis proceeded. 2,4-D was also found to accumulate

primarily as amide conjugate(s) when 2,4-D was included in the medium. The levels of 2,4-D within the cells declined dramatically after transfer to 2,4-D free medium.

Because the IAA levels were extraordinarily elevated, it became important to investigate in the second series of experiments the biosynthesis and turnover of the IAA. The most direct way to monitor the IAA biosynthetic pathway is to measure the incorporation of precursors labeled with stable isotopes into IAA. In particular, the cells were grown in either D₂O (which labels IAA and its precursors early in the pathway), ¹⁵N indole (which is a key intermediate in the pathway) or D₅-L Trp (which is the putative amino acid precursor of IAA). Using GC-MS-SIM as described above, the incorporation of D₂O into IAA was determined to be the same in suspension cultures with 2,4-D and in embryogenic cultures without 2,4-D. Similarly, ¹⁵N indole labeled 25% of the IAA in the embryogenic culture and 21% of the IAA in the cultures proliferating in 2,4-D. D₅-L Trp, however, shows greater incorporation into IAA in the presence of 2,4-D in suspension cultures (35%) than in embryogenic cultures devoid of 2,4-D (<5%).

In order to interpret these results on IAA incorporation, it becomes necessary to assess the indole incorporation into Trp and the ability of D₅-L Trp to label the bulk Trp pool. Tryptophan was extracted using 65% 2-propanol/0.2 M imidazole buffer (pH 7) and D₅-L Trp was added as an internal standard. The sample was applied to a Dowex 50W-X2 200-400 (hydrogen form) column, washed with distilled water and then eluted with 2 N NH₄OH. The extract was then filtered through a "Bakerbond SPE" amino disposable column, evaporated to dryness and dehydrated by multiple azeotropic distillations using absolute ethanol and dichloromethane. The Trp was then derivatized in anhydrous methanol and acetic anhydride (Supelco) for 1 hr at 55 C. The residue was subsequently washed on a Fisher PrepSep disposable C₁₈ column and purified by HPLC using a Waters 4 lm NovaPak C₁₈ column 15 cm X 3.9 mm with 30% methanol/water. The appropriate fractions were then quantitatively analyzed using GC-MS-SIM. Gas chromatography was run essentially at the same conditions as described (Chen *et al.*, 1988) for IAA analysis except that the initial 140 C oven temperature (1 min) was followed by a ramp at 30 C/min up to 280 C. Ions monitored were at m/z 130 and 135 (unlabeled and labeled quinolinium ions) and 260 and 265 (unlabeled and labeled molecular ions). The results of the Trp experiments show that ¹⁵N indole is equally incorporated into Trp in suspension cultures with 2,4-D and embryogenic cultures, and that D₅-L-Trp equally labels the Trp pool under both experimental conditions.

The most interesting finding from the IAA and Trp incorporation studies is the differential incorporation of Trp into IAA depending on whether 2,4-D is present or not. This could be attributable to differences in uptake of the precursor into the different cultures, differences in compartmentation of either the precursor or the enzyme responsible for IAA biosynthesis or a change in the IAA biosynthetic pathway promoted by 2,4-D. Uptake was eliminated as a cause by measuring isotopic enrichment in the free Trp pool of the cells in both proliferating suspensions with 2,4-D and in embryogenic cultures without 2,4-D. The possibility of a change in compartmentation of Trp brought about by 2,4-D was minimized by using long-term rather than pulse chase type feeding (2 weeks in MS media + 10 mg/l D₅-L Trp +/- 2,4-D) but this question needed to be addressed directly.

Compartmentation was investigated by comparing the D₅-L Trp labeling of the cytoplasmic Trp pool and the plastid Trp pool. Conventional techniques for isolating plastids were not successful with the plastids in carrot suspension culture cells which are starch containing. However, using the GR buffer described for amyloplast isolation (1 mM NaP₂O₇, 50 mM HEPES, 330 mM sorbitol, 2 mM Na₂EDTA, 1 mM MgCl₂ and 1mM MnCl₂), plastids can be isolated by homogenizing cells in a Waring blender equipped with razor blades using three 5 sec pulses at low speed. The plastids are then obtained by filtration through Miracloth and a 43 lm screen, followed by differential centrifugation to separate the cellular debris (25 x g for 20 min) from the plastids which are then pelleted by centrifugation at 650 x g for 20 min.

The plastids isolated from cells that had been grown on D₅-Trp were analyzed for the percent labeling of the Trp pool by D₅-L-Trp and compared with the labeling of the total cellular Trp pool. A small differential labeling of the total cellular Trp pool and the plastid localized Trp pool was noted, but this difference was not changed in the presence versus the absence of 2,4-D in the media. This shows that 2,4-D was not affecting the compartmentation of Trp

and thus the differential incorporation of Trp into IAA is attributable to separate pathways for IAA biosynthesis.

***Petunia* Organogenesis**

The uptake and metabolism of exogenous tritium-labeled benzyl adenine was studied during the shoot induction period of petunia leaf explants in tissue culture. These experiments form the complement to our other work on auxin or auxin removal induction of organogenesis or embryogenesis. Transfer experiments with leaf explants (1 cm²) on MS media with 2.2 μ M BA show that 27% and 100% of the leaf explants are committed to shoot induction on days 6 and 10, respectively. To study BA uptake and metabolism, leaf explants were placed on media containing ³H-BA for 1,3,6 and 10 days. BA was taken up from the media on days 1-6; the rate of BA uptake increased to 1826 pmol/day at day 2 but was constant from day 2 to 6. BA metabolites were analyzed using HPLC and identity confirmed by mass spectrometry. Metabolites produced include: BA, BAdo, BA 7G, BA 9G, BAdoMP, BAdoDP, BAdoTP and three still unidentified compounds. BA was detected at day 1 but was not detectable from days 3-10. The ribotide metabolites decreased from days 1 to 10. For example, BAdoMP decreased from 193 to 31 pmol/g. Glucosylated compounds, BA 7G and BA 9G, increased continuously throughout the culture period. The concentration of BA 7G increased from 85 to 1055 pmol/g. These data suggest that particular stages of organogenesis are related to the changes in BA metabolites present in the tissue.

These studies were continued by the study of the uptake and metabolism of BA into two lines of *Petunia hybrida* differing in their shoot organogenic response. Leaf transfer experiments using shoot induction medium containing 4.4 μ M BA showed that leaf explants from line St40 required a shoot induction period of 6-10 days for commitment to shoot organogenesis; whereas leaf explants from petunia TLV1 required 12-28 days. The shoot induction period of St40 and the higher organogenic response was positively associated with a threefold higher absorption of BA from the medium, an increased BA ribotide metabolite pool, the presence of BA within the explant during the shoot induction period, and the production of 6-benzylamino-9-[O-glucopyranosyl-(1-3)-ribofuranosyl]-purine (metabolite C). However, the study of TLV1 leaf explants showed that neither BA nor metabolite C were required during the shoot induction period for eventual shoot development. The longer shoot induction period of TLV1 was associated with low BA uptake during 24 days, a decreasing ribotide metabolite pool, the absence of BA triphosphate and metabolite C throughout the study and the absence of BA within the explant during the shoot induction period. Differences in the shoot organogenic response of these related plant lines are thus associated with differences in exogenous cytokinin uptake and the subsequent metabolism of that hormone.

Auxin Biosynthesis

Lemna studies: An aspect of the work in the Beltsville laboratory has concerned the development of methodology for using stable isotope technology with which to then determine the *in vivo* pathway(s) for IAA biosynthesis (Cohen *et al.* 1990; Baldi *et al.* 1991). Differences in pool sizes, rates of turnover, and low incorporation of label make it difficult to analyze and interpret labeling patterns in classic studies of IAA biosynthesis. Use of stable isotope labeling allows a more complete picture to be developed of the kinetics of labeling and the molecular identity of labeled compounds. As an important foundation and test project we used normal and the IAA overproducing mutant, R₁, of *Lemna gibba* to examine the theory that D, not L, tryptophan was the closer precursor to IAA (Law, 1987). In these studies we used ¹⁵N-labeled D and L-tryptophan supplied to the plants and then analyzed the incorporation of ¹⁵N into IAA by mass spectrometry (Baldi *et al.*, 1991; see Appendix 1). These studies showed that L-tryptophan could be converted to IAA without labeling of the D-tryptophan pool and that labeled D-tryptophan, when given so as to yield an equal enrichment as was used in the L-tryptophan studies, was not an effective label. In addition, it was possible to make quantitative evaluations as to the *rate* of

labeling of the IAA pool from L-tryptophan. These considerations indicated that while D-tryptophan was ineffective, L-tryptophan was also converted at relatively slow rates. These results suggested to us that the widely held views about IAA biosynthesis from tryptophan needed a critical reevaluation.

Maize studies: In cooperative studies with Drs. Allen Wright and M. Gerald Neuffer (University of Missouri) we investigated the maize mutant *orange pericarp* (Wright *et al.*, 1991). This is a seedling lethal mutant which lacks tryptophan synthase B activity due to the mutation of two unlinked loci, *orp1* and *orp2*. The mutant can be rescued by treatment with tryptophan and accumulates indole and anthranilate. The orange color is due to indolic products which accumulate in the pericarp (Wright and Neuffer, 1989). We have used this mutant to test the hypothesis that tryptophan is the precursor to IAA. IAA levels in aseptically grown mutant seedlings (10 d) were determined by quantitative GS-MS and were found to be 50 times that of the normal seedlings. When mutant and normal seedlings were grown on media containing 30% $^2\text{H}_2\text{O}$, only the normal seedlings showed significant incorporation of deuterium into stable ring positions of tryptophan, while both the normal *and the mutant* showed substantial deuterium incorporation into stable positions on IAA. Similar experiments with ^{15}N -anthranilate feeding confirmed these results and established the fact that in mutant seedlings *de novo* IAA biosynthesis occurs without going through tryptophan. Together, these studies offer a fundamental modification of current thought as to the origin of this plant hormone.

The Volcani Center

Work in the Epstein laboratory at the Volcani Center has focused on the comparison of indolic auxin metabolism in tissues and species differing in their ability to form adventitious roots in response to auxin treatment. Several different species in which cultivars are known with differences in ease of rooting formed an important aspect of many of these experiments. Experiments were continued with mung bean as a standard for comparison and with cultivars of olive with different rooting ability (both these studies were used to obtain the preliminary data used in the initial grant proposal). More recently, these ideas were tested further by studies on a number of diverse species, as detailed below.

Mung bean studies: It has been well established that IBA induces adventitious root formation in mung bean hypocotyls and that *de novo* transcription starts following IBA induction. The resulting mRNA's are believed to be translated into proteins that are responsible for the altered developmental program resulting in root formation. In order to begin to understand this developmental program better it is important to look at genes regulated by the auxin involved and understand the promoters and their interaction with the signal molecule. We found that endo-1,4-beta-glucanase (EG), coded in bean by a small family of genes, are closely related to the activity of ethylene and auxin. In particular, the EG with an isoelectric point at 4.5 is closely related to the promotive effects of auxin. Hypocotyls treated with IBA formed roots along the hypocotyl and total EG activity reached a peak between 12-24 h after treatment, however other enzymes such as protease and amylase were not affected by IBA treatment. We showed that the acidic EG activity was positively correlated to IBA activity and that the basic EG activity correlated with ethylene activity and that the activity of both enzymes changed over the course of treatment and only the basic activity was affected by ethylene inhibitor application. Using tomato and avocado EG probes we have obtained putative clones for bean EG from cDNA libraries we constructed.

Petunia cell suspension cultures: The uptake and metabolism of IAA and IBA were studied in a model system using suspension cultures of *Petunia hybrida*. The initial uptake of ^3H -IBA was higher than that of ^3H -IAA and after 10 min of incubation with labeled IBA and IAA, 4.6 pMol vs 0.35 pMol (39% vs 12% of total applied radioactivity) of IBA and IAA, respectively, were found in the cell extracts. The uptake of IBA reached a plateau of 6.0 pMol (62%) after 2 h while that of IAA increased continuously up to 1.5 pMol (46%) after 24 h. Following the

addition of 40 μ M of unlabeled auxin more IBA was initially than IAA (39% vs 12%) but the level almost equalized after 24 h of incubation when IBA uptake reached 890 nMol (55%) and IAA reached 840 nMol (46%). IBA was metabolized very rapidly by *Petunia* cell suspensions to new products. HPLC of the cell extracts demonstrated a new metabolite after only 2 min of incubation, and after 30 min 60% of the radioactivity was in the new metabolite vs 10% in the IBA. The new compound was resolved by autofluorography into two metabolites but after 24 h only a single compound persisted. The metabolites were tentatively identified as IBA aspartate and IBA glucose. In the medium IBA disappeared rapidly and after 24 h most of the radioactivity was present as IBA aspartate. IAA was also converted into two products, however both products were still present after 24 h. IAA metabolism proceeded at a slower rate and while free IBA disappeared after only 0.5 h, free IAA was still present after an hour incubation. We showed that only intact cells were able to metabolize IBA and that the reaction was inhibited by low temperature and anaerobic conditions. Thus these reactions appear to involve reactions that occur at or near the cell surface.

Citrus leaf: The transport and metabolism of radiolabeled IBA was studied using midrib sections of Cleopatra mandarin (*Citrus reticulata* Blanco) and compared to that of IAA. Exogenous IBA was metabolized by the midribs to a polar ester conjugate. Ethylene pretreatment of the midribs reduced their capacity to metabolize IBA by ca. 70% as compared to air controls. IBA transport capacity in the leaf midribs was ca. two times greater in the basipetal direction than the acropetal. The basipetal transport capacity of ^3H -IBA was lower than that of ^{14}C -IAA (ca. 24% and 39% of the uptake, respectively). While ethylene treatment reduced basipetal transport of IAA by ca. 70% it did not affect the transport of IBA. Most of the transported label was found as free IBA, but the reduction of IBA conjugation by ethylene treatment did not effect the transport capacity.

Sweet cherry: The transport and metabolism of IBA were studied in easy-to-root (ER) and difficult-to-root (DR) cultivars of sweet cherry (*Prunus avium* L.) in an aseptic system. Plantlets of both cultivars (ER=Rainer; DR=40-E-50) were grown on rooting agar medium supplemented with ^3H -IBA. Most of the radioactivity (80%) was found in the bases of the shoots of both cultivars and very little was transported to the shoots and leaves. ER plantlets absorbed more IBA for a longer period than DR plantlets. Both cultivars metabolized IBA rapidly to an ester with glucose. After one day, no free IBA could be detected in the extracts of the DR cultivar. The ER cultivar, on the other hand, metabolized the conjugate slower and after one day an intense spot of IBA was still visible on the autofluorogram. Free IBA disappeared after the second day, but re-appeared after 4 days and disappeared again after 7 days. We postulated that the ER cultivar, as opposed to the DR cultivar, has an increased capacity to hydrolyze the ester conjugate at the appropriate time to release IBA at the appropriate time to promote root initiation. Conjugation was inhibited in the DR cuttings with 1,6-dihydroxyacetophenone and this increased significantly the percentage of rooting.

Protea: IBA greatly enhanced the rooting of an early-flowering variety of protea, *Leucadendron discolor*, but had very little effect on the late-flowering variety. IBA transport and metabolism were studied in both varieties after incubating of the cuttings in ^3H -IBA. More of the radiolabeled IBA was transported to the leaves of the ER variety than the DR variety (35%-45% and 10%, respectively). IBA was metabolized rapidly by cuttings of both varieties and after 24 h most of the label was in the new metabolite. However, free IBA (about 10%) was present in the cuttings during the whole period up to the time of root emergence (4 weeks). More free IBA was accumulated in the base of ER cuttings, while in the DR cuttings IBA was conjugated to glucose. It is probable that IBA-glucose serves as a source of IBA and the difference between the varieties is a consequence of the free IBA which is released, transported and accumulated at the site of root formation.